MCI DEC 2017

1. Which of the following is a Selective estrogen receptor modulator -
    a. Fulvestrant
    b. Flutamide
    c. Tamoxifen
    d. Clomiphine citrate

Explanation :

a. Fulvestrant – selective estrogen receptor down regulators (or) pure estrogen antagonists
b. Flutamide - selective antagonist of androgen receptor
c. Tamoxifen – selective estrogen receptor modulator. Used as a major indication for treatment of breast cancer. Has beneficial effect on bone and lipid profile but increase the risk of endometrial cancer and thromboembolism.
d. Clomiphine citrate – binds to both ER- alpha and ER – beta and acts as a pure estrogen antagonists.

2. which of the following Stone is insensitive to alkaline PH of urine?
    a. Xanthine stones
    b. Cystine stones
    c. Struvite stones
    d. Calcium stones

Explanation:

a. Xanthine stones – due to deficiency of xanthine oxidase leading to xanthinuria. May be either radio opaque or radio lucent .
b. Cystine stones – Due to the defect in renal absorption of amino acid cystine leading to cystinuria. Radio opaque due to the presence of sulphur. They are hexagonal shaped or benzene like crystals.
c. Struvite stones – aka triple phosphate stones. Formed mainly after infection by proteus, which causes alkaline urine leads to precipitation of phosphate crystals. Drugs like acetazolamide can also cause struvite stones.
d. Calcium stones - most common type of stone is calcium oxalate stones. They are radio opaque.

3 Which of the following is Ocp induced liver malignancy ?
    a. Hepatic adenoma
    b. Hepatocellular carcinoma
    c. Hemangioma
    d. Focal nodular hyperplasia

www.medicalcoupons.in

Explanation :

a. Hepatic adenoma: Most common symptomatic benign liver tumour. Most common associated factors are ocp and steroids. High risk of rupture and malignant transformation.

4. Which of the following methods of contraception is used within 1 st 6 months of lactation?
   a. Combined Ocps
   b. Copper T- A380
   c. Progesterone only pills
   d. Condoms

Explanation:

a. Combined ocps – are contraindicated during lacation .
b. Copper T-A380 – can also be used during lactation but most common complication is vaginal bleeding .
c. Progesterone only pills - used mainly within 72 hrs of intercourse to reduce the risk of pregnancy and is considered as safe during lactation.
d. Condoms – not much effective because of its high failure rate 2-14 per HWY.

5. Prosopagnosia means ?
   a. Inability to write
   b. Inability to calculate
   c. Inability to identify face
   d. Inability to speak

Explanation :

a. Inability to write – agraphia
b. Inability to calculate – acalculia
c. Inability to identify faces – prosopagnosia
d. Inability to speak - aphasia

6. Layers of spinal anaesthesia ?
   a. Skin>subcutaneous>supraspinatus ligament>interapinatus ligament> ligamentum flavum
   b. Skin> subcutaneous>supraspinatus ligament >ligamentum flavum>interspinous ligament
   c. Skin>subcutaneous>interspinous ligament> supraspinatus ligament>ligamentum flavum
   d. Skin >supraspinatus ligament >subcutaneous >interspinous ligament >ligamentum flavum

www.medicalcoupons.in

7. Which of the following is the correct order of endometrial blood supply?
    a. **Arcuate>radial>basal>spiral**
    b. Radial>arcuate >spiral > basal
    c. Radial >basal > arcuate > spiral
    d. Arcuate > radial > spiral > basal
8. SIADH is caused by ?
    a. Head trauma
    b. Tb meningitis
    c. Small cell carcinoma of lung
    d. **All of the above**

Explanation :

SIADH : syndrome of inappropriate ADH.

CAUSES: Tumors like- head and neck cancer

Small cell cancer of lung

Bronchial carcinoma

Carcinoid tumour

Head trauma

Drugs like – carbamazepine

Clofibrate

Chlorpropamide

Vincristine

Infections like – pneumonia

TB

Meningitis

SIADH is characterized by gain of water and hence dilutional hyponatremia is seen. Rapid correction of hyponatremia produces “central pontine myelinosis”.

Rx: doc is – demiclocycline

9. RDA of calcium in elderly women?
    a. 600
    b. **1200**
    c. 1800
    d. 2400

www.medicalcoupons.in

Explanation :

Elderly women – 1200mg

Pregnancy – 1300 - 1500mg

10. Calculate the volume of fluid to be replaced in a burn patient of TBS 40% of 50 kg within 1 st 8 hrs?
    - **a. 4 L**
    - b. 8 L
    - c. 12 L
    - d. 16 L

Explanation :

Volume of fluid to be given for burns patient is calculated by parkland formula.

Parkland formula= TBSA% * weight (kg) * 4

= 40% * 50*4

= 8L

In 1st 8 hrs- 50% of fluid should be given i.e., 4L

In next 16 hrs – remaining 50% of fluid should be given .

11. Sun burst appearance in osteosarcoma is due to?
    - a. Abnormal calcium deposits
    - **b. Periosteal reaction**
    - c. Bony destruction
    - d. Highly vascular

Explanation :

Osteosarcoma:

2nd most common malignant bone tumour, arise from metaphysis of lower end of femur. Gross appearance shows mutton leg appearance . periosteal reaction is present.

X ray findings - sun ray appearance

Codmans triangle

Alkaline phosphatase is highly increased.

Rx pre- operative chemotherphy ,limb saving surgery , post – operative chemotherphy .

www.medicalcoupons.in

12. Motor molecule are all of the following except?
    a. Kinesin
    b. Dyenin
    c. Myosin
    d. Actin
13. P value in null hypothesis?
    a. 0.005 %
    b. 0.05%
    c. 0.5%
    d. 5%

Explanation :

Test of significance:

- state the null hypothesis (ie, state that both groups are same)
- state the level of significance (alpha - 0.05)
- choose the appropriate statistical test
- find "P" value
- if "P" value is <0.05 , reject null hypothesis ie, both groups are different .
- if "P" value is >0.05 ,accept null hypothesis ie, both groups are same.

14. Framingham heart study is an example of?
    a. Retrospective cohort
    b. Prospective cohort
    c. Case control study
    d. Cross sectional study

Explanation:

Framingham heart study was initiated to study the relationship between the number of risk factors and subsequent development of cardiovascular disease. Is an example of cohort study.

15. In Which of the following insentives to mother is 1000Rs and Asha 400rs?
    a. Rural area ,low performing states
    b. Urban area , low performing states
    c. Rural area , high performing states
    d. Urban area , high performing states

www.medicalcoupons.in

Explanation : RURAL URBAN

	Mother	Asha	Mother	Asha
Low performing states	1400	600	1000	400
High performing states	700	600	600	400

16. Which of the following is regarding the work of Asha ?
    a. Conduct delivery
    b. Provides resource for anganwadi workers
    c. Builds house hold toilets
    d. For immunization of children

Explanation :

ASHA: ACREDITED SOCIAL HEALTH ACTIVIST

- Asha must be a resident of a village preferably the age group of 25-45.
- 1 Asha works for 1000 population.
- Minimum education required for Asha is- 8 th pass
- Training of Asha is done by ANM and AWW for minimum duration of – 23 days .
- Impact indicators of Asha:- infant mortality rate
- Child malnutrition
- Number of TB/leprosy case detection.
- Escort services of RCH
- Helps in construction of household toilets.

17. Which of the following cancer is highest reported to death worldwide ?
    a. Ovarian ca
    b. Breast ca
    c. Stomach ca
    d. Lung ca

www.medicalcoupons.in

Explanation :

The most common causes of cancer death are cancers of:

- Lung (1.69 million deaths)
- Liver (788 000 deaths)
- Colorectal (774 000 deaths)
- Stomach (754 000 deaths)
- Breast (571 000 deaths)

Reference : http://www.who.int/mediacentre/factsheets/fs297/en/

18. Vaccine contraindicated in pregnancy?
    a. **Rubella**
    b. Yellow fever
    c. BCG
    d. Hep b

Explanation :

All live vaccines are contraindicated in pregnancy except, yellow fever vaccine and OPV. Women are advised not to receive the rubella vaccine during pregnancy as a safety precaution based on the theoretical possibility of a live vaccine causing disease, in this case, 'congenital rubella syndrome'.

19. Black colour triage indicates ?
    a. **Unsalvageable**
    b. Highly prioritized
    c. Can wait for some time
    d. Ambulatory care

Explanation : red – highly prioritized

Yellow – can wait for some time

Green – ambulatory care

Black - moribund/ dead

20. Which of the following is the correct order of Chain of survival ?
    a. Early access> early chest compression>early airway management >early defibrillator>early hospital admission
    b. Early chest compression> early hospital admission >early access >early airway management > early defibrillator
    c. **Early access>early airway management>early chest compression > early defibrillator > early hospital admission**
    d. Early access>early defibrillator > early hospital admission > early airway management > early chest compression

www.medicalcoupons.in

21. Which of the following hormone helps in the regulation of 1,25 dihydroxycholecalciferol ?
    a. Renin
    b. Angiotensin
    c. Parathormone
    d. Vit D

Explanation :

Hormones regulating calcium homeostas

Hormone	Blood Ca	Main actions
PTH Parathyroid hormone	↑	stimulates bone demineralization (o clas s) stimulates renal Ca resorp on (indirect) stimulates syn esis of c citriol in kidneys
Calcitonin	↓	inhibits renal Ca resorpti stimulates bone m alizati
Calcitriol	↑	stimula s intes al Ca re rption stimulate e act of H on kidney

1. Maintaining constant Ca^{2+} plasm centra n (PTH, calcitonin)
exchange between bones and ECF, rine tion of Ca^{2+}, quick regulation
2. Maintaining constant calcium ont nt in body (calcitriol)
control of calcium intake (intestine), s w re ulation

22. Ranolazine is not con raind ated in giving with which of the following drugs?
    a. Ketoconazo
    b. Diltiaz em
    c. Morph n
    d. Clo api

Explanation:

Dose of ran lazine should be limited to 500 mg twice daily in patients on MODERATE C YP3A INHIBITORS ,including diltiazem , verapamil , erythromycin , flucona ole .


Do n t us ranolazine with STRONG CYP3A INHIBITORS, including ketoconazole, itraco azole, clarithromycin etc.

23. Glycemic index of glucose?
    a. 0.5
    b. 1
    c. 1.5
    d. 2

www.medicalcoupons.in

24. Corocoid process of scapula is a example of?
    a. Pressure epiphysis
    b. Traction epiphysis
    c. Atavistic epiphysis
    d. Aberrant epiphysis

Explanation :



25. Imbecile?
    a. Less than 20
    b. 25-34
    c. 25-49
    d. 50-70

Explanation : less than 20 – profound

0 -24 - idiot

25 – 49 – imbecile

50 – 70 – moron

26. Keratic precipitate is seen in which layer ?
    a. Epithelium of cornea
    b. Decemets membrane
    c. Stomach
    d. Endothelium of cornea

www.medicalcoupons.in

Explanation:

Keratic precipitate (KP) is an inflammatory cellular deposit seen on corneal endothelium. Acute KPs are white and round in shape whereas old KPs are faded and irregular in shape. Mutton-fat KPs are large in shape and are greasy white in color and are formed from macrophages and epithelioid cell. They are indicative of inflammatory disease. Mutton fat Kps are due to granulomatous iridocyclitis. Another variant called red KPs may be seen in hemorrhagic uveitis.

27. All of the following shows Increased anion gap are all except?
    a. Aspirin poisoning
    b. Uremia
    c. Diarrhoea
    d. Diabetic acidosis

Explanation : diarrhoea is eg of normal anion gap.

28. Respiratory papillomatosis is caused by?
    a. HPV
    b. EBV
    c. H. Influenza
    d. Klebsiella

Explanation :

Recurrent respiratory papillomatosis (RRP), which is caused exclusively by human papilloma virus (HPV), is a rare condition characterized by recurrent growth of benign papillomata in the respiratory tract. The papillomata can occur anywhere in the aerodigestive tract but most frequently in the larynx, affecting both children and adults. The management of this entity remains still challenging since no specific definitive treatment exists.

29. Positive swartz sign is seen in ?
    a. Meniere's disease.
    b. Active otosclerosis
    c. Acoustic neuroma
    d. Malignant otitis externa

Explanation:

Otosclerosis is an abnormal bone growth in the middle ear that causes hearing loss. Swartz sign is a pink reflex , seen on the promontory through intact tympanic membrane ,in the area of oval window. It indicates active focus with increased vascularity. Otosclerosis usually occurs during pregnancy. Management for active otosclerosis is surgery and inactive otosclerosis is sodium fluoride theraphy.

www.medicalcoupons.in

30. which of the following complement deficiency leads to Meningococcal meningitis?
    a. C3b
    b. C1
    c. C5-C9 [MAC]
    d. C5a

Explanation :

Recurrent meningococcal meningitis is caused by neisseria infections due to deficiency of MAC complex (c5-9).

C1 esterase deficiency leads to hereditary angioedema.

C3b – is a opsonin ,helps in opsonisation of the pathogens.

C5a – chemokine.

31. In which of the following cattle tract appearance is seen?
    a. ARMD (age related macular degeneration)
    b. Diabetic retinopathy
    c. Retinal detachment
    d. CRAO

Explanation :

Atherosclerosis related thrombosis at the level of lamina cribrosa is most common cause (80%) followed by embolization from heart and carotid artery (20%) producing a sudden painless loss of vision.

C/F : sudden painless loss of vision ,cherry red spot at fovea, segmentation of blood in blood vessels producing cattle tract sign .

Rx : ocular massage is most effective method

Decrease IOP by giving manitol

Inhalation of carbogen (95% of O2 and 5% of CO2).

32. Treatment for nasopharyngeal carcinoma?
    a. Surgery
    b. Chemotheraphy
    c. Radiotheraphy
    d. Both chemo and radiotheraphy

Explanation :

Mc tumour of head and neck which give rise to secondaries with occult primary . Mc site is fossa of Rossenmuller . Mc type is non keratinizing undifferentiated carcinoma followed by keratinazing carcinoma. Mc manifestation is upper neck swelling usually unilateral due to cervical lymphadenopathy . earliest lymphnode to be involved is

retropharyngeal lymphnode. Mc cranial nerve palsy in nasopharyngeal carcinoma is 5 th followed by 6 th and also 9 th and 10 th CN. Treatment of choice is irradiation (external beam radiotheraphy) as it is highly radiosensitive. In stage 1 and 2 only radiotherapy is done and in 3 rd and 4 th stage chemoradiation is the treatement.

33. At 24 weeks of gestation the fundus height is at the level of ?
    a. Pubic symphysis
    b. Umbilicus
    c. Xiphoid sternum
    d. Manubrium of sternum

Explanation:

At 12 weeks fundus is above pubic symphysis

At 24 weeks fundus is at the level of umbilicus

At 36 weeks fundus is at the level of xiphi sternum

At 38 weeks fundus is just below the level of xiphi sternum

34. All of the following is true regarding TOF EXCEPT?
    a. Ventricular septal defect
    b. Normal position of aorta
    c. Pulmonary stenosis
    d. Right ventricular hypertrophy

Explanation:

- Tetralogy of fallot is a cyanotic congenital heart defect.
- A hole in the wall between your heart's main pumping chambers (ventricular septal defect or VSD).
- A valve between your heart and lungs that is too narrow (pulmonary stenosis or PS)
- A right heart chamber with walls that are too thick (right ventricular hypertrophy)
- A major blood vessel (aorta) that is misplaced or moved (overriding aorta).

35. All of the following are involved in triple arthodesis except?
    a. Talotibular
    b. Subtalar joint
    c. Calcaneocuboid joint
    d. Talonavicular joint

www.medicalcoupons.in

Explanation:

Triple arthrodesis is a surgical procedure whose purpose is to relieve pain in the rear part of the foot, improve stability of the foot, and in some cases correct deformity of the foot, by fusing of the three main joints of the hindfoot: the subtalar joint, calcaneocuboid joint and the talonavicular joint.

36. Spontaneous or post trauma watery nasal discharge is due to?

a. Allergic rhinitis
b. CSF rhinorrhea
c. atrophic rhinitis
d. Rhinosporidiosis

Explanation:

It is a flow of CSF from nose (due to CSF leakage from sub arachnoid space into nasal cavity). Usual sites of CSF leak are from cribriform plate >frontal sinus > floor of the anterior cranial fossa. Mc cause of CSF rhinorrhoea is iatrogenic trauma during surgery.

C/F: unilateral clear watery discharge dripping on looking down, which increases on coughing , sneezing or exertion.

37. Optic nerve diameter of adult is?

a. O.5 mm
b. 1 mm
c. 1.5 mm
d. 2 mm

38. Which of the following is the carbonic anhydrase inhibitor is anti glaucogenic?

a. Dorsalamide
b. Lantoprost
c. Brimonidine
d. Dipivefrine

Explanation:

Dorsalamide is a carbonic anhydrase inhibitor which is used in the management of glaucoma to decrease the IOP by decreasing the production of aqueous humour. Should not be given to patients who are allergic to sulphur.

www.medicalcoupons.in

39. 1 st trimester miscarriage is most commonly due to?
    - **a. Chromosomal abnormalaities**
    - b. Incompetent cervix
    - c. Fibroids
    - d. Infections

Explanation :

The most common cause of recurrent 1 st trimester abortion is chromosomal abnormalities of which mc is balanced translocation in mother. Mc cause of recurrent 2 nd trimester abortion is incompetent cervix. Infections can't cause recurrent abortions.

40. Quintan fever is caused by?
    - a. Tularaemia
    - b. Coxiella burnetti
    - c. H.ducreyi
    - **d. Bartonella**

Explanation:

Trench fever/quintan fever is a clinical syndrome caused by infection with Bartonella Quintana. The disease is classically a 5-day fever. The onset of symptoms is sudden with high fever, severe headache, back pain and leg pain and a fleeting rash. Recovery takes a month or more. Relapses are common.

41. Most common site of carcinoma esophagus?
    - a. Upper 1/3
    - **b. Middle 1/3**
    - c. Lower 1/3
    - d. Cricopharyx.

Explanation :

Mc site of esophageal carcinoma in india – middle 1/3 rd

Mc site of esophageal carcinoma in worldwide – lower 1/3 rd

Mc type – squamous cell carcinoma

42. Recurrent laryngeal nerve doesn't supply which of the following laryngeal muscles?
    - a. Cricoarytenoid
    - b. Thyroarytenoid
    - **c. Cricothyroid**
    - d. Interarytenoids

www.medicalcoupons.in

Explanation :

All the laryngeal muscles are supplied my recurrent laryngeal nerve except cricothyroid which is supplied by external laryngeal nerve , branch of superior laryngeal nerve .

43. A boy with fever with chills , Swollen tonsil and it is pushed medially is due to which of the following?
    - **a. Quinsy**
    - b. Lushcka tonsil
    - c. Parapharyngeal abcess
    - d. Retropharyngeal abcess

Explanation :

Quinsy/peritonsillar abcess is the collection of pus between the fibrous capsule of the tonsil. Mc occurs as a complication of acute tonsillitis and generally unilateral. Caused by GABHS /mixed flora. Managed by i.v fluids ,antibiotics and analgesics. If not responding to antibiotics, then interval tonsillectomy is done after 6 weeks of quinsy.

44. Karyotype of down syndrome is ?
    - a. Trisomy 13
    - b. Trisomy 18
    - **c. Trisomy 21**
    - d. Trisomy 22

45. From which of the following parafollicular “c” cells are derived from ?
    - **a. Ultimobranchial body**
    - b. Neuroectoderm
    - c. Splanchnic Mesoderm
    - d. pharyngeal arches

Explanation :

Parafollicular “c” cellas are derived from neural crest > 4 th ventral endodermal pouch.

www.medicalcoupons.in

46. Which of the following organ undergoes liquefactive necrosis?
    a. Cerebrum
    b. Kidney
    c. Small intestine
    d. Pancreas

Explanation :

Brain – undergoes liqefactive necrosis due to ischemia /abscess formation in brain .

All other organs of the body undergo coagulative necrosis eg.kidney ,small intestine etc.due to coagulation of proteins.

Pancreas and breast are eg of fat necrosis.

47. Launge hauman classification is for ?
    a. Shoulder joint
    b. Hip joint
    c. Knee joint
    d. Ankle joint

Explanation :

The **Lauge-Hansen classification** system is used for the classification of the ankle injuries based on injury mechanisms which have predictable patterns and imaging findings. Along with the Weber classification, these systems are useful tools for describing and classifying ankle injuries.

Classification

The Lauge-Hansen system uses two-word descriptors:

- the first word describes the position of the foot at the time of injury (i.e. supination or pronation)
- the second word describes the deforming force direction (i.e. abduction, adduction or external rotation)

48. Which of the following is the most cochlear toxic drug ?

    a. kanamycin
    b. gentimicin,
    c. amikacin,
    d. streptomycin

www.medicalcoupons.in

Explanation :

Neomycin has the greatest cochleotoxic effect of all antibiotics. **Kanamycin** are close to neomycin in cochleotoxic potential and are capable of causing profound, permanent hearing loss while sparing balance.
Streptomycin tends to cause more damage to the vestibular portion than to the auditory portion of the inner ear.


49. Creatinine is formed from which of the following amino acids ?
    a. Aspartate, glycine,lucine
    b. Alanine ,glutamate ,methionine
    c. Aspartate ,lycine , glutamate
    d. Arginine ,glycine, methionine

50. Wernike korsakoff is due to the deficiency of?
    a. Vit b2
    b. Vit b12
    c. Vit b1
    d. Vit –b6

Explanation :

Wernicke encephalopathy and korsakoff psychosis is due to deficiency of vit B1 thymine usually seen in alcoholics.


51. Identify the structure shown in arrow.



www.medicalcoupons.in

a. Asterion
b. Pterion
c. Inion
d. Glabella

Explanation :



52. GCS score for dead patient is?
a. 0
b. 3
c. 5
d. 15

Explanation :

GCS : max score is 15 and minimum score is 3 . GCS for a dead patient is 3.

Glasgow Coma Scale

Best eye response (E)	Spontaneous--open with blinking at baseline	4
	Opens to verbal command, speech, or shout	3
	Opens to pain, not applied to face	2
	None	1
Best verbal response (V)	Oriented	5
	Confused conversation, but able to answer questions	4
	Inappropriate responses, words discernible	3
	Incomprehensible speech	2
	None	1
Best motor response (M)	Obeys commands for movement	6
	Purposeful movement to painful stimulus	5
	Withdraws from pain	4
	Abnormal (spastic) flexion, decorticate posture	3
	Extensor (rigid) response, decerebrate posture	2
	None	1

www.medicalcoupons.in

53. Best to evaluate adequate fluid resuscitation in a burn patient?

a. Urine output
b. Thirst
c. Fall in Blood pressure
d. Shock

Explanation :

The key for monitoring fluid resuscitation in burns patients urine output. It should be between 0.5 and 1 ml/kg body weight per hour. If the urine output is below this ,the infusion rate should be increased by 50%.

54. 21 teeth present at the age of?

a. 5
b. 6
c. 10
d. 12

Explanation :

The complete set (ie,20) primary teeth have to come in by the time the child is 2 1/2 to 3 yrs old. Rule of thumb is that ~ 4 teeth will erupt for every 6 months of life. Lower teeth erupt before upper teeth with teeth in both jaws usually erupting in pairs. Primary teeth are smaller and whiter in color than permanent teeth that will follow. After age 4 ,the jaw and facial bones of child begin to grow and creating the space between the primary teeth. Between the age of 6 and 12, a mixture of booth primary teeth and permanent teeth reside in mouth .

55. Another Term of Lesbianism ?

a. Fetischism
b. Eonism
c. Urolagnia
d. Tribadism

Explanation :

Lesbianism /tribadism – gratification of sexual desire of a woman by another women.

Urolagnia – sexual excitement by sight or odor of urine or faeces

Eonism /transvestism – pleasure of wearing clothes of opposite sex.

Fetischism – sexual gratification by article of opposite sex.

www.medicalcoupons.in

56. Movement of superior rectus muscle?
    a. **Elevation , intortion, adduction**
    b. Depression, extortion, adduction
    c. Elevation , extortion, abduction
    d. Depression , intortion, abduction

Explanation:

Muscles	Function		
	Primary	secondary	tertiary
Lateral rectus	Abduction	-	-
Inferior rectus	Adduction	-	-
Inferior rectus	Depression	Extortion	Adduction
Superior rectus	Elevation	Intortion	Adduction
Inferior oblique	Extortion	Elevation	Abduction
Superior oblique	Intortion	Depression	Abduction

57. A Man with bulla on face and neck, flaccid bulla, spreading and nikolsky sign positive is seen in
    a. Bullous pemphigoid
    b. Pemphigus vulgaris
    c. Pemphigus violaceous
    d. Dermatitis herpetiformis

www.medicalcoupons.in

Explanation :

Disease	Biopsy features	DIF
Bullous pemphigoid	Subepidermal oedema/blister. Mixed perivascular infiltrate with eosinophils	Linear IgG, C3 in BMZ (lamina lucida).
Dermatitis herpetiformis	Subepidermal vesicles. Mixed papillary infiltrate with eosinophils.	Granular IgA in tips of papillae.
Pemphigus vulgaris	Suprabasal acantholytic vesicle (tombstone pattern). Mixed perivascular infiltrate with eosinophils. Bulla spread and Nikolsky sign is positive.	IgG within suprabasal intercellular spaces.
Pemphigus foliaceus	Subcorneal acantholytic vesicle (tombstone pattern). Mixed perivascular infiltrate with eosinophils.	IgG within upper epidermal intercellular spaces.

58. waxy hyaline membrane deposits in alveoli is seen in?

a. ARDS
b. Interstitial lung disease
c. Asthma
d. Emphysema

Explanation :

Acute Respiratory distress syndrome (ARDS) of the newborn is an acute lung disease caused by surfactant deficiency, which leads to alveolar collapse and noncompliant lungs. Previously known as hyaline membrane disease, this condition is primarily seen in premature infants younger than 32 weeks' gestation. RDS has improved with the increased use of antenatal steroids to improve pulmonary maturity, early postnatal surfactant therapy to replace surfactant deficiency.

www.medicalcoupons.in

59. Vitamin k undergoes -?
    a. Gamma decarboxylation of arginine
    b. alpha decarboxylation of arginine
    c. **Gamma carboxylation of glutamate**
    d. Alpha carboxylation of glutamate

Explanation :

Vitamin – K : is the coenzyme for gamma carboxylation of glutamate in postsynthetic modification of calcium binding proteins. Vitamin K is also important in synthesis of bone calcium binding proteins. Menadione is water soluble form of Vit – k.

60. Which of the following drug is the antidote for methanol poisoning ?
    a. Flumazenil
    b. Fomepizole
    c. Pralidoxime
    d. Protamine sulphate

Explanation :

Fomepizole is used as an antidote in confirmed or suspected methanol or ethylene glycol poisoning. Fomepizole is a competitive inhibitor of alcohol dehydrogenase, the enzyme that catalyzes the initial steps in the metabolism of ethylene glycol and methanol to their toxic metabolites.

61. GFR is best measured by which of the following substances?
    a. Inulin
    b. Paraaminohippuric acid
    c. Insulin
    d. ADH

Explanation :

Kidney function is determined by measuring glomerular filtration rate (GFR)--the volume of plasma that the kidneys filter through the glomeruli per unit time. The "gold standard" for measuring GFR is through the use of inulin. Para aminohippuric acid is used to measure the renal plasma flow.

62. Which of the following is most common cause for oral cancer ?
    a. Alcohol
    b. Betel nut chewing
    c. Sun exposure
    d. Infections like EBV,HPV etc.

www.medicalcoupons.in

Explanation :

Most cases of oral cancer (~90%) in south east asia are linked to tobacco chewing and tobacco smoking. In India mc form of tobacco chewing is betel liquid which usually consist of betel leaf ,areca nut , lime and tobacco.

63. In which of the following route fentanyl should not be given ?



a. Intramuscular
b. Intravenous
c. Transdermal
d. Intrathecal

Explanation :

Fentanyl can be given as slow iv or im , it can also be given intrathecal administration.

64. Site of gluconeogenesis is ?

a. Muscle
b. Liver
c. Kidney
d. Intestine

Explanation:

Gluconeogenesis is the synthesis of glucose from non carbohydrate compounds. It takes place in liver and kidney . The liver accounts for 90% of gluconeogenesis in the body and the remaining 10% occurs in kidney and other tissues of the body. It takes place in both cytoplasm and mitochondria of liver.

65. Hand foot mouth disease is caused by ?

a. HINI
b. Adenovirus
c. Rotavirus
d. Coxsackie virus

www.medicalcoupons.in

Explanation:

Coxsackie virus is single strand + ve sense RNA virus ,which spread by feco oral route. HFMD, caused by coxsackieviruses, usually causes fever, malaise, rash, and small blisters that ulcerate. The most frequent locations for the blisters/ulcers are on the palms of the hand, soles of the feet, and in the mouth. There is no specific treatment or vaccine available for coxsackievirus infections.

66. Period of viability of the fetus is?
    - **a. 210**
    - b. 220
    - c. 230
    - d. 240

Explanation :

Previously ,28 weeks of geststion is considered as a period of viability ie,210 days. Now it has been changed to 22 weeks to consider fetus as viable. The best option here is 210 days.

67. Cephalic index of 75- 80 represents?
    - a. Aryans
    - b. Negroes
    - **c. Europeans**
    - d. Mangolians

Explanation :

Types of skull	Cephalic index	Race
Dolico cephalic	70 - 75	Aryans, Negroes
Mesaticephalic	75 - 80	Europeans , Chinese
Brachycephalic	80 - 85	Mongols

68. Pyronaridine belongs to which class of drugs?
    - a. Antibiotic
    - **b. Anti malarial**
    - c. Anti viral
    - d. Anti fungal

Explanation :


Pyronaridine has high potency against *Plasmodium falciparum*, including chloroquine-resistant strains. Studies in various animal models have shown pyronaridine to be effective against strains resistant to other anti-malarials, including chloroquine. Resistance to pyronaridine appears to emerge slowly and is further retarded when pyronaridine is used in combination with other antimalarials, in particular, artesunate. Pyronaridine toxicity is generally less than that of chloroquine, though evidence of embryotoxicity in rodents suggests use with caution in pregnancy.

69. G6PD deficiency leads to ?
    a. Shock
    b. Galactosemia
    c. Hemolytic anemia
    d. Hereditary fructose intolerance

Explanation :

G6PD deficiency is an inherited condition in which the body doesn't have enough of the enzymeglucose-6-phosphate dehydrogenase, or G6PD, which helps red blood cells (RBCs) function normally. This deficiency can cause hemolytic anemia, usually after exposure to certain medications, foods, or even infections.


70. Identify the structure shown in image indicated by arrow.



www.medicalcoupons.in

a. 4 th ventricle
b. Cerebral aqueduct
c. Cerebellum
d. Thalamus

Explanation : The arrow in the above image indicates 4 th ventricle.



71. Which of the following is the Antibody for neonatal lupus ?
a. Anti centromere antibody
b. Anti ds DNA
c. Anti Ro
d. Anti mitochondrial antibody

Explanation :

Anti centromere antibody – CREST SYNDROME

Anti ds DNA – SLE

Anti RO – sjogren's syndrome , neonatal lupus

Anti mitochondrial antibody – primary biliary cirrhosis

72. Definition of antiport system ?
a. Two molecules simultaneously enter the cell via same channel
b. One molecules enter and one molecules leave the cell via same channel simultaneously .
c. Two molecules simultaneously leaves the cell via same channel .
d. Only one molecules leave the cell .

www.medicalcoupons.in

Explanation :

An antiport is an integral membrane transport protein that simultaneously transports two different molecules, in opposite directions, across the membrane.


73. All of the following comes under COAD except ?
    a. Asthma
    b. Chronic bronchitis
    c. Emphysema
    d. None of the above

Explanation :

COAD – chronic obstrcutive airway disease includes all the 3 ie, asthma, chronic bronchitis , emphysema. COAD is associated with chronic obstruction of the flow of air out of the lungs. This obstruction is generally permanent and progressive over time. People with this condition often cough up large amounts of sputum for several months of the year because the inflammation stimulates the production of mucus. This mucus further narrows the already inflamed airways. The ideal way to prevent this condition from happening and to stop it getting worse is to not smoke.

74. Which of the following pass through Guyon's canal ?
    a. Radial nerve
    b. Median nerve
    c. Ulnar nerve
    d. Auxillary nerve

Explanation :



75. Germ tube formation is seen in which of the following ?
    a. Candida albicans
    b. Cryptococcus neoformans
    c. Histoplasmosis
    d. Chromoblastomycosis

www.medicalcoupons.in

Explanation :

Candida albicans has evolved mechanisms to evade innate immunity, which is an important reason that candidiasis is a major complication in AIDS patients. *C. albicans* resists macrophage phagocytosis via a mechanism that does not stimulate apoptosis in macrophages (22). *C. Albicans* are defective in making germ tubes and cannot escape the macrophages following phagocytosis and are killed.

76. Ovarian follicular reserve is best accessed by which of the following ?
    a. Urine LH
    b. Antral follicle count
    c. Endometrial biopsy
    d. Measuring HCG

Explanation :

Ovarian reserve plays a crucial role in achieving pregnancy following any treatment in subfertile women. The estimation of ovarian reserve is routinely performed through various ovarian reserve tests (ORTs) in an effort to predict the response and outcome in couples prior to *In Vitro* Fertilization and counsel them. Most widely used tests are basal follicle stimulating hormone and anti-Mullerian hormone and antral follicle count. Here the best answer is antral follicle count.

77. All of the following is true regarding crohn's disease except?
    a. Skip lesions
    b. Feeling of mass or fullness in the abdomen
    c. more common in women than in men
    d. recurrence is uncommon

Explanation :

Crohn's disease is a Inflammation can affect any part of the digestive system, from the mouth to anus. Common symptoms can include:

- diarrhoea
- abdominal pain
- fatigue (extreme tiredness)
- unintended weight loss
- blood and mucus in your faeces (stools).

People with Crohn's disease sometimes go for long periods without symptoms or with very mild symptoms. This is known as remission. Remission can be followed by periods where symptoms flare up and become particularly troublesome.

www.medicalcoupons.in

78. Which of the following is not correct regarding reduced ORS ?
    a. Sodium 75
    b. Trisodium Citrate 10
    c. Potassium chloride 20
    d. Glucose 90

Explanation :

WHO ORS Formulae - comparison
(All values in mmol / litre)

	Standard ORS (g/L)	--- (mEq/L)	Reduced osmolarity ORS (2003)
Sodium Chloride	2.6	90	75
Chloride		80	65
Potassium chloride	1.5	20	20
Trisodium Citrate	2.9	10	10
Anhydrous Glucose	13.5	111	75
Total Osmolarity		311	245

79. Folinic acid is a antidote for which of the following?
    a. 6 – mercaptopurine
    b. Methotrexate poisoning
    c. Cisplatin toxicity
    d. Cyclophosphamide

Explanation :

Methotrexate inhibits dihydrofolate reductase and also inhibits thymidylate synthase. Methotrexate toxicity can be reduced by administration of leucovorin (folinic acid).

80. Which of the following is not a component of superior mediastinum ?
    a. Esophagus
    b. Thoracic duct
    c. SVC
    d. Desending aorta

Explanation :

Superior mediastinum extends upwards terminating at the superior thoracic aperture.it includes arch of aorta , SVC , vagus nerve , phrenic nerve ,thymus, trachea, esophagus and thoracic duct. Desending aorta is a content of posterior part of inferior mediastinum.

www.medicalcoupons.in

81. Identify the given image below?



a. Colorectal carcinoma
b. Diffuse esophageal spasm
c. Intussception
d. Crohn's disease

Explanation :

The "apple core sign" was first described in barium enema examinations of the colon. The apple core appearance is the visual manifestation of an annular lesion of the bowel with irregular overhanging edges and shouldered margins . The apple core sign is classically seen in cases of colon carcinoma.

82. Which of the following k+ sparing diuretics that doesn't cause gynecomastia ?
a. Spironolactone
b. Triamterene
c. Amiloride
d. Eplenorone

Explanation :

Amiloride acts by inhibiting the epithelial sodium channel (ENaC). Its mechanism is different to spironolactone ,which is an aldosterone receptor antagonist. Therefore amiloride doesn't cause gynecomastia. Eplenorone has similar action like spironolactone but it have less adverse effects like gynecomastia and vaginal bleeding when compared with spironolactone.

83. Lid lag sign of hyperthyroidism is?
a. Moebius sign
b. Von graefe's sign
c. Dalyrymptes sign
d. Stellwag's sign

www.medicalcoupons.in

Explanation :



84. In case of hanging , during autopsy which of the following cavity should be opened 1st?
    - **a. Cranial cavity**
    - b. Thoracic cavity
    - c. Abdominal cavity
    - d. Pelvic cavity

85. CEA is a marker for the follow up of which of the following ?
    - a. Pheochromocytoma
    - **b. Colorectal carcinoma**
    - c. Choriocarcinoma
    - d. Prostatic cancer

Explanation:


Carcinoembryonic antigen (CEA) is a glycoprotein, which is present in normal mucosal cells but increased amounts are associated with adenocarcinoma, especially colorectal cancer. CEA therefore has a role as a tumour marker CEA may be elevated in colorectal cancer, which is where it is most clinically useful. However, it may also be elevated in a wide variety of other malignant and benign conditions.

www.medicalcoupons.in

86. Heberdan arthropathy is a feature of ?
    - **a. Osteoarthritis**
    - b. Osteoporosis
    - c. Psoriatic arthritis
    - d. Osteogenesis imperfect

Explanation :

Heberden's nodes seen in distal interphalangeal joints is a characteristic feature of idiopathic osteoarthritis. Bouchard's nodes is also a feature of osteoarthritis seen in proximal interphalangeal joints.




87. Tb of phalanges is called a
    - a. Caries sicca
    - **b. Spina ventosa**
    - c. Pot spine
    - d. Military TB

Explanation :


Tuberculous dactylitis, also known as **spina ventosa**, is a rare skeletal manifestation of tuberculosis where the short tubular bones (i.e. phalanges, metacarpals, metatarsals) are affected.

88. Identify the part in the image indicated by arrow?
    - a. Sphincter of oddi
    - b. Pancreas
    - **c. 3 rd part of duodenum**
    - d. Jejunum



www.medicalcoupons.in


Explanation :



89. Distribution of rash in chickenpox is ?

a. Centripetal distribution
b. Centrifugal distribution
c. Spread downwards
d. Spread upwards

Explanation :



www.medicalcoupons.in

90. Most common tumour to occur in HIV patients is ?
    a. Cervical cancer
    b. Hodgkin's lymphoma
    c. Kaposi scarcoma
    d. Nasopharyngeal carcinoma

Explanation :

Infection with HIV weakens the immune system and reduces the body's ability to fight viral infections that may lead to cancer. The viruses that are most likely to cause cancer in people with HIV are:

- Kaposi sarcoma-associated herpesvirus (KSHV), also known as human herpesvirus 8 (HHV-8), which causes Kaposi sarcoma is the most common cancer to occur in HIV patients and some subtypes of lymphoma.
- Epstein-Barr virus (EBV), which causes some subtypes of non-Hodgkin and Hodgkin lymphoma.
- Human papillomaviruses (HPV), high-risk types (16 & 18) of which cause cervical cancer, most anal cancers, and oropharyngeal, penile, vaginal, and vulval cancer.
- Hepatitis B virus (HBV) and hepatitis C virus (HCV), which both cause liver cancer.

91. Visual Crowding is a phenomenon seen in which of the following ?
    a. Astigmatism
    b. Amblyopia
    c. Presbyopia
    d. Senile cataract

Explanation :

Amblyopia is a developmental visual disorder which is characterized by reduced visual acquity (uncorrectable by lense in the absence of any detectable defect in eye or visual pathway). Crowding phenomenon is a feature of an amblyopic eye. It is partial loss of sight , best detected and treated by 3 or 4 yrs of age for best results. Occlusion of normal eye to encourage the use of amblyopic eye is the most effective treatment.

www.medicalcoupons.in

92. Man with Lepromatous leprosy who had undergone chemotherapy develops various nodules is a example of which type of hypersensitivity ?
    a. Type 2
    b. Type 3
    c. Type 4
    d. Type 5

Explanation :

Leprosy is a chronic granulomatous infection principally affecting the skin and peripheral nerves caused by the obligate intracellular organism *Mycobacterium leprae.* The disease causes skin lesions and neuropathy. Complications secondary to the neuropathy can result in deformity and disability. The immunological response mounted by the host dictates the clinical phenotype that develops. People with leprosy show a spectrum of clinical types. Tuberculoid disease is the result of high cell-mediated immunity with a largely Th1 type immune response. Lepromatous leprosy however is characterized by low cell-mediated immunity with a humoral Th2 response. Type 1 reactions are delayed hypersensitivity(type 4) reactions that occur in borderline disease. Type 2 or erythema nodosum leprosum (ENL) reactions occur in borderline lepromatous and lepromatous disease (type -3).

93. Site of hordeolum externum ?
    a. Meibomian glands
    b. Gland of zeis
    c. Wolfring's gland
    d. Glands of Moll

Explanation :

A hordeolum is a common disorder of the eyelid. It is an acute focal infection (usually staphylococcal) involving either the glands of Zeis (external hordeola, or styes) or, less frequently, the meibomian glands (internal hordeola). In the early stages of disease the gland become swollen, hard and painful. Usually the whole edge of the lid is edematous. Subsequently abscess is formed which points near the base of one of the cilia. Pain is considerable until pus is evacuated.

94. Intermittent ring causing dysphagia is seen in which of the following disease?
    a. Diffuse esophageal spasm
    b. Schatzki's ring
    c. Esophageal web
    d. Streakhouse syndrome

www.medicalcoupons.in

Explanation :

Schatzki rings are fixed smooth, benign, circumferential, and narrow ring of tissue in the lower end of the esophagus. Schatzki rings are a common cause of intermittent dysphagia. Typically, the dysphagia is intermittent or short-lived. Food may become stuck in the esophagus but can be forced down by drinking liquids or may be regurgitated. The dysphagia may not recur for months or years. Patients may have associated symptoms of gastroesophageal reflux. Barium esophagography is the preferred method of detection of Schatzki rings.

95. Identify the procedure given in the picture?



a. Robotic surgery
b. Laproscopic cholecystitis
c. Endoscopy
d. LAZER ablation

Explanation :



Robotic surgery, or robot assisted surgery, is one of the developments that is at the forefront of this shift. Robotic surgery involves the use of surgical robots that can be controlled by doctors through computers. This results in less pain and blood loss for

www.medicalcoupons.in

the patient, and quicker healing time. Even higher accuracy can be achieved when computer-generated images are used to guide surgeries. One such possibility is the use of a real time MRI scan to guide surgeries.

96. Most common site of zollinger Ellison syndrome is?
    a. Cardiac part of stomach
    b. Pylorus part of stomach
    c. Duodenum
    d. Jejunum

Explanation :

Zollinger-Ellison syndrome (ZES) is a disease of the gastrointestinal system. People who have ZES develop tumors known as gastrinomas in the pancreas and duodenum (the first section of the small intestine). The gastrinomas caused by ZES secrete the hormone gastrin. Because gastrin creates excessive stomach acid, 90 percent of patients with ZES develop stomach and duodenal ulcers.

97. Mummification is seen in which conditions?
    a. Moist and humid environment
    b. Lead poisoning
    c. Desiccation of body in hot and dry weather
    d. Electric burns


Explanation :

Mummification is a another modification of putrefaction. Hot and dry climate favours it. Dehydration and drying causes evaporation of water and shrinkage of cadaver. The natural appearances and body features are preserved. The mummified body is odourless, dry and leathery brown. The time required is 3 months to 1 yr. It begins in the exposed parts of the body like face, hands, feet and then extends to internal organs. Arsenic and Antimony poisoning favours mummification .

98. In Gamekeeper's thumb which of the following is damaged?
    a. medial collateral ligament
    b. Ulnar collateral ligament
    c. anterior ligamnet
    d. posterior ligament

www.medicalcoupons.in

Explanation :



Ulnar collateral ligament of thumb is damaged

99. Carotid sinus massage leads to ?
    a. Reflux tachycardia
    b. Reflux bradycardia
    c. Bradypnea
    d. Tachypnea

Explanation :

Carotid sinus massage is performed for three reasons:

• To break junctional tachycardia (AVNRT and AVRT) by blocking AV nodal conduction (and cause Bradycardia).

• To reveal on the P wave pattern of an atrial arrhythmia by reducing the frequency of AV nodal conduction during the tachycardia so that the QRS complexes do not mask the artrial activity.

• To test for carotid sinus hypersensitivity, which is a severe fall in blood pressure or heart rate in response to carotid sinus stimulation.

100. Incidence is
    a. No. Of new case
    b. No. Of existing case
    c. 1st case in community
    d. 1st case that comes to doctor

Explanation :

Incidence is defined as the no. of new cases occurring in a defined population during specific period of time. It can be determined from cohort study .

Incidence = no of new case /population at risk *1000

Incidence is a RATE . (expressed per 1000)

www.medicalcoupons.in

101. Invertogram to be done in a new born:
   a. Immediately
   b. After 2 hrs
   c. After 4 hrs
   d. After 6 hrs

Explanation :

Invertogram is done for detection of imperforate anus. It is done after six hours of birth in infant u sufficient air may have collected in large intestine to cast a X-ray shadow.

102. For hyponatremia select the electrocardiographic finding with which it is most commonly associated?
   a**. No known electrocardiographic abnormalities**
   b. Prolonged QT interval
   c. Short QT interval
   d. Widened QRS complex

Explanation :

Hypokalemia typically increases automaticity of myocardial fibers, which results in ectopic beats or arrhythmias. Electrocardiography in hypokalemia reveals flattening of the T wave and prominent U waves. Hyperkalemia decreases the rate of spontaneous diastolic depolarization in all pace maker cells. It also results in slowing of conduction. One of the earliest electrocardiographic signs of hyperkalemia is the appearance of tall, peaked T waves. More severe elevations of the serum potassium result in widening of the QRS complex. At serum sodium levels compatible with life, neither hyponatremia nor hypernatremia results in any characteristic electrocardiographic abnormalities.

103. Over 75% of the strength of the intact abdominal wall lies in the –
   a. Skin
   b. Subcutaneous tissue
   c. Aponeurosis
   d. Peritoneum

Explanation:

Aponeurosis of rectus sheath is very strong and provides maximum strength of intact abdominal wall.

www.medicalcoupons.in

104. In amblyopia, the treatment of choice is:
   a. Surgery
   b. Orthoptic exercise
   c. Spectacles
   d. Conventional occlusion

Explanation :

Amblyopia is a developmental visual disorder which is characterized by reduced visual acquity (uncorrectable by lense in the absence of any detectable defect in eye or visual pathway). Crowding phenomenon is a feature of an amblyopic eye. It is partial loss of sight , best detected and treated by 3 or 4 yrs of age for best results. Occlusion of normal eye to encourage the use of amblyopic eye is the most effective treatment. Occlusion therapy must be carried out for as short a period as may be expected to give improvement in the acuity of the amblyopic eye and yet avoid the risk of occluded eye itself becoming amblyopic from lack of visual output.

105. Paranoid schizophrenia is seen with which of the following substance abuse?
   a. Cocaine
   b. Amphetamine
   c. Barbiturates
   d. Opiods


Explanation :

Schizophrenia like syndrome may be seen with amphetamine use, mescaline ,phencyclidine abuse, LSD abuse , huntington's chorea, homocystinuria , hemochromatosis , wilson's disease, acute intermittent porphyria. Thought disorder in paranoid schizophrenia are bizzre delusions ,1st rank symptoms , and formal thought disorder. Insight is absent.

106. Which of the following is not a component of carpel tunnel ?
   a. Median nerve
   b. Flexor digitorum profundus
   c. Flexor pollicis longus tendon
   d. Ulnar nerve

www.medicalcoupons.in

Explanation :



107. Which antiviral agent most often causes anemia, leucopenia, or thrombocytopenia?

a. Zidovudine
b. Ribavarin
c. Acyclovir
d. Amantadine

Explanation:

The most serious side effect of zidovudine is bone marrow suppression. Ganciclovir (Cytovene) may also cause these adverse effects, but acyclovir (Zovirax) usually does not. Didanosine and zalcitabine (other dideoxynucleosides used for HIV) are less likely to cause bone marrow suppression than zidovudine and are more frequently associated with pancreatitis and sensorimotor neuropathy.

108. Maximum disinfectant is seen with?

a. Ethylene oxide
b. 2% glutaraldehyde
c. Alcohol 70- 80%
d. Cresol

Explanation:

Cresol emulsions are very powerful disinfectant . Also known as “all purpose general disinfectant “.

www.medicalcoupons.in

109. Hippus is seen in which poisoning ?
   a. Abrus
   b. Aconite
   c. Datura
   d. Alcohol

Explanation :

The condition called hippus is also called pupillary athetosis, or spasmodic, rhythmic dilation and constriction of the pupils.

Aconite- in late stages Hippus is seen.

Alcohol- pupils are earlier dilated, later constricted.

Barbiturates- pupils are earlier constricted, later dilated.

110. Superior vena caval syndrome is most commonly caused by ?
   a. Lung cancer
   b. pheochromocytoma
   c. cirrhotic liver
   d. rib fractures

Explanation :

SVCS is more common for people who have lung cancer, non-Hodgkin lymphoma, or cancers that spread to the chest. But cancer can cause SVCS in other ways:

- A tumor in the chest may press on the superior vena cava.
- A tumor may grow into the superior vena cava, causing a blockage.
- If cancer spreads to the lymph nodes surrounding the superior vena cava, the lymph nodes may enlarge and press on or block the vein.
- A blood clot may appear in the vein. This is caused by a pacemaker wire or an intravenous catheter, which is a flexible tube placed in a vein to take out or put in fluids

111. Common cause for delayed puberty in girls?
   a. Turner syndrome
   b. Constitutional delay
   c. Primary ovarian failure
   d. Addison's disease

Explanation:

A girl who has not started to have breast development by the age of 13 is considered to be delayed. Girls with delayed puberty are simply late maturers, but once they start, puberty will progress normally. This is called constitutional delayed puberty and

www.medicalcoupons.in

is more common in boys than girls. Often, this is something that is inherited from the parents, so it is more likely to occur if the mother started her periods after age 14 (the average is about 12 ½) or if the father was a "late bloomer." Decreased body fat is a major cause of pubertal delay in girls. Here the best option is constitutional delay**.**

112. Treatment of choice for atropic rhinitis is?
    a. Vidian neurectomy
    b. McGovern's technique
    c. Young's operation
    d. FESS

Explanation:

Atrophic rhinitis is defined as a chronic nasal disease characterised by progressive atrophy of the nasal mucosa along with the underlying bones of turbinates. The common symptoms may include foetor, ozaena, crusting/nasal obstruction, epistaxis, anosmia / cacosmia and secondary infection with maggot infestation. The mainstay of treatment is conservative (for example, nasal irrigation and douches; nose drops (e.g. glucose, glycerine, liquid paraffin); antibiotics and antimicrobials; vasodilators and prostheses). Surgical treatment -(Young's operation) aims to decrease the size of the nasal cavities, promote regeneration of normal mucosa, increase lubrication of dry nasal mucosa and improve the vascularity of the nasal cavities.

113. Most common site of peripheral artery aneurysm is ?
    a. Iliofemoral artery
    b. Internal iliac artery
    c. Subclavian artery
    d. Popliteal artery

Explanation :

About 70% of peripheral arterial aneurysms are popliteal aneurysms; 20% are iliofemoral aneurysms. Aneurysms at these locations frequently accompany abdominal aortic aneurysms, and > 50% are bilateral. Rupture is relatively infrequent. Peripheral arterial aneurysms occur in men much more often than in women (> 20:1); mean age at presentation is 65. Aneurysms in arm arteries are relatively rare; they may cause limb ischemia, distal embolism, and stroke.

114. Which organ assessment should be done before CECT?
    a. Liver
    b. Kidney
    c. Heart
    d. Brain

www.medicalcoupons.in

Explanation:

Renal function should be assessed before taking CECT. Iodinated contrast media is highly nephrotoxic.

115. Polyuria is a feature of all of the following except:
   a. Lithium toxicity
   b. Hypocalcemia
   c. ADH deficiency
   d. Hypokalemia

Explanation:

Polyuria is defined, as 24-hour urine volume greater than 3 litres. Hypocalcaemia does not cause polyuria. Hypercalcaemia causes polyuria by solute diuresis. Solute diuresis is also seen with glucose, mannitol, radiocontrast agent, urea, resolving ATN.

Lithium causes nephrogenic diabetes insipidus and hence polyuria. ADH deficiency causes central diabeted insipidus. The various causes are postoperative (removal of pituitary), trauma supra-or intrasellar tumour, Sheehan's syndrome, infections etc.

116. Which one of the following vectors transmits scrub typhus?
   a. Louse
   b. Ticks
   c. Mite
   d. Flea

Explanation:

Scrub typhus is caused by Orientia tsutsugamushi, which is maintained in nature by transovarian transmission in trombiculid mites, mainly of genus Leptotrombidium. After hatching, the infected larval mites (chigger) inoculate organisms into the skin while feeding. Clinical features are fever, headache, myalgia, cough, gastrointestinal symptoms, regional lymphadenopathy etc. Doxycycline or chloramphenicol is used for treatment.

117. Food poison is a example of which kind of epidemics?
   a. Seasonal trends
   b. Multiple source epidemics
   c. Probagated epidemics
   d. Point source epidemics

Explanation:

Seasonal trends - eg. Measles ,URI

Multiple source epidemics – eg. Contaminated well in a village

www.medicalcoupons.in

Probagated epidemics – eg. HIV (person to person spread)

Point source epidemics – eg.food poisoning , Bhopal gas tragedy.

118. Muscle stretch is due to the action of which of the following ?
   a. Muscle spindle
   b. microfilaments
   c. Golgi tendon organ
   d. Sarcomere

Explanation:

Muscle spindle is present in skeletal muscle. It has a nuclear bag fibre and nuclear chain fibre. It helps in the contraction of skeletal muscles. Its function is to change the length of the muscle.

119. Pre exposure schedule of rabies vaccine is
   a. 0, 3, 7, 14, 21, 28, 90
   b. 3, 7 , 21
   c. 0, 3, 7, 14 ,28
   d. 0, 7 , 28

Explanation:

Hydrophobia is pathognomic.

Causative agent is – Lyssavirus type-1.

Types of rabies – street virus and fixed virus.

Vaccine schedule :

post exposure prophylaxis : 0, 3, 7, 14 ,28, 90

pre exposure prophylaxis : 0,7,28

post exposure prophylaxis in those vaccinated previously : 0,3,7

120. Which of the following drug cause uterine relaxation ?
   a. Dinoprost
   b. Carbetocin
   c. Methergine
   d. Atosiban

Explanation:

Preterm birth is strongly associated with neonatal death and long-term neurological morbidity. The purpose of tocolytic drug administration is to postpone threatening preterm delivery for 48 hours to allow maximal effect of antenatal

www.medicalcoupons.in

corticosteroids and maternal transportation to a center with specialized neonatal care facilities. There is uncertainty about the value of atosiban (oxytocin receptor antagonist) and nifedipine (calcium channel blocker) as first-line tocolytic drugs in the management of preterm labor.

121. A 3 yr old boy was brought to ER with multiple coalescing lesions on his face and extremites.



a. Herpes simplex
b. Impetigo
c. Dermatitis herpetiformis
d. Phemphigus.

Explanation :

Impetigo is the most common bacterial infection in children. primarily caused by *Streptococcus pyogenes* or *Staphylococcus aureus*. Impetigo is classified as either nonbullous (impetigo contagiosa) (about 70% of cases) or bullous. nonbullous impetigo commonly have honey-colored crusts with a moist erythematous base as shown in the above image. Bullous impetigo is considered to be less contagious than the nonbullous form.

122. Skin of the parotid gland is supplied by which of the following nerve?

a. Greater auricular nerve
b. Auriculotemporal nerve
c. Mandibular nerve
d. Vagus nerve

Explanation :

The parotid gland receives **sensory** and **autonomic** innervation. The autonomic innervation controls the rate of saliva production.
Sensory innervation is supplied by the **auriculotemporal nerve** (gland) and the **great auricular nerve** (fascia and skin). The parasympathetic innervation to the parotid gland has a complex path. It begins with the **glossopharyngeal nerve.** The **auriculotemporal nerve** then carries parasympathetic fibres from the otic ganglion to the parotid gland. Parasympathetic stimulation causes an increase in saliva production. Sympathetic innervation originates from the superior cervical ganglion, part of the paravertebral chain.

www.medicalcoupons.in

123. Which of the following tumour is rathke's pouch derivative ?
   a. Glioblastoma multiforme
   b. Craniopharyngioma
   c. Vestibular schwanoma
   d. Oligodentroglioma

Explanation :

A craniopharyngioma is a benign (noncancerous) tumor arising from small nests of cells near the pituitary stalk. Adamantinomatous (ordinary) craniopharyngioma occurs in children and tends to be less solid than papillary craniopharyngioma. Papillary craniopharyngioma occurs in adults and is a more solid tumor. Increased pressure within the brain causes many of the symptoms associated with this tumor. Other symptoms result from pressure on the optic tract and pituitary gland Surgery to remove the tumor is usually the first step in treatment. If hydrocephalus (excess water in the brain) is present, a shunt (drainage system) may be placed during surgery. The shunt will help remove excess cerebrospinal fluid from the brain and ease the pressure.

124. For diagnosis of schizophrenia symptoms should be present for how long period?
   a. 2 weeks
   b. 1 month
   c. 6 month
   d. 1 year

Explanation :

Two or more of the following each present for a significant portion of time of during 1 month

- Delusions
- Hallucinations
- Disorganized speech
- Negative symptoms ie, affective flattening, alogia or avolition .

Continous signs of disturbance persist for atleast 6 months.

This 6 month period must include atleast 1 month of symptoms.

125. Leucocyte rolling over endothelial cells of the blood vessel is carried out by which of the following?
   a. Selectins
   b. Integrins
   c. PECAM 2
   d. E- Cadherins

Explanation:

Rolling is a process of transient adhesion of leucocytes with the endothelial cells. Selectins are the important molecules responsible for it. They interact with complementary molecules resulting in transient adhesion.

126. Which of the following tumour is most common in post renal transplant patients?
    a. Non Hodgkin's Lymphoma
    b. Hepatocellular carcinoma
    c. Non melanoma skin cancer
    d. Kaposi's sarcoma

Explanation :

There is an overall 3–5-fold increase in cancer risk in transplant recipients compared with the general population, with skin cancers and lymphoma particularly prevalent. Cancers in transplant recipients are often more aggressive than those in the general population, with poor prognosis, particularly for gastrointestinal tumours and lymphomas.

127. In ERG 'A' wave corresponds to?
    a. Rods and cones
    b. Pigment epithelium
    c. Nerve bundle layer
    d. Artefact

Explanation:

Electroretinography: A test in which the electrical potentials generated by the retina of the eye are measured when the retina is stimulated by light. Abbreviated as ERG. In an ERG, an electrode is placed on the cornea at the front of the eye that measures the electrical response of the rods and cones, the visual cells in the retina at the back of the eye. An ERG may be useful in the evaluation of hereditary and acquired disorders of the retina. The instrument used to conduct ERG is an electroretinograph, and the resultant recording is called an electroretinogram

waves	Function
A wave	Activity of rods and cones
B wave	A response arising from bipolar cell layers
C wave	Retinal metabolism of the pigment

www.medicalcoupons.in

128. Which of the following is not associated with cephal hematoma?
   a. Usually involves parietal and occipital bones
   b. Bleeding is sub periosteal
   c. Swelling is present at birth
   d. Usually reabsorbed in 2-3 months.

Explanation:

Cephalhaematoma is a collection of blood between pericranium and skull bones following forceps delivery or normal labour due to rupture of small emissary veins or fracture of scalp bones. It appears 12 to 24 hours afterbirth. Cephalhaematoma is never present at birth. It appears a few hours after birth and at times on the second day. It causes hyperbilirubinaemia due to release of bilirubin from collected blood. Caput succedaneum is present at birth.

129. Culex is a vector for which of the following diseases?
   a. Japanese encephalitis
   b. Yellow fever
   c. Brugian filariasis
   d. Chickengunya

Explanation :

Japanese encephalitis :

Causative agent : Group- B Arbovirus

Host factors : Pigs are "Amplifier host" .

Cattles and Buffalos are 'mosquito attractants'.

Birds are also involved

Man is a 'Incidental Dead End Host'.

Vectors of JE: Culex tritaeniorhynchus (most important vector)

Culex vishnuii

Incubation period of JE: 5- 15 days (9-12 days in mosquito)

Case fatality rate : 20 – 40%

Vaccines : nakayama strain /Beijing strain

SA-14-14-2 strain

www.medicalcoupons.in

130. Which of the following structure is not visualized in hysteroscopy ?
   a. endocervix
   b. Uterine surface
   c. Tubal ostia
   d. Endometrial cavity

Explanation:

Hysteroscopy permits full visualization of the endometrial cavity and endocervix and is helpful in diagnosing focal lesions that are missed with endometrial sampling. Uterine surface, which is the outermost layer of uterus ie, perimetrium cannot be visualized by hysteroscopy.

131. Presenting case of infant with intussception , management to be done is?
   a. fluid resuscitation + NG tube decompression + antibiotic + hydrostatic reduction
   b. fluid resuscitation + antibiotic + surgery
   c. fluid resuscitation + NG tube decompression + surgery
   d. fluid resuscitation + NG tube decompression + antibiotic

Explanation:

Intussusception is the most common cause of intestinal obstruction in children between ages 6 months to 3 years old. A segment of intestine invaginates into the adjoining intestinal lumen causing bowel obstruction. Mc site is ileocolic. The symptoms includes classic triad of vomiting, abdominal pain, and red current jelly stools. Intussusception is managed by hydrostatic and pneumatic reduction of the obstructed bowel. If manual reduction is not possible or perforation is present, a segmental resection with an end-to-end anastomosis is performed.

132. Chancroid is caused by ?
   a. Campylobacter jejuni
   b. N. Gonorrnoea
   c. H.Ducreyi
   d. Treponema Pallidum

Explanation:

Chancroid (soft chancre)- caused by haemophilus ducreyi. Single or multiple painful ulcers with deep ulceration. The ulcer is soft ,non-indurated, dirty looking. It produces painful, inguinal lymphadenitis k/a inflammatory bubo. Management is with ceftriaxone i.m or azithromycin 1 gm single dose.

133. Drug of choice for acute gout is ?
   a. Naproxen
   b. Probenacid
   c. Allopurinol
   d. Febustat

www.medicalcoupons.in

Explanation:

Naproxen is a NSAIDS, is used to relieve pains from various conditions such as headaches, muscle aches , tendonitis , dental pains ,mensural cramps. It also reduces pain caused by arthiritis , bursitis ,and gout attacks. Probenecid and allopurinol is used for chronic gout attacks.

134. Identify the above given image :



a. Pithris pubis
b. Scabei
c. Head louse
d. Tick

135. Which of the following is not a metabolites of tyrosine?

a. Melanin
b. Melatonin
c. norepinephrine
d. T3 and T4

Explanation :

Tyrosine is required for the formation of thyroid hormones, melanin and catecholamines such as epinephrine, norepinephrine, dopamine. Melatonin is a metabolite of tryptophan.

136. Incubation period of hepatitis – B is

a. 10- 14 days
b. 30- 180 days
c. 15- 60 days
d. 3- 35 days

www.medicalcoupons.in

Explanation :

Type	Causative agent	Incubation period	Common modes of transmission
Hep -A	Enterovirus -72 (picornavirus)	15 - 45 days	Fecal – oral , sexual
Hep -B	Hepadnavirus	30 – 180 days	Sexual ,perinatal, percutaneous
Hep -C	Hepacivirus (flavivirus)	15 – 160 days	percutaneous
Hep –D	Viriods like	30 – 180 days	Sexual , perinatal , percutaneous
Hep -E	Calcivirus (alpha virus like)	15 – 60 days	Fecal – oral

137. Haemorragic conjunctivitis is caused by ?
   a. Cornebacterium diphtheria
   b. Adenovirus
   c. Enterovirus 70
   d. Chlamydia

Explanation:

Acute haemorragic conjunctivitis is a dark red spots on conjunctiva caused by Enterovirus 70.

Pseudomembranous conjunctivitis is caused by adenovirus.

True membranous conjunctivitis is caused by infection with cornebacterium diphtheria.

Swimming pool conjunctivitis is caused by Chlamydia.

138. Which of the following drugs is preferred in patients with hypertriglyceridemia?
   a. Simvastatin
   b. Colestipol
   c. Gemfibrozil
   d. Alirocumab

www.medicalcoupons.in

Explanation :

Fibric acid derivatives such as gemfibrozil ,clofibrate, fenofibrate, bezafibrate stimulate PPAR – alpha and it increase LPL and decrease triglycerides and hence preferred in patients with hypertriglyceridemia.


Cloestipol is a bile acid resins is bad for hypertriglyceridemia.

Simvastatin is a HMG coA reductase inhibitors are the doc for dyslipidemia.

139. Which of the following is not the characteristic feature of kwashiorkor?
   a. Edema
   b. Pluckable hair
   c. Preserved appetite
   d. Flaky paint dermatosis

Explanation :

Protein energy malnutrition is a spectrum of conditions ranging from mild undernutrition to extreme forms of malnutrition. It includes kwashiorkor and marasmus.



140. Which of the following is the tumour marker for hepatocellular carcinoma?
   a. CEA
   b. a – feto protein
   c. B- HCG
   d. calcitonin

www.medicalcoupons.in

Explanation :

CEA	Colon cancer
	Pancreatic cancer
	Lung and stomach cancer
AFP	Yolk sac tumour
	Hepatocellular carcinoma
B – HCG	Trophoblastic tumors like choriocarcinoma , seminoma etc.
Calcitonin	Medullary carcinoma of thyroid

141. H.pylori is a risk factor for causing which of the following cancers?
   a. GIST
   b. Gastric adenocarcinoma
   c. Squamous cell carcinoma of esophagus
   d. Hepatic adenoma

Explanation :

H.pylori is a gram –ve flagellated bacteria producing enzymes like phospholipase and urease. H.pylori is associated with peptic ulcer disease, gastric cancer and gastric mucosa associated lymphoma.

142. Which of the following class of drugs is not used in treatement of uterine fibroids?
   a. Nafarelin
   b. Tranexemic acid
   c. Misoprostol
   d. Mifepristone

Explanation :

Nafarelin – GnRH agonist is used to decrease the size of fibroids.


Tranexemic acid is a non hormonal drug which can be used to reduce the blood loss over short periods.

Mifepristone is also used to treat fibroids.

143. Which of the following nerve passes through cavernous sinus ?
   a. Occulomotor
   b. Abducens
   c. Trochlear
   d. Trigeminal V1

www.medicalcoupons.in

Explanation :



144. Schober's test is done for?

a. Flexion of lumbar spine
b. Pain with motion of hip
c. Neck pain and stiffness
d. Chest expansion

Explanation :

Schober's test is measure of flexion on lumber spine. This test is done in ankylosing spondylitis

145. Presumptive treatment is given in

a. Leprosy
b. Tuberculosis
c. Malaria
d. AIDS

Explanation :

Presumptive treatment is given for Malaria when API is ≥ 2.

• For leprosy, multidrug therapy is used.

• Short course chemotherapy is given for tuberculosis as DOTS regime.

• Universal precaution with body fluids is necessary in case of AIDS.

www.medicalcoupons.in

146. Dancing carotid is seen in ?
   a. Thyrotoxicosis
   b. Av fistula
   c. Hypothyroidism
   d. Blow out carotid

Explanation:

A jerky carotid pulse characterized by full expansion followed by quick collapse. Also known as Dancing carotid. Prominent carotid pulsations due to the wide pulse pressure in aortic regurgitation (Corrigan's sign). It is also seen in thyrotoxicosis.

147. Blood supply to the Lower end of esophagus is supplied by ?
   a. Inferior thyroid artery
   b. Left gastric artery and inferior phrenic artery
   c. Left pulmonary artery
   d. Short gastric artery

Explanation:

Blood supply of esophagus :

- The cervical portion is supplied by the inferior thyroid artery
- The thoracic portion is supplied by bronchial and esophageal branches of the thoracic aorta
- The abdominal portion is supplied by ascending branches of the left phrenic and left gastric arteries.

148. PR interval in ECG corresponds to ?
   a. Ventricular repolarization
   b. Atrial depolarization and AV nodal conduction
   c. Late repolarization of papillary muscle
   d. IDIO- ventricular rhythm

Explanation :

PR interval – normal duration : 0.12- 0.2 sec

This represents the time that it takes for the electrical impulse generated in the sinus node to travel through the atria and across the atrioventricular node to the ventricles.

149. Blood supply to the dorsum of penis is?
   a. Cavernosal artery
   b. Common penile artery
   c. Internal pudendal artery
   d. Inferior gluteal artery

www.medicalcoupons.in

Explanation :

The arterial blood supply of penis happens at the expense of external and internal pudendal arteries. The external pudendal artery starts from an internal wall of femoral artery on 2.5-2.7 cm below inguinal ligament. The internal pudendal artery is main source of blood supply of penis of the man.

150. Drug of choice for ovulation induction is
   a. Asoprisnil
   b. Tamoxifen
   c. Levonorgestrel
   d. Clomiphene citrate

Explanation :

It is non steroidal triphenylethylene compound related to DES. It binds to estrogen receptors in the hypothalamus , so the negative feedback of estrogen on hypothalamus is abolished resulting in increased pulsatile GnRH frequency which causes a rise in FSH and LH. It is used in cases of anovulations, pcos, anovulatory DUB with infertility. It is the drug of choice for postpill amenorrhoea. Most common side effect of clomiphene citrate is multiple pregnancy.

151. Most common cause of facial nerve palsy is ?
   a. Infection
   b. Idiopathic bell's palsy
   c. Head trauma
   d. Stroke

Explanation :

Facial nerve (7th cranial nerve) palsy is often idiopathic (formerly called Bell palsy). Idiopathic facial nerve palsy is sudden, unilateral peripheral facial nerve palsy. Symptoms of facial nerve palsy are hemifacial paresis of the upper and lower face. Tests (eg, chest x-ray, serum ACE level) are done to diagnose treatable causes. Treatment may include lubrication of the eye, intermittent use of an eye patch, and, for idiopathic facial nerve palsy, corticosteroids.

152. Curdy white discharge from vagina is due to ?
   a. Candidial vulvivaginitis
   b. Trachomonas vaginitis
   c. Bacterial vanginosis
   d. Atrophic vaginitis

Explanation:

Candidal vaginitis is most common during pregnancy , in OCP users , corticosteroid theraphy, diabetes and in all immune compromised and immunodeficient states.

www.medicalcoupons.in

Caused by – candida albicans , candida galbrata , candida tropicales.

Symptoms: Vaginal vulvar pruritus, burning, or irritation (which may be worse during intercourse) and dyspareunia are common, as is a thick, curdy white, cottage cheese–like vaginal discharge that adheres to the vaginal walls. Symptoms and signs increase the week before menses. Erythema, edema, and excoriation are common.

Diagnosis : Vaginal pH - <4.5

wet mount – budding yeast cells and filamentous elements.

Treatment : topical miconazole / cotrimazole

Systemic fluconazole

Both partners should be treated.

153. VMA in urine is used to diagnose which of the following conditions?
   a. Adrenal adenoma
   b. Cushing syndrome
   c. Pheochromocytoma
   d. Addison's disease

Explanation :

Vanillylmandelic acid (VMA) is one of the breakdown products (metabolites) of catecholamines. This test measures the amount of VMA that is passed into the urine, typically over a 24-hour period, to detect excess epinephrine and norepinephrine. It is used to detect tumors called neuroblastomas and other neuroendocrine tumors. Neuroendocrine tumors, such as neuroblastomas and pheochromocytomas, can sometimes produce large amounts of catecholamines, resulting in greatly increased concentrations of the hormones and their metabolites.

154. Horrock's apparatus is used to ?
   a. Disinfect the action of chlorine in water
   b. Chlorine demand estimation of water
   c. Remove the hardness of water
   d. Indicate the quality of water

Explanation :

It is used to find out the dose of bleeching powder required for disinfetion of water ie, chlorine demand estimation of water. Bleeching powder contains 33% of available chlorine. Orthotoludine – arsenite test is used for testing residual chlorine at the end of 1 hr contact. Chlorine acts best as a disinfectant when the PH of water is around 7. The standard prescribed for chloride is 200 mg/ litre. Maximum permissible level is 600 mg/litre.

www.medicalcoupons.in

155. Which of the following drug is used in the treatment of post menopausal osteoporosis?

a. Clomiphene citrate
b. fluoxetine
c. veralipride
d. Raloxifene

Explanation :

Raloxifene is SERM , that act as both estrogen agonist and antagonists depending on the tissue. Raloxifene has estrogen like action on bones and lipids and estrogen antagonists action on breast and endometrium. Due to its agonist effect on bones, it reduces the risk of fracture by 50% , specially in vertebra by increasing bone mineral density by 2-3%. 1st line treatment of post menopausal osteoporosis is bisphosphonates. Major adverse effect is predisposition of thromboembolism.

156. Which of the following cell is the Sole efferent fibre from cerebellum?

a. Stellate cells
b. Golgi cells
c. Purkinje cells
d. Granules cells

Explanation :

The three-layered cerebellar cortex contains six main neuronal cell types: **stellate** and **basket cells** in the molecular layer, **Purkinje cell** somata in the Purkinje cell layer, and **granule cells**, **Golgi cells**, and **unipolar brush cells** in the granular layer. The cerebellar nuclei and the lateral vestibular nucleus constitute the sole output (efferent) of the cerebellum, and play a central role in cerebellar circuitry and function. Purkinje cells represent the sole output neuron of the cerebellar cortex and thus changes in their function have significant impact on the function of the cerebellum as a whole. Understanding pathologic processes within these cells is crucial to develop strategies to prevent their loss and thus it reduce ataxia. Only excitatory cell of cerebellum is granules cells.

157. Which part of eye is used in biometric scans ?

a. Lens
b. Iris
c. Cornea
d. Retina

Explanation :


Biometrics is the science of human recognition. In biometrics, “Iris and Retinal scanning” technologies are known as ocular based identification technologies, means that they rely on unique physiological characteristics of eye to identify an individual. The structure of the iris is unique to the individual and can be used for

www.medicalcoupons.in

identification purposes. Iris recognition and retinal scanning are both very reliable modalities for biometric identification.


Iris recognition – It takes the reading of the characteristics of the iris	Retinal scanning – It is the analysis of blood vessels at the back of eye.

158. Identify the structure marked below



a. Azygous vein
b. Hemiazygous vein
c. Minor fissure of right lung
d. Major fissure of left lung

Explanation :



lac operon coding region is done by ?

a. Inducer
b. Promoter
c. Regulator
d. Operator

www.medicalcoupons.in

Explanation :

The *lac* operon is an **operon**, or group of genes with a single promoter (transcribed as a single mRNA). The genes in the operon encode proteins that allow the bacteria to use lactose as an energy source.

The **promoter** is the binding site for RNA polymerase, the enzyme that performs transcription.

The **operator** is a negative regulatory site bound by the *lac* repressor protein.

The ***lac* repressor** is a protein that represses (inhibits) transcription of the *lac*operon.

160. Which of the following condition doesn't cause gynecomastia ?
   a. Leprosy
   b. Tb
   c. Klinefelter syndrome
   d. Hyperthyroidism

Explanation :

Gynecomastia is triggered by a decrease in the amount of the hormone testosterone compared with estrogen. The hormones testosterone and estrogen control the development and maintenance of sex characteristics in both men and women. The pathologic causes of gynecomastia are diverse and may include Klinefelter syndrome, certain cancers, endocrine disorders such as hyperthyroidism , metabolic dysfunction, various medications, or may occur due to a natural decline in testosterone production. Gynecomastia is the most obvious clinical manifestation of hormone dysfunction in leprosy.

161. True statement about bronchopulmonary segment is
   a. It is according to the segmental bronchus.
   b. It is surgically unresectable
   c. Pul. Artery is intersegmental
   d. 10 segments in each lung

Explanation :

A bronchopulmonary segment is a division of the lung. These segments are the largest subdivisions of a lung lobe. Each supplied by a tertiary (segmental) bronchus. Segments are pyramidal-shaped, with their apices facing the lung root and their bases at the pleural surface. Supplied independently by a segmental bronchus and a segmental pulmonary Artery .10 segments on the right side and 8-10 segments on the left side. They are surgically resectable.

162. Which of the following is the anaerobic branched filamentous non acid fast bacilli ?

www.medicalcoupons.in

a. Clostridium tetani
b. Nocardia
c. Actinomyces
d. Atypical mycobacteria

Explanation:

Actinomyces : It is a anaerobic gram +ve and filamentous branching bacilli. It is non- acid fast bacilli , while Nocardia is partially acid fast. Actinomyces normal flora is in female genital tract , gingival. Transmission is endogenous due to tooth pick injuries. Most common organism is Actinomyces Israeli that produce indurated swelling and draining sinus that contains yellow sulphur granules. Most common site is lower jaw. It also produces solitary brian abcess and also cause Actinomycetoma. Doc for Actinomyces infection is penicillin.

163. Thalassemia in pregnancy is diagnosed by ?
a. Nestrof test
b. Hba2>3.5 %
c. Mutational analysis test
d. Chorionic villus sampling

Explanation :

Prenatal diagnosis of thalassemia can be done by chorionic villus sampling or by fetal blood sampling. Both tests are conducted under ultrasound guidance.

Chorionic villus sampling involves obtaining some cells from the placenta for testing. This is done during the 10th to 12th week of pregnancy.

Fetal blood sampling involves obtaining a small amount of fetal blood from the umbilical cord for testing. This procedure is done at about 18 - 20 weeks of pregnancy.

164. Fox - Fordyce spots are?
a. Seen due to blockade of apocrine gland
b. Variant of Sebaceous gland in lips
c. Pilo sebaceous units
d. Type of Eccrine sweat glands

Explanation :

Fordyce spots are visible sebaceous glands without hair follicles. They are present in 80–95% of adults. They are probably present at birth but become bigger and more visible from about puberty onwards. They are present on most of the body, and sometimes are visible on the penis, scrotum, lips, or inner (buccal) cheeks. These are completely normal and are not a sexually transmitted infection (STI).

www.medicalcoupons.in

165. Which of the following is seen in catarrhal stage of typhoid ?

a. High leucocyte
b. Eosinophilia
c. Leucopenia with relative lymphocytosis
d. Thrombocytopenia

Explanation :

Typhoid is a bacterial infection that can lead to a high fever, diarrhea, and vomiting. It can be fatal. It is caused by the bacteria *Salmonella typhi.*

Clinical features is divided into 4 stages :

1st week – prodromal symptoms similar to URT infection.

Step ladder pattern temperature

Relative bradycardia (faget's sign)

Malaise , headache, abdominal pain

Leucopenia with relative lymphocytosis

2nd week - macula- popular "Rose spot" rash

Constipation , confusional state , hepatospleenomegaly

Positive Widal test

3rd week - "week of complications"

Haemolytic anemia , meningitis ,acute cholecystitis , UTI , intestinal perforation and haemorrhage . Dehydration is a significant risk. Thrombocytopenia with risk of bleeding. Death can occur at this stage from overwhelming toxaemia.

4th week – "week of convalescence"

In the untreated patient the fourth week ceases the fever, mental state and abdominal distension slowly improve over a few days, but intestinal complications may still occur. Convalescence is prolonged, and most relapses occur at this stage.

Treatement :

The only effective treatment for typhoid is antibiotics. The most commonly used are ciprofloxacin (for non-pregnant adults) and ceftriaxone. Other than antibiotics, it is important to rehydrate by drinking adequate water.In more severe cases, where the bowel has become perforated, surgery may be required.

166. Obesity is not seen in which of the following disease ?

www.medicalcoupons.in

a. Prader-willi syndrome
b. Acromegaly
c. Bardet – Biedl syndrome
d. Carpenter syndrome

Explanation :

Acromegaly , a disorder of IGF-1 which causes excessive growth of the hands, feet, jaw and other internal organs in adults. In acromegaly , glucose does not suppress growth hormone. MRI shows pituitary tumours in 90% of acromegalic patients. The best confirmatory test for acromegaly is the oral glucose suppression test. Obesity is not seen with acromegaly.

Gigantism , abnormally high linear growth due to excessive action of IGF-1 before the closure of the epiphyseal growth plate in children.

167. Copycat suicide is mainly carried out by ?
a. School going Children
b. Newborns
c. Adults
d. Adolescent

168. Colipase enzyme is required for ?
a. Bile
b. Pancreatic lipase
c. Gastric secretion
d. Salivary enzymes

Explanation :

Colipase is a protein that functions as a cofactor for pancreatic lipase, with which it forms a stoichiometric complex. It also binds to the bile-salt covered triacylglycerol interface thus allowing the enzyme to anchor itself to the water-lipid interface.

169. Neonatal sepsis is caused by
a. Strep. Pyogenes
b. Strep. Agalactiae
c. Salmonella species
d. Strep.pneumoniae

Explanation :

Group – B streptococcal (strep.agalactiae) are normal commensals of the female genital tract and are important cause of neonatal sepsis and meningitis.

www.medicalcoupons.in

170. Which of the following is necessary in the process of exocytosis?

a. Calcium
b. Magnesium
c. Sodium
d. Potassium

Explanation :

Neurotransmitters are transmitted by exocytosis. They are chemical messages that are transported from nerve to nerve by synaptic vesicles. Synaptic vesicles are membranous sacs formed by endocytosis of the plasma membrane at pre-synaptic nerve terminals. The synaptic vesicle awaits a signal, an influx of calcium ions brought on by an action potential, which allows the vesicle to dock at the pre-synaptic membrane.

171. SERM that is associated with endometrial hyperplasia is ?

a. Raloxifene
b. Clomiphene citrate
c. Tamoxifen
d. Ormeloxifene

Explanation :

SERM – tamoxifen , doloxifene , toremifene

Possess estrogen antagonistic activity in breast and blood whereas agonistic activity in bone, uterus, and liver.

Their major indication is in the treatment of breast carcinoma.


They have beneficial effect on bone and lipid profile but increases the risk of endometrial carcinoma and thromboembolism.

172. Treatment for hyperkalemia is ?

a. Salbutamol
b. Calcium
c. Hemodialysis
d. All of the above

www.medicalcoupons.in

Explanation :



173. Vitamin required for the regulation of the synthesis of collagen is ?
   a. Vit. A
   b. Vit. C
   c. Vit. E
   d. Vit. K

Explanation :

Collagen contains specific amino acids – Glycine, Proline, Hydroxyproline and Arginine. These amino acids have a regular arrangement in each of the three chains of these collagen subunits. These amino acids are found at specific locations relative to glycine and are modified post-translationally by different enzymes, both of which require vitamin C as a cofactor.

174. Breast cancer of 4 cm with axillary lymph node involvement without distant metastasis is in which stage?
   a. 1
   b. 2a
   c. 3a
   d. 3b

Explanation :

www.medicalcoupons.in

STAGES OF BREAST CANCER

STAGE 0	Non-invasive cell malformations
STAGE 1	Tumour ≤2cm, Invasive breast cancer, early stage, invasion limited to breast
STAGE 2	2A -Tumour absent/ Tumour ≤2cm, cancer present in axillary lymph nodes, no spread to body -2cms≤ Tumour ≤5cm, no cancer in axillary lymph nodes, no spread to body 2B - 2cms≤ Tumour ≤5cm, cancer present in lymph nodes, no spread to body -Tumour >5cm, no cancer in axillary lymph nodes, no spread to body
STAGE 3	3A -Tumour absent/ 2cms≤ Tumour ≤5cm/ Tumour >5cm, cancer present in axillary and may be in lymph nodes near sternum 3B -Any size tumour, cancer present in skin or chest wall, cancer present in axillary lymph nodes or lymph nodes near sternum 3C -Any size tumour, cancer spread to lymph nodes near clavicle, , cancer present in axillary lymph nodes or lymph nodes near sternum
STAGE 4	Spread to body (metastasis)

175. Glutaraldehyde is not used as a disinfectant for which of the following ?
   a. Thermometer
   b. Endotracheal tube
   c. Cystoscope
   d. Bronchoscope

Explanation :

Glutaraldehyde is used for cystoscopes, bronchoscopes, rubber tubes, face masks, and endotracheal tubes. Wiping with alcohol was at least as efficient as immersion in a phenolic and is recommended for thermometer disinfection in hospitals.

176. Which of the following is associated with rheumatic heart disease?
   a. Roth's spots
   b. Pulsus paradoxus
   c. Koplik's spots
   d. MacCallum plaques

Explanation :

Rheumatic heart disease describes a group of short-term (acute) and long-term (chronic) heart disorders that occur as a result of acute rheumatic fever. Damage of the heart valves is a common result of rheumatic fever. It is usually seen in children who are 5 to 15 years old. Mural endocardial lesions can be seen as **MacCallum plaques** in rheumatic heart disease. These plaques appear as map-like areas of thickened, roughened, and wrinkled part of the endocardium in the left atrium.

www.medicalcoupons.in

177. Burning sensation in extremites is due to ?

a. Pheripheral neuropathy
b. Poliomyelitis
c. Syringomyelia
d. Multiple sclerosis

Explanation :

Peripheral neuropathy is damage to the peripheral nerves and results in a tingling, painful, or burning sensation in the extremities. It most commonly occurs in the legs. Other symptoms of peripheral neuropathy can include weakness and numbness. Poorly controlled diabetes is one of the most common causes of peripheral neuropathy, but a number of conditions can be responsible for damage to the peripheral nerves.

178. Burned airways, 1st line management is ?

a. Tracheostomy
b. Elective intubation
c. Fluid replacement
d. Prophylactic corticosteroid

Explanation :

In cases of thermal burns to the airway, many burn surgeons prefer ET intubation over blind insertion techniques, but supraglottic airways are acceptable if intubation isn't available or can't be established quickly.

179. Amniocentesis can be done earliest in which of the following weeks?

a. 8 to 10 wks
b. 10 to 12 wks
c. 14 to 16 wks
d. 18 to 20 wks

Explanation

Amniocentesis is an invasive, diagnostic antenatal test. It involves taking a sample of amniotic fluid in order to examine fetal cells found in this fluid. Amniocentesis is usually performed between 14 and 20 weeks. Some medical facilities may perform amniocentesis as early as 11 weeks. Amniocentesis can be used later in the third trimester for a few reasons.

www.medicalcoupons.in

180. Malignant hyperthermia is due to defect in ?

a. Ryanodine receptor
b. T- tubule defect
c. Na+/k+ ATP ase defect
d. Mutation of SERCA pump.

Explanation :


Malignant hyperthermia (MH) is a dominantly inherited disorder of skeletal muscle that leads to a life threatening adverse reaction upon exposure to potent volatile anesthetics (halothane, isoflurane, sevoflurane, desflurane etc) and the skeletal muscle relaxant succinylcholine. The anesthetic drugs trigger an uncontrolled calcium (Ca2+) release from the sarcoplasmic reticulum (SR) through the ryanodine receptor (RYR1) causing a rapid and sustained rise in myoplasmic Ca2+ which cause prolonged muscle contraction. DOC is dantrolene.

181. Identify the image given



a. Ulcus serpens
b. Acanthamoebal ulcer
c. Dentritic ulcer
d. Pseudodentritic ulcer

Explanation



Viral ulcer – caused by herpes simplex usually attacks the epithelium and stroma. In epithelium these dentritic ulcers are seen. These are called true dentrites and are like branch of tree. Corneal sensation is reduced and so it is regarded as most painless ulcer. Steroid are C/I in dentritic ulcer . If given, it can lead to geographical ulcer.

www.medicalcoupons.in

182. Delayed onset of puberty in female age is seen at?
   a. 12 yrs
   b. 13 yrs
   c. 14 yrs
   d. 15 yrs

Explanation :

Puberty starts when the pituitary gland begins to produce two hormones , luteinizing hormone (called LH) and follicle-stimulating hormone (called FSH), which cause the ovaries to enlarge and begin producing estrogens. The growth spurt starts shortly after breasts begin to develop, and the first menstrual cycle begins about 2-3 years later. A girl who has not started to have breast development by the age of 13 is considered to be delayed.

183. Microsporum affects ?
   a. Skin , hair and nail
   b. Skin and hair
   c. Skin and nail
   d. Hair and nail

Explanation :

Microsporum spp. mostly infect the hair and skin, except for Microsporum persicolor which does not infect hair.

184. Who level of Hb con... Of anemia in pregnancy is ?
   a. 10 gm/dl
   b. 11 gm/dl
   c. 12 gm/dl
   d. 13 gm/dl

Explanation :

Anemia is a condition of low circulating haemoglobin (Hb) in which concentration has fallen below a threshold lying at two standard deviations below the median of a healthy population of the same age, sex and stage of pregnancy. The WHO definition for diagnosis of anemia in pregnancy is a Hb concentration of less than 11 g/dl (7.45 mmol/L) and a hematocrit of less than 33%.

185. Radiological test for staging of endometrial carcinoma invasion is?
   a. MRI
   b. CECT pelvis
   c. TAS
   d. TVS

www.medicalcoupons.in

Explanation :

A dedicated MRI protocol is crucial for accurate MRI evaluation of endometrial carcinomas. In endometrial carcinoma, it is important to accurately assess invasion depth and preoperative staging. Fusion of T2-weighted magnetic resonance imaging (T2WI) and diffusion-weighted magnetic resonance imaging (DWI) may contribute to the improvement of anatomical localization of lesions.

186. For how long immunusuppresive therapy should be given post renal transplant?
   a. Lifelong
   b. 3 months
   c. 6 months
   d. 9 months

187. Organism causing whipple's disease ?
   a. Actinomycetes
   b. Giardia
   c. M. Chelonie
   d. M. Kansasii

Explanation :

T. whipplei is one of the actinomycetes, and is a distant relative of Mycobacterium avium-intracellulare, and Mycobacterium paratuberculosis explaining in part why Whipple's disease is similar to the diseases caused by these bacteria. The disease is common in farmers and those exposed to soil and animals, suggesting that the infection is acquired from these sources.

188. What is the ideal time to check for diabetic retinopathy in type 1 IDDM ?
   a. 1 yr
   b. 3 yr
   c. 5 yr
   d. 7 yr

Explanation :

For type – 1 DM , pt can wait for 5 yrs to check the retina

For type – 2 DM , it is necessary to check retina immediately as it is asymptomatic.

189. Most common cause of secondary amenorrhoea is?
   a. Turner syndrome
   b. Pregnancy
   c. Pelvic radiation
   d. Mullerian agenesis

www.medicalcoupons.in

Explanation :

Secondary amenhorrea is defined as a absence of mensuration of 3 normal cycles or for 6 months or more in a woman with previous normal mensural pattern in absence of pregnancy. Most common cause of secondary amenorrhea is pregnancy.

190. Floor of femoral triangle is formed by ?
   a. Inguinal ligament
   b. Adductor longus
   c. Pectineus muscle
   d. Sartorius muscle

Explanation :

Borders of femoral triangle :

Superior border – formed by inguinal ligament

Lateral border – formed by the medial border of Sartorius muscle

Medial border – formed by medial border of adductor longus muscle

Roof – formed by fascia lata

Floor – formed by pectineus, iliopsoas muscle, and adductor longus muscle.

191. Which of the following is not a sign of placental separation?
   a. Gush of blood flow
   b. Firm and globe shaped uterus
   c. Uterus rises in the abdomen
   d. Shortening of the length of umbilical cord

Explanation :

Childbirth involves three stages:

First stage: Begins from the onset of true labor and lasts until the cervix is completely dilated to 10 cm.

Second stage: Continues after the cervix is dilated to 10 cm until the delivery of your baby.

Third stage: Delivery of your placenta. The third stage is the delivery of the placenta and is the shortest stage. The time it takes to deliver your placenta can range from 5 to 30 minutes.

www.medicalcoupons.in

Signs of placental separation :

- A gush of blood comes from vagina
- Umbilical cord gets longer
- Uterus rises over the abdomen
- Uterus becomes round in shaped.

192. Most common site for the development of laryngeal web is ?
   a. Anterior
   b. Posterior
   c. Subglottic
   d. Supraglottic

Explanation :

Laryngeal web is a condition in which child's larynx (voice box) contains a layer of web-like tissue. This partially obstructs child's windpipe, causing frequent shortness of breath and other symptoms. Laryngeal webs occur in the glottic level and affect the vocal cords ™More than 90% of laryngeal webs webs are located anteriorly and extend toward the arytenoids. Other types include the posterior glottic web, causing interarytenoid vocal cord fixation; subglottic webs, which may occur with or without cricoid cartilage involvement and subglottic stenosis; and supraglottic webs.

193. Rise in JVP with inspiration is seen with all except ?
   a. Pericardial effusion
   b. Pleural effusion
   c. Constrictive pericarditis
   d. Cardiac tamponade

Explanation :

JVP will decrease during inspiration in the normal state. The JVP will rise during inspiration (kussmaul's sign)in the presence of pericardial constriction , right ventricular infarction , pericardial effusion, or rarely, cardiac tamponade.

194. Normal refractive power of neonates are ?
   a. Myopia
   b. Pseudomyopia
   c. Hypermetropia
   d. Anisometropes

Explanation :

www.medicalcoupons.in

The eye of the newborn is hypermetropic by 00+2 to +3D. A – P diameter of eyeball is 16.5 mm. Corneal diameter is 10mm. At 2 - 4 months there is a development of fixation reflex , and at 6 -8 months depth perception starts developing

195. Intraabdominal pressure during gynaecological procedure is ?
   a. 5 – 10 mg
   b. 10 – 20 mg
   c. 20 – 30 mg
   d. 40 – 50 mg

Explanation :

The deliberate induction of carbon dioxide pneumoperitoneum during laparoscopic surgery could be a possible source of cardiovascular collapse. An IAP from 8 to 12 mm Hg is recommended for laparoscopic surgery to avoid complications caused by hemodynamic derangements.

196. Which of the following is the marker for hepatocellular carcinoma ?
   a. Alpha feto protein
   b. Beta HCG
   c. TSH
   d. Thyroglobulin

Explanation :

Hepatocellular carcinoma (HCC) is one of the most frequent malignant tumors. Serum tumor markers, as the effective method for detecting hepatocellular carcinoma for a long time, could be divided into 4 categories: oncofetal antigens and glycoprotein antigens; enzymes and isoenzymes; genes; and cytokines. Serum alpha fetoprotein (AFP) is the most widely used tumor marker in detecting patients with hepatocellular carcinoma, and has been proven to have capability of prefiguring the prognosis.

197. Which of the following statement is false regarding gastrinoma?
   a. Associated with MEN 1 syndrome
   b. Most common presentation is Gastric outlet obstruction
   c. Mc site is duodenum
   d. It is relatively silent

Explanation :


Gastrinomas are neuroendocrine tumours that are usually found in the first part of the small intestine or in the pancreas, which produce the hormone gastrin. The signs and symptoms of gastrinomas that most patients experience are abdominal pain, heartburn, nausea, blood in the vomit, bleeding from the back passage and diarrhoea.

www.medicalcoupons.in

198. Damage to which of the following nerves lead to aphonia, aspiration, vocal cord paralysis?

a. inferior laryngeal nerve
b. superior laryngeal nerve
c. Internal laryngeal nerve
d. Recurrent laryngeal nerve

Explanation :



PARALYSIS OF SUPERIOR LARYNGEAL NERVE

(A) UNILATERAL

- Paralysis of cricothyroid muscle & ipsilateral anaesthesia of the larynx above the vocal cord.
- **Causes :**
  - Thyroid surgery
  - Thyroid Tumors
  - Diptheria.
- **Clinical features :**
  - Weak voice with decreased pitch
  - Anaesthesia of the larynx on one side
  - Occassional aspiration.

➢ **Laryngeal findings** include :

- **Askew position of glottis** - Ant. Comissure is rotated to healthy side.
- **Shortening of V.C.** with loss of tensi… V.C. appears wavy
- **Flapping of the paralysed vocal cord** - V.C. …gs down during inspiration & bulges u… during expiration.

(B) BILATERAL

- An uncommon condition …both … cricothyriod muscles are paralysed a…ong with …aesthesia of upper larynx.
- **Causes:**
  - Surgical or accide…al traum…
  - Diptheria
  - Cervical ly… hade…nathy
  - Neoplastic di…se
- **Clini…tures:**
  - Both …C. p…sis
  - Ana…the…ia o…ynx
  - Cough
  - …ockin…
  - Weak & husky voice

Treatm…t:

- **…eostomy** with a cuffed tube & an oesophageal feeeding tube.

Epiglottopexy is an operation to close the laryngeal inlet to protect the lungs from repeated aspiration. It is a reversible precedure.

199. Esmolol is a s…ort acting B- blocker because?

a. Bl…oo…esterase activity
b. …apid renal clearance
c. Rapi… hydrolysis
d. I…creased bioavailablity

Explana…tion :

Esm…ol (Brevibloc) is an intravenous, short-acting, titratable, cardioselective beta b…cke… with a very rapid onset and offset of action (t1/2 = 9.2 minutes). Esmolol-i…duce… beta blockade can be maintained as long as infusion is continued. It exhibits ne…ther intrinsic sympathomimetic activity nor significant membrane-stabilizing ac…vity. It is rapidly metabolized by an esterase in the erythrocyte cytosol to an …nactive acid metabolite. Its hemodynamic and electrophysiologic effects are similar to those of other beta blockers.

www.medicalcoupons.in

200. Route of spread of TB to kidney is through?

a. Lymphatic
b. Hematogenous
c. Both
d. None

Explanation :

Renal tuberculosis (renal TB) is a type of extra-pulmonary tuberculosis that affects the kidney. It is caused by *Mycobacterium tuberculosis*, which is a common cause of respiratory infection especially in the low socio-economic regions of the underdeveloped and developing countries. The kidneys can get infected through the spread of infection from other parts of the body especially the lungs via the blood. This type of infection of kidneys that spreads from other sites in the body that are already affected is referred to as secondary tuberculosis.

201. A 36 yr old man who was taking some drug develops a patch of redness and swelling of the skin. which of the following is possible diagnosis?

a. TEN
b. Fixed drug eruption
c. Pemphigus vulgaris
d. Steven Johnson syndrome

Explanation:

A fixed drug eruption is an allergic reaction to a medicine that characteristically recurs in the same site each time when a particular drug is taken. It presents as well defines, round or oval patches of redness and swelling of skin, sometimes surrounded by blisters. In mucosal sites, extensive ulceration can occur. The hands, feet, lips, eyelids, genitalia, and perianal regions are common sites.

202. Choice of contraception in patients with history of gestational diabetes ?

a. Combined contraceptive pills
b. DMPA
c. IUD
d. Barrier method

Explanation :

Women with prior gestational diabetes are at risk of developing overt diabetes predominantly type- 2 after pregnancy often during their reproductive cycles. Contraception is essential component to be able to plan pregnancy when glucose status is normalized and may also confer protection from developing diabetes by preventing a subsequent pregnancy. IUD is a very effective and reversible contraceptive method without metabolic disturbances and therefore is an ideal contraceptive for women with prior GDM.

www.medicalcoupons.in

203. Laboratory findings shows high PTH , high ALP, low calcium and low PO_4 Which of the following is the possible diagnosis of the patient?

a. Paget's disease of bone
b. Osteoporosis
c. Vit. D deficiency
d. Hypoparathyroidism

Explanation :

Among the patients with severe vit .D deficiency, prevalence of hyperparathyroidism is high (38%). Severe vit. D deficiency was associated with elevated levels of alkaline phosphatase, a marker for increased bone turnover.

Causes of Hypocalcemia

Causes	Serum concentrations of markers PTH	ALP	Phosphate	25-(OH)D
Commom				
Postsurgical hypoparathyroidism	L	N	H	N
Autoimmune hypyoparathyroidism	L	N	H	N
Vitamin D deficiency	H	N or H	L	L
Renal disease	H	N or H	H	N or L
Rare				
Parathyroid hormone resistance	H	N	H	N
Vitamin D resistance	H	N	L	N
Autosomal dominant hypocalcemia	N	N	N	
Hypomagnesemia	N or L	N	N	N or L
Sclerotic metastasis	H	H	L	N
Other				
Hungry bone syndrome, critical illness, intravenous bisphosphonate				

204. A 9 yr old school going girl has a difficulty in focusing on one task, day dreamer, easily distracted, talk nonstop. Possible diagnosis is ?

a. Conduct disorder
b. ADHD
c. AUTISM
d. Pervasive development disorder

Explanation :

ADHD is one of the most common conditions in childhood. It impacts how kids function in school and in everyday life. symptoms include :

- Difficulty paying attention (inattention)
- Being overactive (hyperactivity)
- Acting without thinking (impulsivity)

There is no cure for ADHD, currently available treatments may help reduce symptoms and improve functioning. ADHD is commonly treated with medication, education or training, therapy, or a combination of treatments.

www.medicalcoupons.in

205. False statement regarding peutz jegher syndrome is?
   a. Associated with mucosal, lip and skin melanosis
   b. It is autosomal disorder
   c. Has multiple polyp in intestine that require surgery to stop bleeding
   d. It is a malignant condition

Explanation :

Peutz-Jeghers syndrome (PJS) is an autosomal dominant inherited disorder characterized by benign intestinal hamartomatous polyps in association with a distinct pattern of skin and mucosal macular melanin deposition. People with Peutz-Jeghers syndrome have a high risk of developing cancer during their lifetimes. Cancers of the gastrointestinal tract, pancreas, cervix, ovary, and breast are among the most commonly reported

206. B – blocker ointment is used for which of the following conditions?
   a. Superficial burns
   b. Infantile hemangiomas
   c. Contaminated wounds
   d. Clean wounds

Explanation :

The treatment of infantile hemangiomas has changed from the use of oral corticosteroids to oral propranolol. Topical beta blockers are now used for lesions with both deep and superficial components and those that are amblyogenic. When initiated in the proliferative phase of the lesion, the effectiveness of the treatment can be seen within days.

207. NRHM tracking system is for ?
   a. Mother and child linkage
   b. Tracking leprosy cases by ASHA
   c. Tracking HIV patients
   d. Monitoring TB patients.

Explanation :

Mother and Child Tracking System (MCTS) is an initiative of Ministry of Health & Family Welfare (national rural health mission) for ensuring delivery of full spectrum of healthcare and immunization services to pregnant women and children up to 5 years of age. Generation of work plans of ANMs, sending regular alerts to the service providers as well as beneficiaries about the services and to monitor delivery of services will go a long way in ensuring quality service delivery, micro birth planning, ensuring universal immunization and will have positive impact on important health indicators like Infant Mortality Rate and Maternal Mortality Ratio.—

www.medicalcoupons.in

208. Equilibrium potential of chloride ions in the cell according to nerst equation?
   a. +60mv
   b. -70mv
   c. +80 mv
   d. – 90 mv

Explanation :

Nerst equation gives the equilibrium potential or isoelectric potential. Equilibrium potential is the membrane potential at which equilibrium is reached. ie, there is no net flux of that ion.

E Na+ = +60 mv (tendency of sodium is to diffuse in till potential reaches +60 mv)

E K+ = -90 mv tendency of potassium is ti diffuse out till potential reches -90mv)

E cl- = - 70 mv . This is the closest to the RMP

209. In which of the following eating disorder amenorrhoea is present?
   a. Anorexia nervosa
   b. Bullemia nervosa
   c. Schizophrenia
   d. Binge eating disorders

Explanation :

Anorexia nervosa	Bulimia nervosa
Mainly refers to starving on purpose to create a skeletal like body.	It refers to binge eating followed by purging, resulting in an average or over weight.
Very less eating	Attempts to counteract the effects of over eating by self induced vomiting.
Restrictive type is commoner	
Binge eating 25- 50%	Binge eating is required for diagnosis
May result in conditions such as Amenorrhoea, osteoporosis, infertility etc.	May result in heart failure and damage to esophagus and teeth.

www.medicalcoupons.in

210. Fetal biophysical profile doesn't include?
   a. Non- stress test
   b. Fetal breathing movements
   c. To estimate placental function
   d. Fetal tone


Explanation :

BIOPHYSICAL PROFILE (BPP)

- **Composes of 5 components**
  1. **Non stress test (NST)**
  2. **Amniotic Fluid Volume (vertical pocket of 2cm or more)**
  3. **Fetal breathing movements (30 secs or more in 30 mins)**
  4. **Fetal movements (3 or more in 30 mins)**
  5. **Fetal tone (Extension/Flexion of an extremity)**

- **Each carry a score of 2 points, a total of 8 or 10 is Normal, 6 is Equivocal, and 4 or less is abnormal**

211. Identify the organism in the slide image given below:



   a. Chlonorchis sinensis
   b. Schistosoma hematobium
   c. Fasciola hepatica
   d. Paragonimus westermani

www.medicalcoupons.in

Explanation :




Image courtesy: Gryseels B, et al., Human schistosomiasis, *Lancet* 006; 368 1106.

Schistosoma hematobium causes schistosomiasis or bilharziasis . Symptoms associated are allergic dermatitis due to cercariae known as swimmers itch . Symptoms due to eggs are egg granuloma and hematuria . It also causes bladder cancer.

212. In modern management techniques in which of the following analysis Benefits are measured in monetary terms ?

a. Cost effective analysis
b. Cost benefit analysis
c. System analysis
d. Network analysis

Explanation :

Cost benefit analysis .

It is a management technique where economic benefits of any programme are compared with cost of that programme. The benefits are expressed in monetary terms. The main draw back of this technique is that all benefits in field of health cannot be expressed in monetary terms.

213. Which of the following is a short acting synthetic opoid?

a. Loperamide
b. Buprenorpine
c. Remifentanil
d. Tramadol

Explanation :

Remifentanil is a potent ultra short-acting synthetic opioid analgesic drug. It is given to patients during surgery to relieve pain and as an adjunct to an anaesthetic.

www.medicalcoupons.in

Remifentanil is a specific mu-type-opioid receptor agonist. Hence, it causes a reduction in sympathetic nervous system tone, respiratory depression and analgesia.

214. True statement regarding erythropoietin is ?
   a. Senses hypoxia
   b. Transfers oxygen to tissues
   c. Stimulates bone marrow
   d. Mainly produced by liver

Explanation :

Erythropoietin (EPO) is a hormone produced by the kidney that promotes the formation of red blood cells by the bone marrow. Erythropoietin stimulates the bone marrow to produce more red blood cells. The resulting rise in red cells increases the oxygen-carrying capacity of the blood. Only about 10% of erythropoietin is produced in the liver.

215. Common site of carcinoid tumour is ?
   a. Appendix
   b. Ileum
   c. Rectum
   d. Lungs

Explanation :

Carcinoid tumors arise from neuroendocrine cells and may develop in almost any organ. The gastrointestinal tract accounted for 54.5% of the tumors. Within the gastrointestinal tract, the small intestine was the most common site (44.7%), followed by the rectum (19.6%), appendix (16.7%), colon (10.6%), and stomach (7.2%).

216. Which among the following cell is activated form of macrophage?
   a. Stellate cells
   b. Langerhan cells
   c. Giant cell
   d. Epitheloid cells

Explanation :

Epithelioid cells are cells of the mononuclear phagocyte system found in certain granulomas mainly associated with intense immunological activity. These cells show little phagocytic activity.

217. Function of SGLT 1 is to transport ?
   a. Sodium
   b. Glucose
   c. Potassium
   d. Bicarbonate

www.medicalcoupons.in

Explanation :

SGLT -1 , is a high-affinity Na+/glucose co-transporter . SGLT1 transports glucose and galactose across the luminal (gut) side of enterocytes, and is the first step in the absorption of sugars from nutrients. In the kidney, SGLT1 is located on the apical (urine) side of the proximal tubule, and facilitates the reabsorption of urinary glucose from the glomerular filtrate.

218. Hypervirulent klebsiella pneumonia causes?
   a. URTI
   b. Conjunctivitis
   c. endopthalmitis
   d. Bronchitis

Explanation :

Hypervirulent *Klebsiella pneumonia* (hvKP) infection have been reported worldwide . Such strains are notorious for their capacity to cause serious and metastatic infections in young and healthy individuals, such as pyogenic liver abscesses and endophthalmitis.

219. Most common site of Puerperal infection is?
   a. Uterus
   b. Placenta
   c. Vagina
   d. Umbilical cord

Explanation :

Puerperium is a period after child birth. It starts after delivery and extends 6-8 wks after child birth.

Puerperal pyrexia – rise of temp reaching 100.4 ‘ F (38’C) on separate occations at 24 hrs apart within 1st 10 days following delivery.

Most common site of puerperal infection – placental site.

Most common manifestation of puerperal infection – endometritis

Most common cause of puerperal sepsis – streptococcus

Most common route of infection – direct spread

220. Main function of DNA POLYMERASE 3 is?
   a. To initiate replication
   b. To fill the gap after removing primer
   c. Helps in proof reading
   d. Endonuclease activity

www.medicalcoupons.in

Explanation :

DNA polymerase 3 is responsible for replication but it doesn't start the process of replication. It will read the parent DNA from 3' to 5'. Specific feature of DNA polymerase 3 is to

1) Proof reading /rechecking
2) Exonulease activity

DNA polymerase 1 helps to fill the gap once after removing the primer by RNAse H.

221. Most common site of innocent murmur is ?
   a. Apex
   b. Left lower sternal border
   c. Above clavicles
   d. 2nd rt intercoastal space

Explanation :

Still's murmur is by far and away the most common innocent heart murmur. These signs are unreliable in under age 1 year due to higher Incidence of Congenital Heart Disease.

Common Innocent Heart Murmurs

Type (Timing)	Description of murmur	Age group
Classic Vibratory Murmur (Still Murmur) Systolic	Maximal at MLSB or between LL and apex Low frequency vibratory, twanging, groaning, squeaking murmur	3-6 year Occasionally in infancy
Pulmonary ejection murmur (Systolic)	Maximal at ULSB, Early to mid-systolic, Grade 1-3/6 in intensity. Blowing in quality	8 – 14 year
Pulmonary flow murmur	Maximal at Transmits well to left and right chest, axilla and back. Grade 1-2/6 in intensity	Premature and full-term newborns Usually disappears by 3 – 6 months of age
Venous hum (Continuous)	Maximal at (over left) supraclavicular or infraclavicular areas. Grade 1-3/6 in intensity. Inaudible in supine position. Intensity changes with rotation of head and compression of jugular v	3 – 6 year
Carotid bruit (Systolic)	Right supraclavicular area and over carotid. Grade 2-3/6 in intensity. Occasional thrill over carotid	Any age

222. Afferents of cremasteric reflex are present in ?
   a. Genitofemoral
   b. Iliohypogastric
   c. Ilioinguinal
   d. Pudendal

Explanation :

The cremasteric reflex is dependent upon the nerve roots L1 and L2. Genitofemoral nerve carries both the afferent and efferent components of reflex. This

www.medicalcoupons.in

reflex is elicited by stroking the superior and medial part of the thigh in a downward direction. The normal response in males is a contraction of the cremaster muscle that pulls up the scrotum and testis on the side stroked. In children, this reflex may be over exaggerated and this can lead to mistaken of diagnosis of undescented testis. Upper and lower motor neuron disorders can cause an absence of cremasteric reflex.

223. Chance of metastasis to liver in colon cancer is?
   a. Less than 25%
   b. 25- 50%
   c. 75%
   d. 90 %

Explanation:

More than 50% of the patients with colorectal cancer (CRC) have or will develop metastasis, with a quarter having distant metastatic disease at the time of diagnosis, most frequently in the liver. Liver metastasis is the leading cause of cancer-related morbidity and mortality in colorectal cancer. Left sided colon cancer has 70% increased risk of isolated liver metastases. The only potentially curative treatment for liver metastasis is liver resection but only 15% to 20% of the patients are suitable for surgical resection.

224. False statement regarding crohn's disease is?
   a. Skip lesions are seen
   b. Presence of non caseating granuloma
   c. Intestinal resection is curative
   d. Ileum is the most common site to be affected.

Explanation :

COMPARISON OF CROHN DISEASE VERSUS ULCERATIVE COLITIS


	CROHN DISEASE	ULCERATIVE COLITIS
Site of origin	Terminal ileum	Rectum
Pattern of progression	"Skip" lesions/irregular	Proximally contiguous
Thickness of inflammation	Transmural	Submucosa or mucosa
Symptoms	Crampy abdominal pain	Bloody diarrhea
Complications	Fistulas, abscess, obstruction	Hemorrhage, toxic megacolon
Radiographic findings	String sign on barium X-ray	Lead pipe colon on barium X-ray
Risk of colon cancer	Slight increase	Marked increase
Surgery	For complications such as stricture	Curative

www.medicalcoupons.in

225. Which of the following is not a absolute contraindication of OCP?

a. Breast cancer
b. Liver disease
c. Age <30 yrs
d. Congenital hyperlipdemia

Explanation :



226. Which of the following is not a time resistant strategy ?

a. Aim
b. Goal
c. Objective
d. Mission

Explanation :

Goal : ultimate desired state towards which objectives and resources are directed .It is not constrained by time or exisiting recources and shows “all or none phenomenon”.

Objective : is planned en – point of all activities . It is concerned with problems itself.

Mission : is a disruption of fundamental principle of existence of a programme. It is usually time bound.

Aim/ Target : a discrete activity which helps measure extent of attainment of objectives. It is a concept of degree of achievement.

www.medicalcoupons.in

227. A 70 yr old patient underwent TURP for BPH under spinal anaesthesia. One hr later he develops vomiting and altered sensorium. The most possible cause is ?

a. Rupture of bladder
b. Hyperkalemia
c. Over dosage of spinal anaesthetic agent
d. Water intoxication

228. Weight of respired lung to total body weight is?

a. 1/35th
b. 1/70th
c. 1/50th
d. 1/90th

Explanation :

Character	Unrespired Lungs	Respired Lungs
Weight in relation to body weight	1/70	1/35
Volume	Small	Large and covers the heart
Consistency	Liver Like: Dense firm not crepitant	Soft, spongy, elastic, Crepitant
Extension	Up to the level of 4th and 5th rib	Up to the level of 6th and 7th rib
Specific gravity	1.04	0.94
Margin	Sharp	Rounded
color	Uniform reddish	Mottled/ marbled appearance
Air vesicle	Not inflated	Inflated
Section	Little froth less blood exudates on pressure	Abundant frothy blood exudates
Breslow life test	Whole or part sinks	Expanded

229. Which of the following statement is correct regarding fast breathing in a child?

a. > 60 /min in a child > 12 months
b. >50 /min in a child of age between 2 months to 1 yr
c. > 40/min in a child of > 2 months
d. > 70 /min in a child of > 18 months

www.medicalcoupons.in

Explanation :

According to ARI control programme, a case of pneumonia is considered only if child has fast breathing.

Upto 2 months - >60/min

2- 12 months - > 50/min

>12 months - >40/min

230. Which of the following hormone is increased in ZES?
   a. Histamine
   b. Pepsin
   c. Gastrin
   d. Secretin

Explanation :

Zollinger-Ellison syndrome is a rare disorder that occurs when one or more tumors form in the pancreas and duodenum. The tumors, called gastrinomas, release large amounts of gastrin that cause the stomach to produce large amounts of acid. Patients with ZES show gastric acid hypersecretion, which induces refractory peptic ulcers, severe gastroesophageal reflux disease, diarrhea, and even death. The plasma concentration of gastrin is measured by RIA or ELISA , and can be used to diagnose gastrin-secreting tumors, atrophic gastritis, gastric ulcers, and pernicious anemia. Zollinger-Ellison syndrome treated with medications to reduce gastric acid secretion and with surgery to remove gastrinomas.

231. Febuxostat is used in the treatment of?
   a. Gout
   b. Rheumatoid arthritis
   c. Polyarteritis nodosa
   d. giant cell arteritis

Explanation

Febuxostat is used to lower uric acid levels in people with gout. It inhibits xanthine oxidase . Febuxostat works by reducing the amount of uric acid made by the body. Increased uric acid levels can cause gout.

232. Dense bodies in smooth muscle is analogues to which part of skeletal muscle?
   a. I band
   b. M line
   c. Z line
   d. A band

www.medicalcoupons.in

Explanation :

Smooth muscle fibers are spindle-shaped and have a single nucleus; they range from about 30 to 200 μm and they produce their own connective tissue endomysium. Although they do not have striations and sarcomeres, smooth muscle fibers do have actin and myosin contractile proteins, and thick and thin filaments. These thin filaments are anchored by dense bodies. A dense body is analogous to the Z-discs of skeletal and cardiac muscle fibers and is fastened to the sarcolemma.

233. Excessive vomiting is an example of which type of acid base disorder?
   a. Metabolic acidosis
   b. Repiratory acidosis
   c. Metabolic alkalosis
   d. Respiratory alkalosis

Explanation:

In a case of vomiting , H+ ions are lost and because of this, bicarbonate ions remains and the patient develops metabolic alkalosis.

234. Under mental illness act, detaining of mentally ill person is punishable for how many yrs?
   a. 1 yr
   b. 2 yr
   c. 5 ys
   d. 7 yr

Explanation :

The mental heath act (1987) repels the Indian lunacy act (1912).

Defines, “mentally ill person” as a person who is in need of treatment by reason of any mental disorder other than mental retardation for treatment as in patient a person can represent himself or represented by family or friends in writing to the director of a psychiatric hospital. For a preliminary diagnosis , he can be kept in hospital for 10 days, if more time is required then it has to permit by magistrate who can give extension of 10 days twice.

Punishment : detains or keeps mentally ill patient otherwise in accordance with the act is punishable with upto 2 yrs imprisonment or Rs.10000 fine or both.

235. In establishing causal association, most essential criterion is?
   a. Consistency of association
   b. Temporal association
   c. Duration of association
   d. Strength of association

www.medicalcoupons.in

Explanation :

The Bradford-Hill criteria are widely used in epidemiology as providing a framework against which to assess whether an observed association is likely to be causal.

1. Strength of the association.
According to Hill, the stronger the association between a risk factor and outcome, the more likely the relationship is to be causal.

2. Consistency of findings.
Have the same findings must be observed among different populations, in different study designs and different times?

3. Specificity of the association.
There must be a one to one relationship between cause and outcome.

4. Temporal sequence of association.
Exposure must precede outcome.

5. Biological gradient.
Change in disease rates should follow from corresponding changes in exposure (dose-response).

6. Biological plausibility.
Presence of a potential biological mechanism.

7. Coherence.
Does the relationship agree with the current knowledge of the natural history/biology of the disease?

8. Experiment.
Does the removal of the exposure alter the frequency of the outcome?

236. Which of the following is the most common form of tularaemia?
   a. Fever
   b. Lymphadenopathy
   c. Ulcer
   d. Chills

Explanation :

Ulceroglandular - This is the most common form of tularemia and usually occurs following a tick or deer fly bite or after handing of an infected animal.

Glandular - Similar to ulceroglandular tularemia but without an ulcer.

Oculoglandular - This form occurs when the bacteria enter through the eye. This can occur when a person is butchering an infected animal and touches his or her

www.medicalcoupons.in


eyes. Symptoms include irritation and inflammation of the eye and swelling of lymph glands in front of the ear.

Oropharyngeal - This form results from eating or drinking contaminated food or water.

Pneumonic - This is the most serious form of tularemia. Symptoms include cough, chest pain, and difficulty breathing. This form results from breathing dusts or aerosols containing the organism. It can also occur when other forms of tularemia (e.g. ulceroglandular) are left untreated and the bacteria spread through the bloodstream to the lungs.

237. Nyhas classification of direct inguinal hernia is ?
   a. Type - 1
   b. Type - 2
   c. Type - 3 a
   d. Type – 3 b

Explanation :



Nyhus (1991) based the classification on the integrity of the posterior wall and the deep inguinal ring and grouped them into four types:

Type 1: Indirect inguinal hernia with a normal inguinal ring, peritoneal sac is in the inguinal canal

Type 2: Indirect hernia with enlarged deep inguinal ring with the posterior wall intact, sac not in scrotum.

Type 3: Posterior wall (floor) defects:

Type 3a: Direct hernia with a posterior floor defect only

Type 3b: Indirect hernia with enlargement of deep inguinal ring and posterior floor defect

Type 3c: Femoral hernia

Type 4: Recurrent hernia

A: Direct. B: Indirect. C: Femoral. D: Combinations of A-B and C.

238. Which of the following statement is true regarding intermittent claudication ?
   a. Pain starts even at rest
   b. Pain begins when the patient starts to walk
   c. Pain relieved after few mins of walking
   d. Patient was unable to walk

www.medicalcoupons.in

Explanation :

Claudication is pain and/or cramping in the lower leg due to inadequate blood flow to the muscles. It is commonly referred to as "intermittent" claudication because it comes and goes with exertion and rest. In severe claudication, the pain is also felt at rest. Intermittent claudication can be due to temporary artery narrowing due to vasospasm (spasm of the artery), permanent artery narrowing due to atherosclerosis, or complete occlusion (closure) of an artery to the leg. Diagnostic tests include blood pressure measurements to compare the arms and legs, Doppler ultrasonography on the legs, duplex Doppler/ultrasound exam of the extremities to visualize arterial blood flow, an ECG, and arteriography. The prognosis with intermittent claudication is generally favorable because the condition often stabilizes or improves in time. Conservative therapy is advisable.

239. Which of the following is a malignant cerebellar tumour in children?
    a. Craniopharyngioma
    b. Ependymoma
    c. Medulloblastoma
    d. Oligodentroglioma

Explanation :

Medulloblastoma is a fast-growing, high grade tumor located in the cerebellum—the lower, rear portion of the brain. It is unusual for medulloblastomas to spread outside the brain and spinal cord. the exact cause of medulloblastoma is not known .The most common symptoms of medulloblastoma include behavioral changes, changes in appetite, symptoms of increased pressure on the brain (eg, headache, nausea, vomiting, and drowsiness as well as problems with coordination). Unusual eye movements may also occur. Treatment consists of surgical removal of as much tumor as possible, radiation, and then chemotherapy.

240. A patient was brought to the hospital with a history of RTA 8 hrs back. A few drops of blood were noted at the external urethral meatus. He had not passed urine and his bladder palpable per abdomen. The probable diagnosis is.?
    a. Rupture bladder
    b. Urethral injury
    c. Anuria due to hypovolemia
    d. Extravasation of urine in the retroperitoneum due to fracture pelvis

Explanation :

Membranous urethra is injured in this case. Common cause is due to fractured pelvis due to trauma or RTA. Blood is seen at urethral meatus. Investigation of choice is RGU. Managed with supra pubic catheter and drain the urine.

www.medicalcoupons.in

241. Thymidylate synthase inhibitor is?
   a. Cladribine
   b. Lomustine
   c. 5 – flurouracil
   d. Paclitaxel

Explanation :

5 – flurouracil inhibits thymidine synthase. Used for – head and neck, breast , and ovarian cancers. It is also used topically for basal cell cancer and keratosis. It causes single strand breaks affecting the action of both DNA and RNA. S/E – bone marrow suppression.

242. For 100 mt sprint energy/ATP that are required is from?
   a. Muscle
   b. Liver
   c. Blood
   d. Lungs

Explanation :

The 100-meter sprint requires intense, all-out effort for a very short time. Cells break down carbohydrates and fats to provide energy, then store that energy in the form of adenosine triphosphate (ATP). ATP is often termed the cell's energy currency. Because the breakdown of stored ATP releases energy almost instantaneously, & it is the primary energy source as a sprinter explodes out of the blocks. However, muscles only store enough ATP for two to three seconds of maximal power output. To rapidly replenish ATP, muscle cells break down another high-energy compound called creatine phosphate (CP). This is called the ATP-CP or phosphagen energy system, sometimes referred to as the alactic anaerobic system because it does not require oxygen.

243. In which of the following conditions large ovaries is not seen?
   a. PCOS
   b. Mucinous cystadenoma
   c. Turner syndrome
   d. Fibroma

www.medicalcoupons.in

Explanation :

Turner syndrome – genotype- 45XO

Normal pathway of development is female soin the absence of y chromosomes the fetus becomes female but for normal ovarian development 2X chromosomes are required. Hence the presence of only one chromosome will make the gonads streak like. It is the most common cause of primary amenorrhoea and delayed puberty in females.

244. Moderately increased risk of invasive breast cancer is associated with which of the following?
   a. Fibroadenoma
   b. Ductal ectasia
   c. Apocrine metaplsia
   d. Atypical ductal hyperplasia

Explanation :

Of the epithelial hyperplasias, only florid hyperplasia and atypical hyperplasia are associated with increased risk of breast cancer.

245. Which type of Anemia seen in rheumatoid arthiritis is ?
   a. Haemolytic nemia
   b. Aplastic anemia
   c. Anemia of chronic disease
   d. Sideroblastic anemia

Explanation :

RA can be associated with different types of anemia, including anemia of chronic inflammation and iron deficiency anemia. Chronic inflammation can lower the production of red blood cells in your bone marrow. This can lead to the release of certain proteins that affect how the body uses iron. Inflammation can affect the way the body produces erythropoietin, a hormone that controls the production of red blood cells.

246. Which of the following statement is true regarding aminocaproic acid?
   a. Is a irreversible inhibitor of plasminogen
   b. Is a derivative and analogue of amino acid arginine.
   c. Used as a fibrinolytic agent
   d. C/I in bleeding disorders

www.medicalcoupons.in

Explanation :

Overdose of fibrinolytic agent is treated by EPSILON AMINOCAPROIC ACID (EACA) & TRANEXEMIC ACID. EACA is a derivative and analogue of the amino acid lysine, and effective inhibitor of plasminogen. EACA has been used in numerous clinical situations to control bleeding and has been claimed to be an effective agent in subarachnoid hemorrhage, genitourinary bleeding from many causes, and in dental surgery in hemophiliacs. It may also be effective in several less well studied situations such as prophylaxis of bleeding episodes in hemophiliacs, control of menorrhagia, gastrointestinal bleeding, obstetrical bleeding and in bleeding following cardiac and thoracic surgery. Major side effects from EACA include hypotension, cardiac arrhythmias, rhabdomyolysis, and generation of thrombi.

247. "PAD sign " is a feature of which disease?

a. **Ca. Head of pancreas**
b. Pyloric stenosis
c. Chronic pancreatitis
d. Duodenal peptic ulcer

Explanation :

Pancreatic calcifications may be seen concurrently in approximately in 2% of patients who have chronic pancreatitis complicated by pancreatic carcinoma. Upper GI barium studies reveal an extrinsic impression of the mass on the posteroinferior aspect of the antrum of stomach. This is known as "Antral Pad sign".

248. Correct Sequence of hematopiesis is ?

a. Placenta >liver > yolk sac
b. Placenta> bone marrow> yolk sac
c. Yolk sac > bone marrow> liver
d. **Yolk sac > liver> bone marrow**

Explanation :

Erythropoiesis begins in the yolk sac from 3rd week of intrauterine life.

From 3rd month – liver

From 5th month – bone marrow

www.medicalcoupons.in

249. Which of the following is not associated with pain in sub mucosal fibroids?
   a. Torsion
   b. Large size
   c. Associated with endometriosis
   d. Degeneration

Explanation :

A patient with fibroid usually doesn't complain pain. If a fibroid patient presenting with pain it deserves to be evaluated against :

- Malignancy
- It is being extruded from body as a polyp
- Associated endometriosis
- Torsion of a pedunculated fibroma
- Degeneration

250. Most common cause of pulsatile proptosis is?
   a. craniopharyngioma
   b. Cavernous fistula
   c. Prolactinoma
   d. Meningitis

Explanation :

Proptosis – forward displacement or bulging of eyeball. Usually caused by carotid-cavernous fistula which may occur after trauma and cause arterial blood to get into the superior or inferior ophthalmic veins that empty into the sinus. In addition to pulsatile proptosis may also get palsy of CN 6 (most common) or CN 3 or 4 less commonly, along with loss of vision.

251. Type of liver disease seen in kwashiorkor is ?
   a. Fatty liver
   b. Cirrhotic liver
   c. Hepatitis
   d. Liver abcess

Explanation :

Kwashiorkor, also called protein malnutrition, condition caused by severe protein deficiency. It is common in young children weaned to a diet consisting chiefly of cereal grains, cassava, plantain, and sweet potato or similar starchy foods. The latter term comes from the reddish orange discoloration of the hair that is characteristic of the disease. Other symptoms include dry skin and skin rash, potbelly and edema, weakness, nervous irritability, anemia, digestive disturbances such as diarrhea, and fatty infiltration of the liver.

www.medicalcoupons.in

252. Opportunistic screening for HTN & DM starts at the age of ?

a. 30 yrs
b. 40 yrs
c. 50 yrs
d. 60 yrs

Explanation :

As per the guidelines issued by the National Program for Prevention and Control of Cancer, Diabetes, Cardiovascular disease and Stroke, opportunistic screening for hypertension and diabetes in ≥30 years of age is to be done by the health workers at the subcenter level (the most peripheral health center) and referred to the Community Health Centre (CHC).

253. Immediate effect of insulin on DKA type - 1 DM patient is?

a. High bicarbonate
b. Glycosuria
c. Hyperkalemia
d. Increased PH

254. Which statement is false about carcinoid tumour?

a. Presenting with flushing, wheeze, diarrhoea
b. Most common site ileum
c. Conservative management is curative
d. Present at multiple sites

Explanation :

Carcinoid tumors are of neuroendocrine origin and derived from primitive stem cells in the gut wall, but they can be seen in other organs, including the lungs, mediastinum, thymus, liver, pancreas, bronchus, ovaries, prostate, and kidneys. The etiology of carcinoid tumors is not known, but genetic abnormalities are suspected.

Sometimes these tumours can produce hormones, which can cause a collection of symptoms known as ‘carcinoid syndrome'. This can include:

- flushing
- severe diarrhoea
- loss of appetite
- weight loss
- increased heart rate
- wheezing

www.medicalcoupons.in

- sudden changes in your blood pressure.

Surgery, if feasible, is the only curative therapy. If the tumor has metastasized (most commonly, to the liver) and is considered incurable, there are some promising treatment modalities, such as radiolabeled octreotide or the radiopharmaceutical for arresting the growth of the tumors and prolonging survival in patients with liver metastases.

255. Gynecomastia in a newborn is due to the effect of which hormone?
   a. Maternal Estrogen
   b. Fetal estrogen
   c. B – HCG
   d. AFP

Explanation :

Neonatal gynecomastia is caused by the passage of maternal hormones (both estrogen and progesterone) through the placenta during pregnancy. In infants the breast enlargement usually progresses during the first 2 months of life. Also sometimes is observed secretion of the breasts. The breasts should not be red, hot and soft in neonatal gynecomastia. Diagnosis is made by physical examination , there are laboratory tests that are needed. No **treatment** is necessary because the mother's hormones are out of the baby's bloodstream , so will disappear from the breast tissue.

256. Important Step at waste management in hospital is?
   a. Segregation
   b. Transportation
   c. Dumping
   d. Generation

Explanation :

Segregation is one of the most important steps to successfully manage HCW. Segregating hazardous from nonhazardous waste also reduces the great risks of infecting workers handling HCW. Actually, the part of the HCW that is hazardous and requires special treatment could be reduced to some 2-5% if the hazardous part were immediately separated from the other waste. About 10-25% of the HCW is hazardous, treatment and disposal costs could be greatly reduced if a proper segregation were performed.

www.medicalcoupons.in

257. Causative agent of recurrent juvenile laryngeal papillatosis is?
   a. EBV
   b. Respiratory syncytial virus
   c. HPV 6 and 11
   d. Adenovirus

Explanation :

Age group affected – 2- 4 yrs. Caused by HPV serotype of 6 and 11. It is multiple in nature. Symptoms include hoarseness of voice. Managed with co2 lazer therapy. Serotype 11 is more aggressive form which may transform to malignancy.

258. P53 inhibitor acts in which phase of cell cycle?
   a. G1 phase
   b. G1- S phase
   c. M phase
   d. S – G2 phase

Explanation :

P53 is tumour suppressor gene & most gene mutated in most human cancers. P53 is also known as the "Guardian of Genome". P53 is a check point at both G1 – S phase and G2 – M phase.

259. Schilling's test is positive for which of the following conditions?
   a. Iron deficiency anemia
   b. Thalasemia
   c. AML
   d. Vit b12 deficiency

Explanation :

The Schilling test is used to diagnose the site of the disorder causing cobalamin deficiency. The Schilling test is performed to evaluate vitamin B_{12}absorption. Ingested vitamin B_{12} combines with intrinsic factor (produced in the stomach) and is absorbed in the distal ileum.

260. Best parameter to determine the fetal age in 1st trimester is?
   a. Crown rump length
   b. LMP
   c. Level of AFP
   d. Gestational sac mean diameter

Explanation :

Estimation of the gestational age by sonographic measurements of fetal parameters is usually done by measuring mean sac diameter (MSD), Crown-rump length (CRL), biparietal diameter (BPD), head circumference (HC), femoral length (FL) and

abdominal circumference (AC) depending on the trimester of pregnancy. MSD and CRL in 1st trimester and BPD, HC, FL and AC in 2nd and 3rd trimester. Gestational age was calculated by MSD using Rempen normogram and Hadlock normograms were used for the rest of fetal parameters. Gestational age by LMP was also calculated at the same time. Correlation of different fetal parameters in estimating gestational age in different trimesters was done by Pearson correlation. Pearson correlation showed that the CRL was the best fetal parameter (correlation coefficient of 0.909) in the first trimester.

261. All are true about gastric carcinoma except?
   a. H.pylori infection increase risks
   b. Most common site is at fundus
   c. Vit.C protects
   d. More in low socioeconomic group

Explanation :

Stomach cancer usually begins in the mucus producing cells that line the stomach. This type of cancer is called adenocarcinoma. The most common site of gastric cancer is antrum of stomach. Symptoms include postprandial heaviness in the abdomen (earliest symptom), weight loss (late symptom), vomiting and anorexia.IOC is endoscopy with biopsy and brush cytology.mets most commonly to liver, followed by lungs, bone, ovary.

262. Gastric lavage is not indicated in all cases of acute poisoning ideally because of?
   a. Danger of cardiac arrest
   b. Danger of respiratory arrest
   c. Fear of aspiration
   d. Inadequate ventilation

Explanation :

The main danger of consuming a poison, other than reducing the effects of poison itself on the body due to absorption, is death due to aspiration pneumonitis. It is because of this that gastric lavage is not indicated in all cases of acute poisoning.

263. ...m of an enzyme is?
   a. The substrate concentration that produces half maximum velocity
   b. The normal physiological substrate concentration
   c. Numerically identical for all isozymes that catalyze a given reaction
   d. Dissociation constant

Explanation :

"The substrate concentration that produces half maximal velocity (v termed the K value or Michaelis constant"

www.medicalcoupons.in

264. Unmet need of family planning are all except?

a. Women who are sexually active but not using any methods of contraception
b. Women who doesn't want to have any children in future
c. Women not using contraception but want to have children in future
d. Women who are sexually inactive

Explanation :


Unmet need for family planning is an important indicator for assessing the potential demand for family planning services. Currently married women who are not using any method of contraception but who do not want any more children are defined as having an unmet need for limiting and those who are not using contraception but want to wait two or more years before having another child are defined as having an unmet need for spacing.

265. A infant with respiratory distress, excessive drooling of saliva and inability to swallow. On physical examination the child abdomen was distended. Chest x- ray reveals tracheal compression and deviation & Absence of a gastric air bubble. What is the possible diagnosis?

a. ARDS
b. Esophageal atresia with tracheoesophageal fistula
c. Spontaneous pneumothorax
d. Achalasia

Explanation :

Tracheoesophageal fistula (TEF) is a developmental anomaly characterized by an abnormal connection between the trachea and the esophagus that usually accompanies esophageal atresia and in which the esophagus is closed off at some point. Abnormal development of the septum during this time can lead to TEF.



www.medicalcoupons.in

The most common abnormality is type C TEF with esophageal atresia, in which the upper section of the esophagus terminates in a blind pouch, and the lower section ascends from the stomach and connects with the trachea by a short fistulous tract. Abdominal X-rays may show the absence of air in the stomach and lower GI tract (types A and B) or distention (types C and D). Esophagoscopy identifies strictures or fistulas in the upper pouch; bronchoscopy confirms the presence of a fistula. Upright positioning
Treatment - Suction catheter placement to reduce secretions
Diet - Nothing by mouth (NPO)
Gradual introduction of oral feedings postoperatively, initially via a nasoduodenal tube and then orally after anastomoses heal.

266. True statement about frank breech is ?
   a. Most commonly associated with congenital anomaly
   b. Has hip extended and knee flexed
   c. ECV done before 34 weeks
   d. Has hip flexed and knee extended

Explanation :

There are three types of breech presentation: complete, incomplete, and frank.

Complete breech is when both of the baby's knees are bent and his feet and bottom are closest to the birth canal.

Incomplete breech is when one of the baby's knees is bent and his foot and bottom are closest to the birth canal.

Frank breech is when the baby's legs are folded flat up against his head and his bottom is closest to the birth canal.

267. False statement about aminoglycoside is?
   a. Used to treat gram +ve bacterial infection
   b. Acts by inhibiting protein synthesis
   c. kidney failure and hearing loss are the most common side effects
   d. Poorly absorbed by oral intake

Explanation :

Aminoglycosides are active against most gram-negative aerobic and facultative anaerobic bacilli but lack activity against anaerobes and most gram-positive bacteria, except for most staphylococci. Aminoglycosides are thought to work by inhibiting protein synthesis inside bacteria. Impairment of kidney function and hearing loss are the most common side effects of aminoglycosides. Aminoglycosides are not well

www.medicalcoupons.in

absorbed when given by mouth, so need to be given in injection by IV. Eg tobramycin , gentamicin, amikacin, Streptomycin, neomycin, and kanamycin.

268. Migratory motor complex of GIT is seen in which condition?
   a. Full fed state
   b. Fasting
   c. While eating
   d. 3 hrs after food

Explanation :

Migratory motor complex are waves of electrical activity that sweep through the intestines in a regular cycle during fasting. An increase in gastric, biliary and pancreatic secretion is also seen in conjunction with the motor activity. The periodic nature of the migrating motor complex is thought to be controlled from the central nervous system and may be implemented in part by the enteric hormone motilin. The MMC occurs every 45–180 minutes during the interdigestive phase (i.e., between meals) and is responsible for the rumbling experienced when hungry.

269. For DNA fingerprinting blood collected to be in which of the following?
   a. Glass bottles
   b. Syringes containing Na F
   c. EDTA vials
   d. Testubes with Na C

Explanation :

DNA fingerprinting is a technique, by which an individual can be identified at molecular level. Scientists have chosen repeating sequences in the DNA, which are present in all individuals on different chromosomes, and are known to vary from individual to individual except in identical twins. These are used as genetic markers to identify the individual. Any biological material such as a drop of blood, saliva, semen, and any body part such as bones, tissue, skull, teeth, hair with root etc can be used as specimen for DNA fingerprinting. Blood samples (2-3 ml) can be collected in the sterile blood collection material (EDTA vials)

270. False statement about fibroadenoma of breast is?
   a. It increases a risk of malignancy
   b. Surgery in a treatment of choice if size is > 5cm
   c. Surgery done on patients request
   d. Popcorn calcification is seen

Explanation :

A fibroadenoma is a noncancerous tumor in the breast that's commonly found in women under the age of 30. The tumor consists of breast tissue and stromal, or connective, tissue. Fibroadenomas can occur in one or both breasts.

www.medicalcoupons.in

271. Confirmatory test for Diagnosis of gout is ?
   a. Urine analysis
   b. Blood test
   c. Synovial fluid analysis
   d. X- ray of joints

Explanation :

Diagnosis of gout :

Synovial fluid examinations involve taking fluid samples from a joint through a needle and examining them under a microscope for urate crystals. This test can confirm the diagnosis but isn't always practical – it can be difficult and sometimes uncomfortable to draw fluid from a small joint such as the big toe.


A **blood test** can measure the amount of urate in your blood. The critical serum level of urate (the saturation point) is around 360 µmol/L. A raised level of urate strongly supports a diagnosis of gout but can't confirm it – not everyone with a raised level of urate will develop crystals in their joints, and it's possible for urate levels in the blood to be normal at the time of an acute attack.

X-rays of joints will reveal joint damage if you have long-standing and poorly controlled gout. However, x-rays are rarely helpful in confirming the diagnosis because they're usually normal in the early years of having gout.

272. Which of the following is not a cause of haemolytic anemia in newborn?
   a. Enzyme deficiency
   b. ABO/ Rh incompatability
   c. Breast feeding
   d. Hemoglobinopathies

Explanation :

Hemolytic disease of the newborn (HDN) is a blood problem in newborns. It occurs when your baby's red blood cells break down at a fast rate. It's also called erythroblastosis fetalis.



www.medicalcoupons.in

273. A 4 yr old child presents to ER with inspiratory stridor. X- ray reveals clear lung fields and respiratory tract obstruction. Which is the most common site of obstruction that cause inspiratory stridor is?

a. Nasopharynx
b. larynx
c. Trachea
d. Bronchus

Explanation :


Stridor is a symptom, not a diagnosis or a disease, and the underlying cause must be determined. It may be inspiratory (most common), expiratory, or biphasic, depending on its timing in the respiratory cycle, and the three forms each suggest different causes, as follows:

- Inspiratory stridor suggests a laryngeal obstruction
- Expiratory stridor implies tracheobronchial obstruction
- Biphasic stridor suggests a subglottic or glottic anomaly

274. Which of the following is not the adverse effects of morphine?

a. Midryasis
b. Orthostatic hypotension
c. Bradycardia
d. Depression

Explanation:



27[illegible]. Relaxation of mesangial cells of kidney is brought about by ?

a. PGF2
b. Endothelin
c. Vasopressin
d. CAMP

www.medicalcoupons.in

Explanation :

Agents causing contraction or relaxation of mesangial cells: -

Contraction	*Relaxation*
Endothelins	ANP
Angiotensin II	Dopamine
Vasopressin	PGE2
Nor-epinephrine	cAMP Q
Platelet -activation factor	
Platelet-derived growth factor	
Thromboxane A_2	
PGF_2	
LT-C_4 and D_4	
Histamine	

276. In breast feeding , hind milk portion is richer in ?
   a. Proteins
   b. Vitamins
   c. Fats
   d. Lactose

Explanation

Hind milk is rich in fat and provides more energy and satisfies the hunger of the baby. It comes towards the end of the feed. Fore milk is secreted at the start of the feed and is rich in protein, sugar, vitamins, mineral and water that satisfies the thirst of the baby. Colostrum is secreted in first three days after delivery and contains more antibodies, cells and high amounts of vitamins A, D, E and K. In transitional milk the immunoglobulin and protein content decreases while fat and sugar content increases. Preterm milk of mother who delivers prematurely contains more proteins, sodium iron, immunoglobulin and calories.

www.medicalcoupons.in

277. Kiesselbach's area has been damaged in a fist fight. The main artery to the area is?
   a. Anterior ethmoidal
   b. Posterior ethmoidal
   c. Spheno – palatine
   d. Greater – palatine

Explanation :

- Spheno-palatine artery is a branch of maxillary artery and is the chief artery to the nose.
- Antero-inferior part of the nasal septum has the Little's area (Kiesselbach's plexus), which is an important site of arterial anastomosis between the branches of: spheno-palatine, anterior ethmoidal and the labial branch of facial.
- This is a common site for nasal bleeding (epistaxis).

278. Increased suicidal tendency is associated with alteration in the brain levels of?
   a. Dopamine
   b. Serotonin
   c. GABA
   d. Noradrenaline

Explanation :

Decreased serotonin and its metabolite levels in brain is associated with depression whereas elevated levels of 5 – HT are associated with anxiety.

279. Vaccine given in adolescence is ?
   a. Hep - B
   b. Td/Tdap
   c. MMR
   d. Yellow fever

Explanation :

Indian Academy of Pediatrics (IAP) recommended vaccination for adolescents

Recommended Vaccination Schedule (2016)

Vaccine Tdap/Td	Schedule 10 years 9 to 14 years 15 years & olderor immunocompromiesd
• Adolescent & preadolescent girls of 9 -10 years of age only 2 doses needed • Either of the two vaccines can be given with minimum interval between doses to be 6 months • Females 15 years or older or immunocompromised adolescents, three doses at 0, 1 or 2 and 6 months to be given	

www.medicalcoupons.in

280. Commonest post operative complication of intubation is?
   a. Aspiration pneumonitis
   b. Sore throat
   c. Vocal cord granuloma
   d. Tracheal stenosis

Explanation :

Most common post – operative complication of intubation is sore throat which usually subsides in 2-3 days without any specific treatment.

281. Hormone responsible for diabetogenic state in pregnancy is?
   a. B- HCG
   b. AFP
   c. HPL
   d. Maternal progesterone

Explanation :

Human placental lactogen a/k as human chorionic somatotropin that is secreted by syncytiotrophoblast. It is mainly responsible for diabetogenic state in pregnancy, as it antagonises the action of insulin. It leads to maternal lipolysis and increased levels of circulating free fatty acids thus provides a source of energy for maternal metabolism and fetal nutrition. It is a potent angiogenic hormone therefore,may play important in fetal vasculature formation.

282. All of the following are the causes of metabolic encephalopathy except?
   a. Carbon monoxide poisoning
   b. Electrolytic imbalance
   c. Diabetic neuropathy
   d. Increased catecholamines levels

Explanation :


Metabolic encephalopathy (toxic metabolic encephalopathy) is a broad category that describes abnormalities of the water, electrolytes, vitamins, and other chemicals that adversely affect brain function. Other causes of metabolic encephalopathy include carbon monoxide or cyanide poisoning, which prevents hemoglobin from carrying oxygen in the bloodstream and results in tissue anoxia. Toxic metabolic encephalopathy can also include medication side effects or drug ingestions affecting the chemical transmitters in the brain. Called neurotransmitters, dopamine, GABA, serotonin, acetylcholine, and glutamine help nerve endings pass electrical signals between brain cells.

www.medicalcoupons.in

283. Adult Polycystic disease of kidney may have cysts in all of the following organs except?

a. Liver
b. Lungs
c. Pancreas
d. Spleen

Explanation :



284. Most common inherited cause of mental retardation is ?

a. Down syndrome
b. Hypothyroidism
c. Fragile x- syndrome
d. Neurofibromatosis

Explanation :

Fragile x- syndrome – due to defects in FMR 1 gene that is present in long arm of X-chromosome and is associated with CGG repeats.

On karyotyping , the x – chromosomes appears broken. It is the important inherited cause of mental retardation. Patient presents with – long face, large mandible, large ears, large testes.

285. Which of the following drug is used in treatment of cachexia?

a. Cyprohepatidine
b. Megestrol acetate
c. 5- flurouracil
d. Cetrorelix

www.medicalcoupons.in

Explanation :

Cachexia is a complex metabolic syndrome associated with many chronic or end-stage diseases, especially cancer, and is characterized by loss of muscle with or without loss of fat mass. Among the treatments proposed in the literature for cancer-related cachexia, some proved to be ineffective, namely, cyproheptadine, hydrazine, metoclopramide, and pentoxifylline. Among effective treatments, progestagens are currently considered the best available treatment option for cancer-related cachexia.

286. Cerebellar degeneration is seen in ?
   a. Renal cell carcinoma
   b. Hodgkin lymphoma
   c. Thymoma
   d. Ca.stomach

Explanation :

Paraneoplastic syndromes :

Syndrome	**Associated tumours**
Retinal degeneration	Small - cell carcinoma of lung
Opsoclonus myoclonus	Ovarian, lung Neuroblastoma (in children)
Sensory neuropathy	Small-cell carcinoma of lung Hodgkin's disease
Limbic encephalitis	Small-cell carcinoma of lung, Hodgkin's disease
Myelitis	Small-cell carcinoma of lung
Cerebellar degeneration	Small – cell carcinoma of lung, Ovarian, Breast Hodgkin's disease
Subacute motor neuronopathy	Hodgkin's disease, Small – cell carcinoma of lung
Sensorimotor peripheral neuropathy	Small – cell carcinoma of lung, Breast Other carcinoma

www.medicalcoupons.in

www.medicalcoupons.in

Lambert – Eaton myasthenic syndrome	Small-cell carcinoma of lung
Dermatomyositis / Polymyositis	Lung, breast, ovary
Guillain – Barre	Hodgkin's disease

287. Which of the following is a bony lesion in vitamin - A intoxication?
   a. Bone pain
   b. Bone demineralization
   c. Hyperostosis
   d. All of the above

Explanation :

Vitamin A. intoxication produces following bony manifestation -

Bone demineralization

Bone pain ,


Hypercalcemia .

Hyperostosis

Vitamin D intoxication (delayed effect) causes persistent hypercalcemia and hyper phosphatemia which may produce:

1). Urinary Lithiasis.
2). Metastatic calcification which affects kidneys, bronchi, pulmonary alveoli, muscles, arteries and gastric mucosa & Renal failure leading to death.

288. Identify the given image



   a. Candida tropicalis
   b. Pityriasis rosea
   c. Tuberous sclerosis
   d. Roseola infantum

www.medicalcoupons.in

Explanation :

Acute symmetrical self limiting papulosquamatous disorder. Seen specifically on the trunk in children and young adults common during spring and fall.1st manifestation is 1 2-6 cm annular lesion – herald patch with fine scaling, fine cigarette paper like collarette of scales usually over the chest. Typical truncal presentation is chirstmas tree pattern /fir tree pattern. Hnging curtain sign is seen. Associated with HHV 7. As mentioned previously, it is self limiting and hence no treatment is required. If pruritic , antihistamines and tropical steroids and UV – B phototherapy.

289. Chorionic component in early stages of pregnancy is composed of?
   a. **Cytotrophoblast +syncytiotrophoblast + somatic mesoderm + fetal blood vessels**
   b. Cytotrophoblast + syncytiotrophoblast
   c. Syncytiotrophoblast + somatic mesoderm + fetal blood vessels
   d. None of the above

Explanation :

Placenta

The placenta is a fetomaternal organ. The fetal portion of the placenta is known as the villous chorion. The maternal portion is known as the decidua basalis. The two portions are held together by anchoring villi that are anchored to the decidua basalis by the cytotrophoblastic shell.

Function

The placental membrane separates maternal blood from fetal blood. The fetal part of the placenta is known as the chorion. The maternal component of the placenta is known as the decidua basalis.

Although the placental membrane is often referred to as the placental barrier, many substances, both helpful and harmful, can cross it to affect the developing embryo.

Structure

· **Primary chorionic villi** are solid outgrowths of cytotrophoblast that protrude into the syncytiotrophoblast.

· **Secondary chorionic villi** have a core of loose connective tissue, which grows into the primary villi about the third week of development.

· **Tertiary chorionic villi** contain embryonic blood vessels that develop from mesenchymal cells in the loose connective tissue core. These blood vessels connect up with vessels that develop in the chorion and connecting stalk and begin to circulate embryonic blood about the third week of development.

www.medicalcoupons.in

290. Earliest markers for down syndrome is?
   a. Unconjugated estriol
   b. Ultra sound markers (nuchal translucency)
   c. AFP
   d. Amniocentesis

Explanation :

ultrasound (also called sonography) is to confirm the gestational age of the fetus. It can also pick up problems of a serious medical nature, such as blockage of the small intestine or heart defects. Knowing these defects exist as early as possible will benefit the treatment of the child after birth.

Studies showed that there was a strong association between the size of a collection of fluid at the nack of the fetal neck, called nuchal translucency, and the risk of Down syndrome. Early attempts to use a measurement of the nuchal area were limited by a wide variety in measurement techniques. Recently, standardized guidelines on measuring nuchal translucency along with specific training and certifications have been instituted, making this ultrasound measurement useful as part of the first trimester screen.

291. Pulmonary eosinophilia is caused by ?
   a. Wuchereria bancofti (microfilaria)
   b. Eccinococcus granulosus
   c. Ascaris lumbricoides
   d. Taenia solium

Explanation :

Simple pulmonary eosinophilia (SPE), also known as Loeffler syndrome, is a rare, temporary (transient) respiratory disorder characterized by the accumulation of eosinophils in the lungs. They are usually produced in response to allergens, inflammation or infection and are particularly active in the respiratory tract. Most cases of SPE are believed to be due to an allergic reaction to drugs and infection caused by Wuchereria bancrofti, a filarial infection.

292. which of the following is the most common complication of extracapsular cataract surgery ?
   a. Vitreous hemorrage
   b. Opacification of posterior capsule
   c. Retinal detachment
   d. None

Explanation :

Patient presents with painless loss of vison. Post operatively, opacification of posterior capsule occurs in 10- 50% of cases after 3-5 yrs.

www.medicalcoupons.in

293. Specific congenital abnormality associated with DM is?

a. ASD
b. VSD
c. Caudal regression syndrome
d. TOF

Explanation :

Sacral agenesis may be expected in 1% of offspring of diabetic mothers. In 16% of cases of caudal regression syndrome, the mother is diabetic; rarely, the father is diabetic instead. The syndrome appears sporadically but is 100 times more common in one sibling of twin births.

294. Which of the following joint disease/radiograph finding relationships is correct?

a. Osteoarthritis – erosions with overhanging margins
b. Rheumatoid arthritis – ankylosis
c. Gout – osteophytes
d. Psoriatic arthiritis – "bamboo spine"

Explanation :

Rheumatoid arthritis is associated with ankylosis of the joint space caused by the development of pannus, which overgrows and destroys the articular cartilage

295. Sudden and irresistible force compelling a person to commit an offence consciously is known as ?

a. Illusion
b. Obsession
c. Twilight state
d. Impulsiveness

Explanation :

Impulse: This is a sudden and irresistible force compelling a person to the conscious performance of some action without motive or fore-thought. A sane person is capable of controlling an impulse, but an insane person may do things on impulse.

Types:

a. **Kleptomania**Q- An irresistible desire to steal articles of little value.

b. **Pyromania**- an irresistible desire to set things to fire.

c. **Mutilomania**- An irresistible desire to mutilate animals.

d. **Dipsomania**- an irresistible desire for alcoholic drinks at periodic intervals.

e. **Sexual impulses**- including sexual perversions.

f. **Suicidal and homicidal impulses.**

www.medicalcoupons.in

g. **Twilight state** is seen in visual hallucinations, Epilepsy, Head injury, Hysteria, Punch-drunkenness.

296. Sphingomyelin and phosphatydil choline are precursors of?
   a. Surfactant in lung
   b. Mucopolysaccharides
   c. Cholesterol
   d. Ion transport proteins

Explanation :

Surfactant is dipalmitoyl phosphatidylcholine. Surfactant is secreted by type II pneumocytes by exocytosis.

297. All of the following are produced by hepatocytes except?
   a. Von Wille brand factor
   b. Prothrombin
   c. Factor 7
   d. Fibrinogen

Explanation :

VWF →synthesized in endothelial cells

Protein synthesized by liver

Antithrombin III, fibrinogen, Factor II, VII, IX and X (vit.K. Dependent) protein-C (vit. K.
dependent)

298. Most common cause of shock in surgical patient is ?
   a. Anaphylactic shock
   b. Compressed cardiogenic shock
   c. Neurogenic shock
   d. Hypovolumic shock

Explanation :

The most common cause of shock in the surgical or trauma patient is loss of circulating volume from hemorrhage. The clinical and physiologic response to hemorrhage has been classified according to the magnitude of volume loss.

www.medicalcoupons.in

a. Loss of up to 15% of the circulating volume (700 to 750 mL for a 70-kg patient may produce little in terms of obvious symptoms, while

b. Loss of up to 30% of the circulating volume (1.5 L) may result in mild tachycardia, tachypnea, and anxiety.

c. Hypotension, marked tachycardia [i.e., pulse >110 to 120 beats per minute (bpm)], and confusion may not be evident until more than 30% of the blood volume has been lost;

d. Loss of 40% of circulating volume (2 L) is immediately life threatening, and generally requires operative control of bleeding.

299. Chemical peeling is done by?

a. Trichloroacetic acid
b. Phosphoric acid
c. Carboxylic acid
d. Kojic acid

Explanation :

CHEMICAL PEEL

1). Chemical peel produces a controlled partial thickness injury to the skin.

2). Following the insult to the skin, a wound healing process ensues that can regenerate epidermis from surrounding epithelium and adnexal structures, decrease solar elastosis, and replace and reorient the new dermal connective tissue.

3). The result is an improved clinical appearance of the skin, with fewer rhytids and decreased pigmentary dyschromia.

Peeling Agents

1). Glycolic acid.
2). Jessner's solution.
3). Pyruvic acid.
4). Resorcinol.
5). Salicylic acid.
6). Trichloracetic acid.
7). Deep chemical peels.
8). Combinations peels: Salicylic acid +Trichloracetic acid.

www.medicalcoupons.in

300. WHO defines adolescent age between ?

a. 10 – 19 yrs
b. 10 – 14 yrs
c. 10 – 25 yrs
d. 9 – 14 yrs

Explanation :

Adolescence

Adolescence is defined as the period from the onset of puberty to the termination of physical growth and attainment of final adult hight, i.e. adolescence is considered as a period of transition from childhood to adulthood. Adolescence is usually the period 10 to 20 yr.

Puberty

Puberty is the biological process in which a child become adult, i.e. biological process which occurs during transition from childhood to adulthood. Changes include appearance of secondary sex characteristics and development of reproductive capacity.

WHO/UN definitions

Adolescent 10-19 years

Early adolescent 10-13 years

Middle adolescent 14-15 years

Late adolescent 16-19 years

Youth 15-24 years

www.medicalcoupons.in

FMGE JUNE 2018

ANATOMY JUNE 2018

1. Coracoid process of the scapula is which type of epiphysis ?

a) Pressure
b) Traction
c) Aberrant
d) Atavistic

Atavistic epiphysis: A bone that is independent phylogenetically but is now fused with another bone. These types of fused bones are called atavistic, e.g., the coracoid process of the scapula, which has been fused in humans.

2. Pointing index finger is seen in which nerve injury

a) Axillary
b) Radial
c) Median
d) Ulnar

"Pointing finger" deformity is caused due to injury to **the median nerve** in the mid-forearm by paralysis of flexor digitorum superficialis.

3. NOT a content of carpal tunnel

a) Flexor pollicis longus
b) Flexor digitorum profundus
c) Ulnar nerve
d) Median nerve

The carpal tunnel contains **the median nerve**, **the four tendons of the flexor digitorum profundus, the four tendons of the flexor digitorum superficialis, and the flexor pollicis longus (FPL) tendon**. Carpal tunnel syndrome (CTS) is the result of median nerve compression within the carpal tunnel.

4. Which of the following structure passes through Guyon's canal

a) Flexor carpi radialis
b) Radial nerve
c) Ulnar nerve
d) Median nerve

www.medicalcoupons.in

The ulnar nerve and ulnar artery pass through the Guyon canal as they pass from distal forearm to the hand.

Roof - Formed by palmar carpal ligament.

Floor - Formed by flexor retinaculum.

5. NOT a content of superior mediastinum

a) Pulmonary trunk
b) Thymus
c) Oesophagus
d) Arch of aorta

Contents of Superior Mediastinum:

→ **Organs: thymus, trachea, esophagus**
→ **Arteries: aortic arch**, brachiocephalic trunk, left common carotid artery, left subclavian artery.
→ **Veins and lymphatics:** superior vena cava, brachiocephalic veins, the arch of the azygos, thoracic duct.

6. Abdominal part of esophagus is supplied by


a) Bronchial artery
b) Abdominal aorta
c) Right gastric artery and inferior phrenic artery
d) Left gastric artery and inferior phrenic artery

The abdominal esophagus is supplied by the **left gastric artery (a branch of the coeliac trunk)** and left inferior phrenic artery.

7. Which of the following artery is a direct branch of abdominal aorta?

a) Superior suprarenal artery
b) External iliac artery
c) Internal iliac artery
d) Testicular artery

www.medicalcoupons.in



8. **Brunner's gland is a histological finding of which structure in the. gastro-intestinal tract ?**

a) Esophagus
b) Stomach
c) Duodenum
d) Liver

Brunner Gland: - Present in submucosa of proximal duodenum, secretes alkaline secretions to neutralize acid

9. Secondary ossification centre appears before birth at

a) Upper end of femur
b) Lower end of femur
c) Lower end of tibia
d) Lower end of fibula

Secondary ossification centers **develop at later times**. Ossification of the **distal end of the femur** to form the condyles and epicondyles, begins shortly before birth.

www.medicalcoupons.in

10. A 44 year old male came in to your clinic post hernia repair and wants you to evaluate his condition. While eliciting the cremasteric reflex, you notice that the scrotum did not ascend on the side of the surgery, suggesting an iatrogenic injury to the:-

a) Sacral plexus
b) Femoral branch of genitofemoral nerve
c) Genital branch of genitofemoral nerve
d) Femoral nerve

11. In the cremasteric reflex, it is noticed that the scrotum did not ascend on the side of the surgery, suggesting an iatrogenic injury to the Genital branch of genitofemoral nerve.

The cremasteric reflex is a **superficial reflex found in human males** that is elicited when the inner part of the thigh is stroked. Stroking of the skin causes the cremaster muscle to contract and pull up the ipsilateral testicle toward the inguinal canal.

- Cremaster muscle Lies in scrotum
- Sensory supply to cremaster:
  → Laterally -Femoral Branch of genitofemoral nerve
  → Medially - Ilioinguinal nerve
- Motor supply to cremaster.
  → **Genital branch of genitofemoral nerve.**

12. Dorsum of first webspace is supplied by which nerve ?

a) Superficial peroneal
b) Deep peroneal
c) Sural
d) Posterior tibial

The superficial peroneal nerve innervates the peroneus longus and peroneus brevis muscles and the skin over the antero-lateral aspect of the leg along with the greater part of the dorsum of the foot **(with the exception of the first web space, which is innervated by the deep peroneal nerve).**


13. Nerve damaged in fracture neck fibula :

a) Superficial peroneal
b) Deep peroneal
c) Common peroneal
d) Tibial


www.medicalcoupons.in

The common fibular nerve is most commonly damaged by a fracture of the fibula, or the use of a tight plaster cast. The anatomical course of the common fibular nerve causes it to wrap round the neck of the fibular, and so any fractures of the fibular neck can cause nerve palsy.

14. Identify the marked area in the image below



a) Nasion
b) Glabella
c) Pterion
d) Lambda



www.medicalcoupons.in

15. A 43-year-old male came in with complaints of headache, vomiting, blurring of vision, and drowsiness. His son claimed that he had a fall a few hours ago. If you suspect an increased intracranial pressure as the root cause of this, which cranial nerve do you first check since it is commonly injured in cases of raised ICP?

a) 3rd
b) 4th
c) 5th
d) 6th

Mc nerve involved in increased ICT Abducens (VI).

Reasons :

→ Longest intradural course
→ Taking a sharp bend over apex of the petrous part of temporal bone.
→ Passing through the cavernous sinus, not supported laterally by any dura mater.

16. Sensory innervation of cornea is by nerve

a)3rd
b)4th
c)5th
d) 6th

The cornea is one of the most sensitive tissues of the body, as it is densely innervated with sensory nerve fibres via **the ophthalmic division of the trigeminal nerve** by way of 70–80 long ciliary nerves.

17. Masseter is supplied by nerve

a) Mandibular
b) Facial
c) Glossopharyngeal
d) Hypoglossal

Along with the other three muscles of mastication (temporalis, medial pterygoid, and lateral pterygoid), the masseter is innervated by **the anterior division of the mandibular division (V3) of the trigeminal nerve**.

www.medicalcoupons.in

18. Deepest nucleus of cerebellum:-

a) Dentate
b) Emboliform
c) Fastigial
d) Globose

The four deep cerebellar nuclei are in the center of the cerebellum, embedded in the white matter. Dentate, Emboliform, Globose, Fastigial from lateral to medial location in the cerebellar white matter**.**

An easy mnemonic device to remember their names and positions relative to the midline is the phrase **"Don't Eat Greasy Food",** where each letter indicates the lateral to medial location in the cerebellar white matter**.**

PSM JUNE 2018

1. A 25 year old young male was travelling from abroad to India. At the airport he was diagnosed with some disease condition for which India has quarantine. Disease usually not seen in a country but brought from abroad is called as?

a) Endemic
b) Epidemic
c) Zoonotic
d) Exotic

→ **Exotic:** Disease usually not seen in a country but brought from abroad.
→ **Zoonotic:** A disease which is transmitted to man from animals e.g. Rabies, plague.
→ **Epidemic:** When disease has exceeded its normal expectancy.
→ **Endemic:** Constant or continuous presence of a disease in a population.

2. Incidence is best measure by.

a) Cross sectional study
b) Cohort study
c) Case control study
d) Double blind study

We do fellow up, useful parameters obtained e.g. Incidence, relative risk, Attributable risk, population attributable risk.

www.medicalcoupons.in

Case control - odds ratio

Cross sectional - prevalence

Ecological study Group characteristic

3. Primary prevention aims to prevent disease or injury before it ever occurs. Primary prevention in myocardial infarction are all of the following EXCEPT?

a) Maintenance of normal body weight
b) Change in life style
c) Change in nutritional habits
d) Screening for hypertension

All screening & diagnostic tests comes under secondary level of prevention.

Hence, Screening for hypertension is a secondary level of prevention. It aims to reduce the prevalence of a disease by shortening its duration

All other options given protects against development of MI. Hence, primary level of prevention.

4. Installation and usage of sanitary latrines by general public constitutes which level of prevention:

a) Health promotion
b) Specific protection
c) Early diagnosis and treatment
d) Disability limitation and rehabilitation

- Include: safe water
- Installation of sanitary latrines
- Environmental: Improving in housing condition
- Modification: Insect / Rodent control

Level of prevention	Modes of intervention
Primary level	Health promotion & specific protection
Secondary level	Early diagnosis & treatment
Tertiary level	Disability limitation & rehabilitation

www.medicalcoupons.in

5. A microbiologist is preparing BCG vaccine in the laboratory. Which of the following is diluent used with BCG vaccine?

a) Normal saline
b) Distilled water
c) Dextrose
d) Colloids

Lyophilized vaccines & their diluents:

BCG vaccine → Normal saline

Yellow fever vaccine → cold saline

Japanese Encephalitis → phosphate buffer saline

HiB → DPT / saline

Measles, MMR → Distilled /sterile water

6. A 'Case' in TB is defined as:

a) X-ray positive
b) Sputum positive
c) Mantoux positive
d) History of cough

Case of tuberculosis

A definite case of TB (defined below) or one in which a health worker (clinician or other medical practitioner) **has diagnosed TB** and has decided to treat the patient with a full course of TB treatment. Note. Any person given treatment for TB should be recorded as a case.

TB is diagnosed by sputum smear

CBNAAT

ALCPA, LED - FM

CXR, LST

All test used in revised national tuberculosis control programme.

www.medicalcoupons.in

7. Cyclopropagative life cycle is defined as the disease agent which undergoes cyclical change, and multiplies in the body of the arthropod. This type of life cycle is seen in which of the following disease?

a) Malaria
b) Plague
c) Filarial
d) Yellow fever

Biological transmission of arthropod Borne disease:

Transmission Definition	**Propagative (Only multiplication)**	**Cyclo propagative (Multiplication + development)**	**Cyclo development (Only development)**
Examples	Plague Yellow fever	Malaria	Filaria Guinea worm

8. Causative agent for kala-azar/ visceral leishmaniasis:

a) Leishmania donovani
b) Leishmania tropica
c) Leishmania braziliensis
d) None of the above

- Disease transmitted vector is sandfly.
- Transaminases of **Leishmania donovani**, the causative organism of kala-azar.
- Leishmania tropica - caused cutaneous leishmaniasis commonly known As oriental sore or delhi boil or Baghdad boil.
- Leishmania braziliensis muco cutaneous leishmaniasis all three transmitted By sand fly.

9. Which of the following is true for DOTS?

a) Treatment under supervision
b) Every patient given the same treatment
c) Streptomycin is given to all the patients
d) Daily regimen

Treatment under supervision→ YES it is a directly observed treatment, Short - Course (DOTS). Pts has to consume medication in front of DOTS provider.

www.medicalcoupons.in

10. Sensitive parameter of combined obstetric and pediatric care in the countries:

a) Post neonatal mortality rate
b) Infant mortality rate
c) Neonatal mortality rate
d) Perinatal mortality rate

Perinatal mortality rate: its death occur during 28 weeks to till day post delivery

11. Number of live births per 1000 women in the reproductive age reproductive age group in a year refers

a) Gross reproduction rate
b) Total fertility rate
c) Net reproduction rate
d) General fertility rate

General fertility rate is defined as **number of live births per thousand women in the age group (15-49 years) in a given year**.

Total fertility rate → Total children born to a women or couple during there life span

Gross reproduction rate → Total number of girl children born to a women in entire reproduction

Net reproduction → Total girl children produced during entire reproduction life span taking into a amount mortality

12. Maximum biological value is of:

a) Pulses
b) Soyabean
c) Milk
d) Egg

Biological value tells N2 absorption V5N2 Excretion

- Egg – 94
- Milk – 84
- Soya bean – 73

www.medicalcoupons.in

13. Mode of disposal of sewage ground water:

a) Activated sludge process
b) Soakage pit
c) Oxidation pond
d) All of the above

The activated sludge process is still the most widely used biological treatment process for reducing the concentration of organic pollutants in sewage. It's aerobic oxidation process used as a secondary treatment. Heart of this method its Aeration tank. Well-established design standards based on empirical data have evolved over the years.

14. Value of free residual chlorine in Drinking water should be:

a) 0.2 mg/L
b) 0.3 mg/L
c) 0.4 mg/L
d) 0.5 mg/L

It's should be minimum 0.5 mg/l (ppm) in drink water

- ≥ 1.0 mg /l swimming pool
- ≥ 2.0 mg/l kill cyclops
- Measured by chloroscope

15. **Mark the correct statement regarding Ayushman Bharat**.

a) Umbrella scheme includes Pradhan Mantri Jan Aarogya Yojna and subcenters.
b) ↑Benefit to 40 % of poor and vulnerable population
c) Covers only tertiary health care centres
d) Cashless paperless service by only government hospitals.

Ayushman Bharat

- Under Ministry of Health and Family Welfare
- Umbrella scheme
  → Health and wellness centre (HWC)
  → Earlier subcentres were there
  → Comprehensive PHCs
  → Mid level Heath provider -Nurse Ayurvedic practitioner

www.medicalcoupons.in

- **Pradhan Mantri Jan Arogya Yojana (PMJAY)**
  → 23 September 2018
  → ↓ Out of pocket expense
  → ↑Benefit to 40 % of poor and vulnerable population
  → Cover secondary and tertiary health
  → Coverage of 5 lakh/family
  → No restriction of family size
  → Cashless ,paperless service by government and private hospitals

16. The frequently occurring value in a data is:

a)Median
b) Mode
c)Standard deviation
d)Mean

→ The **mode** is the value that appears most frequently in a data set.
→ **Median** is middle value
→ The ***mean*** is the ***average*** or the most common value in a collection of numbers.
→ **Standard deviation** measure of dispersion.

17. Problem village is all EXCEPT.

a)Where no water source in a distance of 1.6 km from community
b)Water is more than depth of 15
c)There is excess of Na, K, salts
d) Risk of guinea worm infection

Guinea worm Eliminated from India Feb 2000.

18. A 25 year old female is using vaginal sponge as a barrier contraceptive during sexual intercourse. Which of the following Spermicide is used in contraceptive TODAY?

a)NET-EN
b)DMPA
c)Nonoxynol - 9
d)Nor-ethinosterol

- "TODAY" is vaginal sponge is barrier type of contraceptive.
  NON OYYNOL - 9 → kill the sperm by rupture of plasma membrane of Acrosomal cap of sperm.
- Failure rate of today → 9-20 /Healthy Women Years.

www.medicalcoupons.in

- TODAY also associated with toxic shock syndrome .

19. Bagassosis occurs with:

a)Silica fibers.
b)Carbon particles
c)Sugarcane fibers
d)Cotton

One of these lesser-known illnesses is Bagassosis. Defined as the **inflammation of the lung caused by the inhalation of dust from bagasse (sugar cane dust)**, the disease was initially discovered more than 50 years ago in India.

20. Cytotoxic drug is disposed in:

a) Yellow bag
b) Red bag
c) Blue bag
d) Black bag

- **Yellow bag**
  → Human anatomical waste
  → Animal waste
  → Soiled waste (cotton / cloths)
  → Chemical waste
  → Discarded medicine (expired medicine, cytotoxic drug)
  → Liquid chemical waste
  → Micro biological, lab waste
- **Red category** → solid contaminated (plastic / rubber)
- **White category** → wasted sharp (needle, Sx blades, and scalpels)
- **Blue category** → glasses, metallic body implants

21. A 24 year old reproductive age female visited to the rural PHC for IUCD insertion to prevent pregnancy. Which of the following is the most common side effect of IUCD insertion?

a) Ectopic pregnancy
b) Pelvic infection
c) Pain
d) Bleeding

- **Mc side effect bleeding** (give to female iron tablet for 2 months).
- Pain is 2nd mc. but its mc cause of removal of IUCD.

22. The Factories Act, 1948 is a social legislation which has been enacted fo occupational safety, health and welfare of workers at work places. Wh is th maximum number of work hours/ week, prescribed under the Factor es a t?

a) 42 hours/week
b) 48 hours/week
c) 54 hours/week
d) 60 hours/week

- The Factories Act. 1948 is a social legislation which has een nacted for occupational safety, health and welfare of workers at work places.
- Its 9 hours a day & maximum 48 hours. week.
- 60 hours , week then paid extra.

23. A 36 year old male presented to the OPD wi h di ficulty in breathing while walking, back pain, unable perform physical activit / and excessive laziness. On examination his body weight is 106 k s. W hich of the following is height independent obesity index?

a) Corpulence index
b) Quetlet index
c) Brocca index
d) Ponderal index

Corpulence Index = Actual w eigh / Desirable weight

So, **Corpulence Index is " eigh independent index" of obesity**; cutoff ≤ 1.2.

24. Cafeteria approa h is elated with:

a) Diet progra
b) Child and mat nal health
c) National v ctor borne disease control programme
d) Contracep n

Fam planning & contraception - "Cafeteria approach" has new name "GATHER"

G: Gra de the couples

A: Ask them their contraception requirement

T: ell them about available methods

H: Help them choose best method for them

E: Explain

www.medicalcoupons.in

R: Returns

25. A 42 year old obese female presented to the rural PHC. Which of the following is the appropriate statistical test to find out obesity as a significant risk factor for breast cancer?

a) Student's paired 't' test
b) Student's unpaired 't' test
c) Chi-square test
d) Wilcoxon's signed rank test.

- Obesity is qualitative variable : breast cancer also qualitative.
- So, appropriate statistical test to find out obesity as significant risk factor for breast cancer is Chi-square test.

26. An important measure of communicability of a disease is:

a) Incidence rate
b) Case fatality rate
c) Prevalence rate
d) Secondary attack rate

SAR =(Number of secondary cases in 1 incubation period / Total Susceptible) x 100

Measles SAR – 80 %

Chicken pox – 90%

Petrussis – 90 %

Mumps – 86%

27. Which one of the following is FALSE regarding confounding factor in epidemiological studies?

a) Associated both with exposure and disease
b) Distributed equally between study and control groups
c) Independent risk factor for disease in question
d) Source of bias is interpretation

Confounding factor Any factor associated with both exposure, outcome and has an independent effect in causation of outcome is confounder.

BEST METHOD TO REMOVE CONFOUNDING:-

→ **MATCHING** [MC used/ simplest method]- Removes confounding bias by distributing known confounding factors equally in two groups
→ **RANDOMISATION** (2ND BEST)
→ **RESTRICTION**
→ **STRATIFICATION**
→ **STATISTICAL MODELING** / MULTIVARIATE ANALYSIS
→ **STRATIFIED RANDOMIZATION** (BEST METHOD).

28. Denominator in calculation of case fatality rate is:

a)Total number of deaths due to all causes
b)Total number of hospital admissions
c)Total number of cases due to the disease concerned
d)Total number of deaths due to the disease concerned

Case Fatality :

- Refers to proportion of fatal cases among those who have the disease.
- Provides an index of the deadliness of a particular disease within a specific population.

CFR = (Number of deaths due to disease / Number of cases of disease) x 100

29. An epidemic is the rapid spread of disease to a large number of people in a given population within a short period of time. A well of contaminated water resulting in an epidemic of acute watery diarrhoea is a typical example for which of the following?

a) Common source, single exposure epidemic
b) Common source, continuous exposure epidemic
c) Slow epidemic
d) Propagated epidemic

- A well of contaminated water resulting in an epidemic of acute watery diarrhea is an example of **Common source, continuous exposure epidemic** as there is continuous and repeated exposure.
- Multiple peaks known as **SECONDARY WAVES** are seen.

www.medicalcoupons.in

ORTHO JUNE 2018

1. **A 65 years old male brought to casualty with history of fall. The patient is in severe pain, there is tenderness on the greater trochanter and also the patient is not able to walk. The lateral border of the foot touches the bed. What is the most common complication in the given clinical scenario?**

a)Non union
b)Ischemic necrosis
c)Malunion
d)Pulmonary complications

The given clinical scenario depicts the Fracture neck of femur (intertrochanteric)

Fractured neck of femur presents with the following clinical features:

- Painful movements of the hip joint
- Shortening and external rotation of the affected limb
- **Extra-capsular (inter-trochanteric)** fracture is more common than intra-capsular.
- Most common complication of **Extra-capsular fracture of neck of femur of neck is mal-union.**
- In **inter-trochanteric fractures**, there is usually gross deformity and pronounced lateral rotation.
- In Intra-capsular (neck of femur fracture)
- Most common complication is avascular necrosis > > non-union.

2. **8 years old boy sustained fracture to the hand while playing with his sister. In which of the following condition, the Triangular relation of the elbow is maintained?**

a) Fracture ulna
b) Anterior dislocation of elbow
c) Posterior dislocation of elbow
d) Supracondylar fracture of humerus

- Triangular relation of elbow includes medial epicondyle, lateral epicondyle and tip of olecranon.
- The relationship is destroyed by dislocations of elbow
- **It is maintained in supracondylar fracture of Humerus (occurs above elbow joint)**

www.medicalcoupons.in

3. Which of the following statement about tubercular osteomyelitis is not true?

a)It is a type of secondary osteomyelitis
b)Sequestrum is uncommon
c)Periosteal reaction is characteristic
d)Inflammation is minimal

- Tuberculous osteomyelitis is a type of secondary osteomyelitis.
- Sequestrum is uncommon in tubercular osteomyelitis.
- **Periosteal reaction is common in Osteomyelitis, but there is little or no periosteal reaction in tubercular osteomyelitis.**
- In tubercular osteomyelitis, Inflammation is rare.

4. Most common site of osteomyelitis in children

a) Diaphysis
b) Metaphysis
c) Epiphysis
d) Growth plate

- Overall Most common location of osteomyelitis is metaphysis. It is due to high vascularity, hair-pin loop system of blood supply. more prone to trauma and monocyte macrophage system is weaker.
- Most common part of osteomyelitis is lower end of Femur.
- Most common organism of osteomyelitis is staphylococcus aureus.

5. Primary osteoarthritis affect all except:

a) Hip Joint
b) Knee joint
c) Distal interphalangeal joint
d) Metacarpophalangeal joint

- **Osteoarthritis involves DIP and PIP with sparing of MCP.**
- Rheumatoid arthritis involves PIP and MCP with sparing of DIP.
- **Osteoarthritis commonly involves knee joint in India.**
- Hip joint is rarely involved in osteoarthritis.

www.medicalcoupons.in

MICRO JUNE 2018

1. A liquid media used to suppress commensal bacteria while allowing the pathogen to remain viable and grow. Which of the following is an enrichment media?

a) Alkaline peptone water
b) Monsour's taurocholate Tellurite peptone water
c) Selenite F broth
d) All of the above

Enrichment media -

- LIQUID culture media
- These media are used to suppress commensal bacteria while allowing the pathogen to remain viable and grow.
- Example –
  - → Alkaline peptone water - V. cholerae
  - → Selenite F-broth - Salmonella, shigella
  - → Monsour's taurocholate tellurite peptone water - V. cholerae.

2. Biological indicator for determining efficacy of autoclaving is

a) Pseudomonas aeruginosa
b) Clostridium perfringenes
c) Bacillus stearothermophilus
d) Salmonella typhi

There are three indicators that may be used to detect the efficacy of the autoclave process:
(1) Physical: pressure and temperature recording devices,
(2) Chemical: indicators that change color after being exposed to specific temperatures, such as temperature sensitive tape. The color change upon exposure to the given temperature, and
(3) Biological: Bacillus stearothermophilus spores are used, due to its resistance to heat, for the testing that measures the biological performance of the autoclave process.

3. Yaws is caused by:

a) **T. Pertunae**
b) T. Pallidum
c) T. Caraetum
d) Hemophilus

www.medicalcoupons.in

Yaws is an infectious disease caused by a spiral-shaped bacterium (spirochete) known as **Treponema pertenue**. Yaws is usually transmitted by direct contact with the infected skin sores of affected individuals.

4. Babesiosis is most commonly transmitted by:

a) Pigs
b) Rats
c) Sand fly
d) Ticks

The deer tick (Ixodes dammini or scapularis) is the most common vector that transmits babesiosis. In extremely rare cases, babesiosis may be transmitted following a blood transfusion with blood that is contaminated with the microorganism.

5. Which bacteria shows Quellung reaction?

a) Pneumococcus
b) Gonococcus
c) Staphylococcus
d) All of the above

The quellung reaction, also called the Neufeld reaction, is a biochemical reaction in which antibodies bind to the bacterial capsule of **Streptococcus pneumoniae, Klebsiella pneumoniae, Neisseria meningitidis, Bacillus anthracis, Haemophilus influenzae, Escherichia coli, and Salmonella**.

6. Molluscum contagiosum is caused by

a) Bacteria
b) Fungus
c) Protozoa
d) Virus

Molluscum contagiosum is an infection caused by **a poxvirus (molluscum contagiosum virus)**. The result of the infection is usually a benign, mild skin disease characterized by lesions (growths) that may appear anywhere on the body.

www.medicalcoupons.in

7. SSPE is a complication of

a) Measles
b) Mumps
c) Rubella
d) Rabies

Subacute sclerosing pan encephalitis (SSPE) is a very rare, but fatal disease of the central nervous system that results from **a measles virus infection** acquired earlier in life. SSPE generally develops 7 to 10 years after a person has measles, even though the person seems to have fully recovered from the illness.

8. Which subtype of human papilloma virus has maximum chances of causing carcinoma cervix:

a) HPV 16&18
b) HPV 6&11
c) HPV 31&32
d) HPV 1&2

→ **Maximum chance of developing the carcinoma cervix (16 & 18) HPV**
→ HPV 6 & 11 cause genital warts (Condyloma acuminata)
→ HPV 1 & 2 cause common wa
→ Rx - Podophyllin resin Cry surgery

9. Stain used in staining fungal elements

a) Acid fast stain
b) Gram stain
c) Methenamine silver
d) All of the above

Fungal stains

- 10% KOH
- LPCB (lactophenol cotton blue)
- GMS (Gomori Methenamine Silver)
- Calcofluor white (Fluorescent stain)
- Mucicarmine stain (used for capsule of cryptococcus neoformans)
- Indian ink staining (used for capsule of cryptococcus neoformans)
- H&E (hematoxylin/eosin stain)

www.medicalcoupons.in

- PAS -Periodic acid Schiff.

DERMA JUNE 2018

1. Spot diagnosis based on the clinical image:



a) Secondary syphilis
b) Lupus vulgaris
c) Acanthosis nigricans
d) Scrofuloderma

Acanthosis nigricans is a **skin condition that causes a dark discoloration in body folds and creases**. It typically affects the armpits, groin and neck. Acanthosis nigricans is a skin condition characterized by areas of dark, velvety discoloration in body folds and creases. The affected skin can become thickened

In the given image:

→ Neck folds are involved
→ Hyperpigmented lesions
→ Thickened skin
→ Velvety appearance.

2. Drug of choice for syphilis:

a) Ceftriaxone
b) Erythromycin
c) Tetracycline
d) Benzathine penicillin

→ **DOC for syphilis is Benzathine penicillin except for Neurosyphilis - Procaine Penicillin (Aqueous / Crystalline Penicillin)**

www.medicalcoupons.in

3. Dermatophytes can affect:

a) Hair
b) Nail
c) Scalp
d) All of the above

A dermatophyte infection, also known as dermatophytosis or tinea, refers to a group of fungal infections that can affect **the skin, hair, and nails**. Tinea infections are one of the most common causes of superficial fungal infections around the world and are distinguished by the area of the body affected.

4. Wavelength of light produced by wood's lamp

a) 320nm
b) 360nm
c) 400nm
d) 480nm

A traditional Wood lamp is a low-output mercury arc covered by a Wood filter (barium silicate and 9% nickel oxide) and emits wavelength **320–450 nm (peak 365 nm)**. The lamp was invented in 1903 by a Baltimore physicist Robert W. Wood.

5. Acne vulgaris involves:

a) Sweat glands
b) Pilosebaceous glands
c) Sebaceous glands
d) Apocrine glands

Acne vulgaris is a common chronic skin disease involving **blockage and/or inflammation of pilosebaceous units (hair follicles and their accompanying sebaceous gland)**. Acne can present as noninflammatory lesions, inflammatory lesions, or a mixture of both, affecting mostly the face but also the back and chest.

www.medicalcoupons.in

OPTHAL JUNE 2018

1. Cornea derives its nutrition chiefly from:

a) Aqueous humour
b) Mucous layer
c) Corneal vessels
d) Perilimbal vessels

Unlike most tissues in the body, the cornea contains no blood vessels to nourish or protect it against infection. Instead, the cornea receives its nourishment from the **tears and aqueous humor** (a fluid in the anterior portion of the eye) that fills the chamber behind it.

2. Maximum cones are seen in:

a) Limbus
b) Fovea centralis
c) Macula lutea
d) Blind spot

The fovea centralis is **located in the center of the macula lutea**, a small, flat spot located exactly in the center of the posterior portion of the retina. As the fovea is responsible for high-acuity vision **it is densely saturated with cone photoreceptors**.

3. A 27-year-old, 38 weeks pregnant woman with height of 130 cm and a weight of 51 kg diagnosed case of cephalopelvic disproportion with fetal distress was posted for emergency cesarean section. A lower segment cesarean section was performed and a live male infant 1.8 kg with an APGAR of 8/9 was delivered. The eye of this newborn would have ?

a) Hypermetropic with regular astigmatism
b) Hypermetropia
c) Hypermetropic with irregular astigmatism
d) Myopia

- At birth, the eye is hypermetropic by +2 to +3D and usually becomes emmetropic by the age of 5 to years
- This is because of the smaller eyeball.
- Eyeball length at birth is 16mm.

www.medicalcoupons.in

4. **Campimetry is used to measure ?**

a) Field of Vision
b) Pattern of retina
c) Malignant melanoma
d) Squint

Perimetry or campimetry is one way to systematically test the visual field. It is the **systematic measurement of differential light sensitivity in the visual field** by the detection of the presence of test targets on a defined background.

5. Drug of choice in eye in trachoma endemic area ?

a) Erythromycin
b) Ganciclovir
c) Clindamycin
d) Azithromycin

Azithromycin is a macrolide antibiotic and is the drug of choice for trachoma. Plasma concentrations are low, but tissue concentrations are higher, giving it value in treating intracellular organisms. It has a long tissue half-life. A single dose is recommended.

6. Herbert's pits are seen in

a) Vernal conjunctivitis
b) Atopic conjunctivitis
c) Gonococcal conjunctiviti
d) Chlamydial conjunctivitis

Herbert's pits are a characteristic and pathognomonic finding of **trachoma**, and are caused by Chlamydia trachomatis.


7. As per WHO classification of Xerophthalmia, X2 is:

a) Night blindness
b) Conjunctival xerosis
c) Bitot's spots
d) Corneal xerosis

World Health Organization Re- Classification of Xerophthalmia Signs

Classification	Ocular Signs
XN	Night blindness
X1A	Conjunctival xerosis
X1B	Bitot's spots
X2	**Corneal xerosis**
X3A	Corneal ulceration Keratomalacia involving one- third or less of the cornea
X3B	Corneal ulceration -Keratomalacia involving one-half or more of the cornea
XS	Corneal scar
XF	Xerophthalmic fundus

8. **Which of the following is true about given condition ?**



a) Painless
b) Affects cornea
c) Sudden loss of vision
d) Bilateral in majority of cases

The given condition is Mooren's ulcer.

Mooren's ulcer:

- Affects cornea
- Painful condition
- Slow loss of vision
- Mostly unilateral
- Peripheral corneal ulceration
- Clear cut overhanging edge seems to be melting from periphery

www.medicalcoupons.in

9. Argon laser trabeculoplasty is done in:

a) Angle closure glaucoma
b) Angle recession glaucoma
c) Secondary glaucoma
d) Open angle glaucoma

The most common type of laser surgery performed **for open-angle glaucoma** is called Argon Laser Trabeculoplasty (ALT). The objective of the surgery is to help fluids drain out of the eye, reducing intra-ocular pressure that can cause damage to the optic nerve and loss of vision.

10. Pupil in acute iritis is:

a) Normal
b) Dilated
c) Constricted
d) Vertically oval

- **In acute iritis (inflammation of iris), iris is irritated and pupil constricts due to stimulus of irritation.**
- Vertically oval, mid-dilated, non-reacting pupil is seen in acute angle closure glaucoma.

11. Cause of sudden loss of vision in a diabetic is due to:

a) Neovascular glaucoma
b) Central retinal artery occlusion
c) Vitreous hemorrhage
d) None of the above

- **Commonest cause of sudden loss of vision in diabetic is vitreous hemorrhage**
- Slow loss of vision in diabetic is due to -
  → Cystoid macular edema
  → Neovascular glaucoma

12. Optic disc changes of retinitis pigmentosa:

a) Consecutive optic atrophy
b) Blurring of disc margins
c) Hyperemia of disc
d) No significant change

www.medicalcoupons.in

Retinitis pigmentosa (RP) is a group of hereditary disorders of the photoreceptors and retinal pigment epithelium (RPE) which gradually causes night blindness and progressive constriction of the visual field. Waxy optic disc pallor, arteriolar narrowing and hyalinization are found in almost all cases. **The optic disc in RP show optic atrophy, classically described as a 'waxy pallor' of the disc**, and is a **consequence of photoreceptor degeneration**.

PHYSIO JUNE 2018

1. Surfactant is secreted by which of the following

a) Type I pneumocyte
b) Type II pneumocyte
c) Sertoli cell
d) Leydig cell

Type I pneumocyte: Involved in gas exchange, makes up 80-90% of alveolar cells.

Type II pneumocyte: Secrete surfactant, makes up 5-10 % area alveolar cells.

Sertoli cell: Present in seminiferous tubules, responsible for nourishment spermatozoa during spermatogenesis, has FSH receptors & secrete inhibin.

Leydig cell: Produces testosterone in response to LH.

2. Which of the following receptors mediate stretch reflex:

a) Golgi tendon organ
b) Merkel's disc
c) Muscle spindle
d) Meissner's corpuscles

The classic stretch reflex involves the stretch receptor in **the skeletal muscle spindle** and its afferent Ia fiber in the posterior nerve root, which directly or indirectly excites the alpha motor neuron in the spinal cord.

www.medicalcoupons.in

3. Resting membrane potential of nerve fibre is close to isoelectric potential of:

a) Potassium ions
b) Sodium ions
c) Chlorideions
d) Magnesium ions

4. Which of the following hormones regulates blood levels of 1,25-OH-cholecalciferol positively:

a) Thyroxine
b) Parathormone
c) Insulin
d) Calcitonin

Serum levels of 1,25-dihyroxyvitamin D have little or no relationship to vitamin D stores **but rather are regulated primarily by parathyroid hormone levels**, which in turn are regulated by calcium and/or vitamin D. In vitamin D deficiency, 1,25-dihydroxyvitamin D levels go up, not down.

Parathyroid hormone (PTH), also called parathormone or parathyrin, is a hormone secreted by the parathyroid glands that regulates the serum calcium concentration through its effects on bone, kidney, and intestine.

5. Enteropeptidase enzyme is secreted by:

a) Stomach
b) Duodenum
c) Jejunum
d) Ileum

Enterokinase, also called Enteropeptidase, proteolytic enzyme (q.v.), secreted **from the duodenal mucosa**, that changes the inactive pancreatic secretion trypsinogen into trypsin, one of the enzymes that digest proteins.

www.medicalcoupons.in

6. Which of the following is the best stimulus for the release of vasopressin ?

a) Hypotonicity of extracellular fluid
b) Hypertension
c) Hypotension
d) Hypo-osmolality of extracellular fluid

The main physiological stimulus to vasopressin secretion is **rising plasma osmolality**, **though significant reductions in arterial blood pressure and blood volume** can also stimulate vasopressin secretion, by unloading of arterial baroreceptors.

7. Electrical impulses originates in:

a) SA node
b) AV node
c) Endocardium
d) Epicardium

The electrical signal starts in **a group of cells at the top of your heart called the sinoatrial (SA) node**. The signal then travels down through your heart, triggering first your two atria and then your two ventricles.

8. Inhibin is secreted by:

a) Leydig cells
b) Sertoli cells
c) Interstitial cells
d) Peg cells

Inhibin is a protein secreted by **the Sertoli cells in men and by the granulosa cells in women**. It inhibits the synthesis and release of the follicle-stimulating hormone in the pituitary gland and reduces the hypothalamic LH - releasing hormone content.

9. Centre controlling vomiting :

a) Apneustic centre
b) Pneumotaxic centre
c) Area Postrema
d) Hypothalamus

www.medicalcoupons.in

Area postrema is the center responsible for controlling vomiting it is located outside the blood brain barrier. and part of the chemoreceptor trigger zone, hence chemicals in blood can directly stimulate this area and induce vomiting.

10. Renin is secreted by:

a) Tubular cells
b) Macula densa
c) JG cells
d) All of the above

Renin is an enzyme secreted by **the juxtaglomerular cells of the kidney**. It interacts with aldosterone in a negative-feedback loop.

11. The rate at which the stomach empties into the duodenum depends on the type of food ingested. Food rich in carbohydrates leaves the stomach in a few hours. Protein-rich food leaves more slowly, and emptying is slowest after a meal containing fat. The rate of emptying also depends on the osmotic pressure of the material entering the duodenum. Hyperosmolality of the duodenal contents is sensed by "duodenal osmoreceptors" that initiate a decrease in gastric emptying, which is probably neural in origin. Which factor is responsible for increasing gastric motility?

a) Presence D1 fatty food in the stomach
b) Distension of stomach
c) Presence of tryptophan in stomach
d) Presence of acid in stomach

Promoters of gastric emptying :

- Chemical → Gastrin
- Mechanical → Gastric Distension.

12. The adrenal cortex is derived from mesoderm and begins posteromedial to the urogenital ridge. The adrenal medulla is derived from neural crest cells. These neural crest cells originate from the dorsal aorta. Clusters of chromaffin cells become the distinct medulla after birth. The adrenal gland first appears at 28-30 days post-conception. Adrenal medulla secretes?

a) Sex hormones
b) Thyroid stimulating hormones
c) Epinephrine
d) Glucocorticoid

www.medicalcoupons.in

The adrenal medulla, the inner part of an adrenal gland, controls hormones that initiate the flight or fight response. **The main hormones secreted by the adrenal medulla include epinephrine (adrenaline) and norepinephrine (noradrenaline)**, which have similar functions.

FORENSIC JUNE 2018

1. Rupture of the skin and subcutaneous tissue is seen in :-

a) Incised wound
b) Abrasions
c) Lacerated wound
d) Contusion

A laceration is the result of a shearing force and causes deeper skin tearing, through **the epidermis and sometimes through the dermis and subcutaneous tissues**.

2. In case of death in lock-up, the inquest is held by:-

a) A police officer
b) Magistrate
c) Panchayat officer
d) District attorney

Magistrate inquest (176 CrPC) is Conducted in these cases :

A. Asylum death

B. Borstal (juvenile home)

C. Custodial death

D. Dowry death

E. Exhumation / Encounter death

3. The time limit for ordering an exhumation in India is:-

a) 1 year
b) 10 years
c) 20 years
d) No limit

www.medicalcoupons.in

In India: No time limit - for exhumation [176 (3) CRPC]

Exhumation :

- It is the digging out of the body out of earth
- Authorization given only by magistrate
- Done in presence of -
  → Magistrate
  → Police
  → Doctor
- Preferred to start in early morning
- No time limit for exhumation
- 500 gms of soil sample taken.

4. First Post Morterm change in a dead body:-

a) Maggot formation
b) Putrefaction
c) Greenish discoloration of right iliac fossa
d) Mummification

- **First postmortem change in dead body is greenish discoloration of right iliac fossa**
- It's because of presence of caecum
- Its first external site of putrefaction (Right Iliac Fossa)
- 1st internal site → Intima of aortic lumen (Reddish brown discoloration)
- Most imp. bacteria involved in putrefaction → Clostridium welchi

5. Superimposition technique is used for:-

a) Skull
b) Pelvis
c) Femur
d) Ribs

Craniofacial superimposition is a technique used in the field of forensic anthropology **to assist in the analysis of an unknown skull**. The process involves superimposing an image of the recovered skull over an ante mortem image of the suspected individual.

6. Pure Aryans have which type of skull:-

a) Brachycephalic
b) Mesaticephalic
c) Dolicocephalic
d) All of the above

www.medicalcoupons.in

- Dolichocephalic → Aryans, Negroids Cephalic index (70. 75)
- Mesaticephalic → Europeans, Chinese -Cephalic index (75 - 80)
- Brachy cephalic → Mongoloids Cephalic index (80 - 85)

7. Gettler's test is positive in:-

a) Hanging
b) Poisoning
c) Strangulation
d) Drowning

A test that traditionally was used is the Gettler Chloride Test where **the amounts of chloride on each side of the heart were compared to determine if the person drowned in fresh or salt water.**

Gettler test not useful in-

→ Dry drowning
→ Hydrocution
→ Patent foramen oval
→ Putrefaction.

8. Rain drop pigmentation Is seen in:-

a) Arsenic poisoning
b) Phosphorous poisoning
c) Mercury poisoning
d) All of the above

Arsenic poisoning :

A - Anemia / Aldrich Mees line / Arsenophagist (Tolerate up to 300mg)

R - Rain drop pigmentation / Reinsch test / Red velvety mucosa

S - Sub endocardial hemorrhages / Sensory neuropathy

E - Eruptions

N - NAA (neutron activation analysis)

I - Imbibition of arsenic (Arsenic imbibed from surrounding soils after death)

C - Cumulative poison / Cholera like symptom / Chelation for Rx

www.medicalcoupons.in




9. A 40-year-old man has committed suicide by consuming some chemical he was able to procure from the paint-making factory he works at. The doctor notices the garlicky odor from the tissues and decided to preserve the hair for further investigation. Poisoning with which one of the following requires preservation of the hair?

a) Arsenic
b) Manganese
c) Phosphorous
d) Alcohol

Poison	Organ preserved
Nux vomica	Spinal cord
Aconite	Heart
Alcohol / any cerebral poison	Brain
Metal (chronic) - arsenic, antimony, thallium	Bone, hair, nail
Volatile poisoning gas)	Lungs

PATHO JUNE 2018

Heart failure cells are:-


a) Lipofuscin granules in cardiac cells
b) Pigmented hepatocytes
c) Pigmented alveolar marcophages
d) Pigmented cells in pancreas

www.medicalcoupons.in

A **siderophage** is a **hemosiderin-containing macrophage**. **Heart failure cells are siderophages** generated in the **alveoli of the lungs** of people **with left heart failure** c chronic pulmonary edema, when the high pulmonary blood pressure causes red blood cells to pass through the vascular wall.

2. Virchows triad includes all except:-

a) Endothelial injury
b) Stasis of blood flow
c) Hypercoagulability
d) Platelet thrombus



3. Mark correct statements regarding surfactant

a) Surfactant production starts at 16-18 wks of fetal life
b) Maximum secretion occurs at 34-35 wks
c) Increase surfactant production by insulin.
d) 30% Dipalmitoyl phosphatidyl choline in composition

- Surfactant is the fluid secreted by type 2 pneumocyte.
- Physiological function of surfactant is [↓ surface tension].
- Composition of surfactant - 70% phospholipids [DPPC]-Dipalmitoyl phosphatidyl choline AKA "lecithin" 30% fibrin, mucopolysaccharides etc.
- Surfactant covers water surface and prevent air - fluid interaction so, [↓ surface tension]

KEY POINTS:

→ Surfactant production starts at 18-20 weeks of fetal life
→ Secretion and action into alveoli occurs at 28 wks.
→ Maximum secretion occurs at 34-35 weeks
→ Increase surfactant production by thyroid, sex steroids, glucocorticoids [most imp. For lung maturation]

www.medicalcoupons.in

→ Decrease surfactant production by insulin.

4. PTT is prolonged in which of the following
A. Hemophilia A

B. Christmas disease

C. Von Willebrand disease

D. ITP

a) **A, B, C**
b) A, B, D
c) A, C, D
d) B, C, D

aPTT is used to monitor the activity of Intrinsic pathway.

Hemophilia :

- Most common type of hemophilia is Hemophilia A
- Hemophilia A - Factor VIII deficiency → aPTT

Von Willebrand disease :

- Associated with reduction in the half-life of Factor VIII - ↑ aPTT

Christmas disease :

- Factor IX deficiency - ↑ aPTT

ITP - Idiopathic Thrombocytopenic Purpura

- Disease associated with ↓ platelets, not a clotting factor problem.
- Parameters affected in ITP -
  → Increase bleeding time*
  → Decrease platelet count

5. AFP (alpha feta protein) is raised in:-

a) Renal carcinoma
b) Pancreatic carcinoma
c) Prostatic carcinoma
d) Hepatic carcinoma

www.medicalcoupons.in

Serum alpha fetoprotein (AFP) is the most widely used tumor marker in detecting patients with hepatocellular carcinoma. When elevated, the AFP is 75-91% specific, and values greater than 400 ng/mL are generally considered diagnostic of HCC in the proper clinical context, including appropriate radiologic findings.

6. Negri bodies are characteristic of:-

a) Tetanus
b) Rabies
c) Polio
d) AIDS

Negri bodies are seen in the Pyramidal neurons within Ammon's horn of the Hippocampus and Purkinje cells in the cerebellar cortex on postmortem of the rabid person.

7. Earliest feature of TB:-

a) Granuloma
b) Langerhan's Giant cells
c) Lymphocytosis
d) Caseation

Ingestion of mycobacterium tuberculosis inside the lung will cause stimulation in the no. of lymphocytes, that will cause **lymphocytosis [earliest feature of TB]**

First cell in the lung which will be affected by myco. TB infection is alveolar macrophage

Transmission of myco. TB - cough & sneeze (droplet infection)

8. Which of the following is not a component of Jones criteria for diagnosing acute rheumatic fever?

a) Pancarditis
b) Migratory polyrthritis
c) Erythema nodosum
d) Sydenham chorea

Jones criteria :

Major criteria

→ Migratory polyarthritis
→ Carditis

www.medicalcoupons.in

→ Erythema marginatum
→ Syndenham chorea
→ Subcutaneous nodules

Minor criteria

→ Arthralgia
→ Fever
→ First degree heart block
→ Elevated inflammatory markers (ESR, CRP).

9. MC site for amoebiasis:-

a) Liver
b) Caecum
c) Transverse colon
d) Sigmoid colon

Amebiasis may involve any part of the bowel, bu**t the cecum and the ascending colon are predilection sites**. The most common extra intestinal manifestation is **liver abscess** caused by hematogenous spread from the GI tract.

10. Crescentic glomerulonephritis is:-

a) FSGS
b) MGN
c) RPGN
d) All of the above

Crescentric glomerulonephritis, also known as **rapidly progressive glomerulonephritis**, is **a condition of the kidney characterized by a rapid loss of renal function**. On histological examination, it has the presence of numerous glomerular crescents (usually greater than 50%).

11. HLA is located on :

a) Short arm of chromosome -6
b) Long arm of chromosome -6
c) Short arm of chromosome -3
d) Long arm of chromosome -3

www.medicalcoupons.in

The human **major histocompatibility complex HLA is located on the short arm of chromosome 6**. It is known to be the most polymorphic genetic system in humans. The biological role of the HLA class I and class II molecules is to present processed peptide antigens.

12. Most common tumor of anterior mediastinum:-

a) Neurogenic cyst
b) Thymoma
c) Neurofibroma
d) Sarcoma

- **MC tumor of anterior mediastinum is THYMOMA**
- Thymoma has a definitive association with a particular type of muscle weakness disorder- Myasthenia gravis.
- Patients suffering from thymoma also going to have a relationship between this tumor and hematological condition which is given the name PRCA [pure red cell aplasia]

13. A 9 Yr old male child came with complaints of an Upper respiratory tract infection. On General examination, he was found to be anemic. The Pediatrician explained to his parents about the causes of Anemia in children. Which of the following is the most sensitive and specific initial laboratory test tc diagnose iron deficiency?

a) Serum iron levels
b) Serum ferritin levels
c) Serum transferrin receptor population
d) Transferrin saturation

In iron deficiency anemia, the earliest parameter that is effected is → decrease in stored iron [ferritin] → measured by serum ferritin levels.

Hence, estimation of Serum ferritin levels is most sensitive and specific test to diagnose iron deficiency anemia in the given options.

14. In beta thalassemia trait, electrophoresis shows:-

a) Increased HbF & normal HbA2
b) Normal HbF & normal HbA2
c) Increased HbF & increased HbA2
d) Decreased HbF & normal HbA2

www.medicalcoupons.in

If thalassemia is suspected, a hemoglobin electrophoresis will help diagnose the condition. The hemoglobin electrophoresis with beta thalassemia trait usually has **reduced or absent HbA, elevated levels of HbA2, and increased HbF**.

15. MC cause of nephrotic syndrome in adults:-

a) Good-pasture syndrome
b) Membranous GN
c) Minimal change disease
d) Focal segmental glomerulosclerosis

Primary causes of nephrotic syndrome are diseases that affect only the kidneys. The most common primary cause of nephrotic syndrome in adults is a disease called **focal segmental glomerulosclerosis (FSGS)**.

16. Preferred biochemical markers in patients presenting with myocardial infarction:-

a) Myoglobin
b) Cardiac specific Troponins
c) CK-MB
d) All of the above

Cardiac troponins T and I are the preferred markers for myocardial injury as they have the highest sensitivities and specificities for the diagnosis of acute myocardial infarction.

17. A known case of peptic ulcer disease presented with repeated episodes of vomiting's. The condition is associated with hypochloremic hypokalemic metabolic acidotic state. What is the most likely part involved in the given condition?

a) Antrum
b) Duodenum
c) Pylorus
d) Fundus

- Peptic ulcer disease , Carcinoma of the stomach → can result in Gastric outlet obstruction
- Peptic ulcer disease, MC site of obstruction - Duodenum [1st part]
- Carcinoma of stomach, MC site of obstruction - Pyloric region of stomach

www.medicalcoupons.in

- Most common cause of Gastric Outlet Obstruction in adult patient - Carcinoma of the stomach.

18. Gold standard method of diagnosing celiac disease is:-

a) Small bowel biopsy
b) Anti-endomysial antibodies
c) Blood picture
d) Biochemical test

Endoscopy with duodenal biopsy showing villous atrophy is the current gold standard for diagnosing celiac disease

19. Most common site of intracranial metastasis is from primary carcinoma of:-

a) Breast
b) Lungs
c) Stomach
d) Testes

Some differences are seen in the types of primary malignancies responsible for the brain metastasis in the two genders. **Lung cancer** is the most common source of brain metastasis in males, whereas breast cancer is the most common source in females.

20. Nephrotic range proteinuria is seen in:-

a) Arteriitis
b) Polycystic kidney disease
c) Pyelonephritis
d) Glomerulonephritis

- Healthy glomerulus is responsible for repulsion of proteins like Albumin
- Any kind of **inflammation affecting the glomerulus will lead to damage to glomerulus**, thus glomerulus will not function properly, **leading to proteinuria**. This type of proteinuria is given the name **"Sub-nephrotic proteinuria"**

Additional findings in patient with glomerulonephritis

- HTN
- Hematuria
- Oliguria

www.medicalcoupons.in

21. Deficiency of which vitamin causes subacute combined degeneration of spinal cord?

a) Vitamin B1
b) Vitamin B6
c) Vitamin B2
d) Vitamin B12

Subacute combined degeneration of the spinal cord can be caused by inadequate oral intake of **vitamin B12**, poor absorption of vitamin B12, or by the use of medications such as metformin, proton pump inhibitors, and nitrous oxide.

22. Transplant of kidney from mother to son is an example of:-

a) Autograft
b) Allograft
c) Isograft
d) Xenograft

Autograft	graft taken from one part of the body of an individual and transplanted onto another site in the same individual, e.g., skin graft
Isograft	graft taken from one individual and placed on another individual of the same genetic constitution, e.g., grafts between identical twins.
Allograft	graft taken from one individual placed on genetically non-identical member of the same species.
Xenograft	graft taken from one individual placed on an individual belonging to another species, e.g., animal to man.

23. Patients with Polycystic disease of kidney presents with all except:-

a) Hematuria
b) Hypertension
c) Renal failure
d) Erythrocytosis

Polycystic kidney disease patients presents with -

→ Hematuria
→ Due to activation of Renin - Angiotensin - Aldosterone System (RAAS) the patient will have HYPERTENSION.

www.medicalcoupons.in

→ Over period of time, renal mass will be reduced so patient will develop Renal failure, as the renal mass is reduced. There will be reduction in the secretion of Erythropoietin, leading to anemia NOT Erythrocytosis.

- **Adult PCKD has Autosomal dominant inheritance**
- **Child PCKD has Autosomal Recessive inheritance**

ANESTHESIA JUNE 2018

1. False statement about post-dural puncture headache (PDPH):

a) Breach of dura
b) Onset of headache is usually 12-72 hours following procedure
c) Commonly occipito - frontal in location
d) Headache is relieved in sitting standing position

Post Dural puncture Headache

- **Pathophysiology** → CSF leak → ICP ↓ → Traction on nerve fibres originating from pia mater
- **Site** → Occipital > Frontal > Retroorbital
- **Duration** → 7-10 days
- **Pathognomic sign** → Change in posture aggravates headache
- **Predisposing conditions -**
  → Dura cutting needle (spinal needle)
  → Most commonly seen with 22G Quincke needle.
  → Wide bore needle, Multiple attempts.
  → Pregnancy (In pregnancy dura mater is very fragile venous engorgement fragility increases, chances of PDPH is more).
  → Note- Early ambulation never predispose to PDPH ***

2. In status asthmaticus, anesthetic agent used as bronchodilator:-

a) Morphine
b) Thiopentone sodium
c) Ketamine
d) Halothane

Ketamine is a short-acting pentachlorophenol derivative that exerts bronchodilatory effects because it leads to an increase in endogenous catecholamine levels, which may bind to beta receptors and cause smooth

www.medicalcoupons.in

muscle relaxation and bronchodilation. Ketamine was used in the management of status asthmaticus in patients with respiratory failure who did not respond adequately to mechanical ventilation. **This agent improved airway resistance, particularly the lower airways, as well as improve lung compliance.**

3. Which of the following inhalational agent is the induction agent of choice in children?

a) Methoxyflurane
b) Sevoflurane
c) Desflurane
d) Isoflurane

Sevoflurane is less soluble than isoflurane and is very pleasant to breathe, which makes it an excellent choice for inhalational induction of anesthesia, particularly in children. The respiratory and cardiovascular effects of sevoflurane are very similar to isoflurane, but sevoflurane does not cause 'coronary steal'.

4. Compression ventilation ratio in case of Neonatal CPR:-

a) 3:1
b) 15:2
c) 30:2
d) 100:8

There should be a **3:1 ratio** of compressions to ventilations, with 90 compressions and 30 breaths to achieve approximately 120 events per minute. Thus, each event will be allotted approximately 1/2 second with exhalation occurring during the first compression following each ventilation.

5. Massive transfusion of stored blood result in:-

a) Citric acid toxicity
b) Hypokalemia
c) Metabolic acidosis
d) Hypercalcemia

Citric acid toxicity: Citrate is used as Preservative in stored blood.

Massive transfusion results in:

→ **Hyperkalemia**: Potassium gets released from stored blood after day 5 from the date of collection.

www.medicalcoupons.in

→ **Metabolic Alkalosis**: Citrate in blood is converted to bicarbonate in liver ,which causes metabolic alkalosis.
→ **Hypocalcemia**: Calcium in patient's blood binds to the chelating agent contained within the stored blood.

PSYCHIATRY JUNE 2018

1. Jet black pigmentation of tongue with tactile hallucinations is feature of:-

a) Heroin
b) Opium
c) Alcohol
d) Cocaine

COCAINE -

→ Derived from erythroxylum coca
→ Has Na blocking properties
→ Seizures
→ FREE BASING
→ Crack (smoked)
→ Jet black pigmentation of tongue

Intoxication-

→ Tactile hallucination - cocaine bugs formication or magnan phenomenon
→ Auditory hallucination
→ Delusion of persecution

2. Not a fundamental symptom of schizophrenia:-

a) Autism
b) Automatism
c) Association Disturbances
d) Ambivalence

The fundamental symptoms, which are virtually present through all the course of the disorder, are also known as the famous Bleuler's four A's

1. **A**utism (autistic thinking and behavior) – excessive fantasy thinking which is irrational and withdrawn behavior.
2. **A**mbivalence- inability to take a decision
3. **A**ffect disturbances -inappropriate affect/emotions
4. **A**ssociation disturbances- of thoughts such as formal thought disorders.

www.medicalcoupons.in

3. Neurotransmitter involved in depression:-

a) GABA and dopamine
b) Serotonin and GABA
c) Dopamine and serotonin
d) GABA and norepinephrine

The three neurotransmitters implicated in Depression are: **Dopamine**. **Norepinephrine**. **Serotonin**.

4. A girl normal language milestones spends her time seeing hel own hands and does not interact with others. Likely diagnosis is.

a) ADHD
b) Autism
c) Asperger's syndrome
d) Rett's syndrome

Asperger's Syndrome, **a form of Autism Spectrum Disorder**, is a developmental disorder. Young people with Asperger's Syndrome have a difficult time relating to others socially and their behavior and thinking patterns can be rigid and repetitive.

PHARMA JUNE 2018

1. Pharmacodynamics deals with:-

a) Mode of excretion of drug
b) Mechanism of action of a drug
c) Transport of drug across the biological membranes
d) Latency of onset

Pharmacodynamics is the branch of pharmacology dealing with **the mechanisms of action of drugs**. Pharmacodynamics involves the study of the biochemical and physiological changes produced by drugs in the body during the prevention and treatment of disease.

www.medicalcoupons.in

2. Drug of choice for paroxysmal supraventricular tachycardia is:-

a) Digoxin
b) Propranolol
c) Adenosine
d) Dilitiazem

Adenosine is the first-line medical treatment for the termination of paroxysmal SVT.

3. Malignant neuroleptic syndrome is caused by:-

a) Antidepressants
b) Anxiolytics
c) Antipsychotics
d) Anti-epileptics

Neuroleptic malignant syndrome comes about most likely, as a result of **"dopamine D2 receptor antagonism"**. **The first-generation antipsychotics work by inhibiting dopaminergic neurotransmission**. Their effectiveness is best when they block about 72% of the D2 dopamine receptors in the brain. Dopamine is a chemical substance (neurotransmitter) found in the brain and elsewhere in the central nervous system that acts to convey messages from one cell to another.

4. Treatment for Zollinger Ellison syndrome is:-

a) Climetidine
b) Omeprazole
c) Misoprostol
d) Aluminium hydroxide

Zollinger-Ellison syndrome managed or treated:

- **Medication**: Drugs called **proton pump inhibitors(Omeprazole)** reduce the production of stomach acid and help heal ulcers.
- **Surgery to remove gastrinomas.**
- **Chemotherapy**: Anticancer drugs to shrink gastrinomas and lower the levels of gastrin in the blood.

5. Proton pump inhibitors suppress gastric acid secretion by acting on:-

a) H+/Cl- ATPase pump
b) H+/K+ ATPase pump

www.medicalcoupons.in

c) Na+/K+ ATPase pump
d) Na+/K+/CI- ATPase pump

Proton pump inhibitors act by irreversibly blocking **the hydrogen/potassium adenosine triphosphatase enzyme system** (the H^+/K^+ ATPase, or, more commonly, the gastric proton pump) of the gastric parietal cells.

6. The drug aspirin has been used for secondary prevention of ischemic heart disease because, it:-

a) Inhibits TXA2 synthesis by platelets
b) Inhibit prostacyclin synthesis
c) Inhibits release of EDRF
d) Inhibit synthesis of endothelins

Aspirin produces its effects **through inhibition of thromboxane A_2** (TXA_2) production, while $P2Y_{12}$ antagonists attenuate the secondary responses to ADP released by activated platelets. The anti-platelet effects of aspirin and a $P2Y_{12}$ antagonist are often considered to be separately additive.

7. Drug of choice for central diabetes insipidus is:-

a) Insulin
b) Desmopressin
c) Diazoxide
d) Chlorpropamide

In patients with central DI, **desmopressin** is the drug of choice. A synthetic analogue of antidiuretic hormone (ADH), desmopressin is available in subcutaneous, IV, intranasal, and oral preparations.

8. A pediatric patient was prescribed erythromycin for prevention of recurrent rheumatic fever in a rural health center. the patient complained of a history of constipation, which medication should be avoided to prevent development of any arrhythmia?

a) Lubiprostone
b) Streptomycin
c) Ebastine
d) Cisapride

www.medicalcoupons.in

- Erythromycin is inhibitor of microsomal enzyme - CYP3A4
- **CAT** drugs increase QT interval (metabolized by CYP3A4).
  → **C**isapride
  → **A**stemizole
  → **T**erfenadine
- If administered with Erythromycin, metabolism is inhibited, drug concentration rises.
- At high plasma concentration, these drugs block K* channel in the heart leading to **QT prolongation →Torsades de pointes.**

9. Which of the following is not an alkylating agent:-

a) Melphalan
b) Cyclophosphamide
c) Busulfan
d) 5-Fluorouracil

Alkylating agents in the classic family include **nitrogen mustard, cyclophosphamide, chlorambucil, melphalan, busulfan, and ifosfamide**.

10. Which of the following drug is not metabolized by acetylation?

a) Isoniazid
b) Hydralazine
c) Phenytoin
d) Procainamide

Drugs metabolized by acetylation are **SHIP**

Sulfonamides

Hydralazine

Isoniazid

Procainamide

11. Therapeutic index of a drug is an indicator of:-

a) Potency
b) Safety
c) Efficacy
d) All of these

www.medicalcoupons.in

The therapeutic index (TI, also referred to as therapeutic ratio) is **a quantitative measurement of the relative safety of a drug**. It is a comparison of the amount of a therapeutic agent that causes the therapeutic effect to the amount that causes toxicity.

12. Beta-blockers are used in all of the following except:

a) Essential tremors
b) AV block
c) Angina pectoris
d) Migraine prophylaxis

Contraindications for beta blockers include:

- Abrupt discontinuations
- Acute bronchospasm
- Acute heart failure
- Asthma
- AV block
- Bradycardia
- Bronchitis
- Cardiogenic shock
- Cerebrovascular disease
- Chronic obstructive pulmonary disease (COPD)
- Diabetes mellitus
- Emphysema
- Hypersensitivity to beta blockers
- Hypotension
- Kidney failure
- Hepatic disease
- Myopathy
- Pheochromocytoma
- Psoriasis
- Stroke
- Vasospastic angina
- Wolff-Parkinson-White syndrome.

13. A 45 year old male was admitted to the ICU after being diagnosed with Sub arachnoid hemorrhage secondary to a ruptured aneurysm. The patient is a known case of stage 2 hypertension and was taking Ramipril for the same. Which of the following calcium channel blockers can be used in this case, which might help relieve the cerebral vasospasm?

a) Nimodipine
b) Felodipine
c) Amlodipine
d) Nitrendipine

www.medicalcoupons.in

Nimodipine (DHPs)- Blocks Ca2+ channel only in cerebral blood vessels which leads to Cerebral vasodilation is it used in subarachnoid hemorrhage (SAH).

14. A hypertensive patient was started on some anti-hypertensive drug but he developed dry cough. Which of the following drug can be responsible for this effect?

a) Calcium channel blockers
b) Beta blockers
c) ACE inhibitors
d) Diuretics

Side effects of ACE inhibitors may include:

→ Dry cough
→ Increased potassium levels in the blood (hyperkalemia)
→ Fatigue
→ Dizziness from blood pressure going too low
→ Headaches
→ Loss of taste

15. Mechanism of action of zileuton:-

a) Cyclooxygenase inhibitor
b) 5-lipooxygenase inhibitor
c) 5-phosphodiesterase inhibitor
d) Prevents mast cell degranulation

Zileuton is a 5-lipoxygenase inhibitor, which in turn, **inhibits the formation of leukotrienes B4, C4, D4, E4**. Limiting these leukotrienes, in turn, helps to reduce inflammation, edema, mucus secretion, and bronchoconstriction in the airways.

16. In the management of diabetes mellitus, lactic acidosis is caused by which of the following?

a) Tolbutamide
b) Glipizide
c) Pioglitazone
d) Metformin

www.medicalcoupons.in

A specific class of oral diabetes medication, called **biguanides**, **can cause a build up of lactic acid levels**. **Metformin (Glucophage) is one of these drugs**. It's used to treat diabetes and may also be prescribed for other conditions, such as renal insufficiency.


17. Antidote for alprazolam is:-

a) Protamine sulphate
b) Flumazenil
c) EDTA
d) BAL

Flumazenil, a **specific benzodiazepine antagonist**, is useful in reversing the sedation and respiratory depression that often occur when **benzodiazepines are administered to patients undergoing anesthesia or when patients have taken an intentional benzodiazepine overdose**. **Alprazolam belongs to a class of medications** called **benzodiazepines**. Benzodiazepines are medications that have anti-anxiety-producing properties.

RADIOLOGY JUNE 2018


1. What is marked by arrow?



a) Right atrium
b) Pulmonary trunk
c) Aortic knuckle
d) Superior vena cava

The aortic knob or knuckle refers to **the frontal chest x-ray appearance of the distal aortic arch** as it curves posterolaterally to continue as the descending thoracic aorta. It appears as a laterally-projecting bulge, as the medial aspect of the aorta cannot be seen separate from the mediastinum.

www.medicalcoupons.in



2. Floating water lily sign is seen in:

a) Lung hydatid
b) Bronchial adenoma
c) Lung abscess
d) Aspergilloma



The water-lily sign, also known as the camalote sign, is seen in **hydatid infections** when there is detachment of the endocyst membrane which results in floating membranes within the pericyst that mimic the appearance of a water lily.

3. String sign is seen in:

a) Crohn's disease
b) TB of ileocecal region
c) Idiopathic hypertrophic pyloric stenosis
d) All of the above

- The **gastrointestinal string sign** (also known as the **string sign of Kantor**) refers to the string-like appearance of a **contrast-filled bowel loop** caused by its severe narrowing.

www.medicalcoupons.in

- Originally used to describe the reversible narrowing caused by spasms in **Crohn disease**, it is now used for any severe narrowing of the bowel lumen, including that seen in **hypertrophic pyloric stenosis, gastrointestinal tuberculosis, carcinoid tumor** and **colon cancer**.

4. Dead bone on an X-ray looks ?


a) More radio-opaque
b) Radiolucent
c) Less radio-opaque
d) Not seen at all

In chronic osteomyelitis there is dead dense bone in center called sequestrum , which is more radio opaque.

5. Multiple punched out lesions in skull is seen in.

a) Acute leukemia
b) Thalassemia
c) Multiple myeloma
d) Ca prostate

A SKULL ROENTGENOGRAM demonstrating “punched-out” lesions of the calvarium is usually associated with the diagnosis of a number of malignant or systemic disorders. The disorders that most frequently come to mind are **metastatic carcinoma or multiple myeloma**.



www.medicalcoupons.in

PEDIA JUNE 2018

1. Earliest indication of sexual maturation in females:

a) Maturation
b) Thelarche
c) Pubarche
d) Menarche

In most girls, **breast budding** is the first visible sign of sexual maturation –**"Thelarche"** means **"the beginning of breast development**, followed closely by the initiation of the growth spurt. Shortly thereafter, pubic and axillary hair appears. Menarche generally occurs about 2 years after onset of breast development and when growth in height slows after reaching its peak.

2. A child of 4 years can do all of the following EXCEPT:

a) Copies Cross
b) Bridge with blocks
c) Copies Circle
d) Copies triangle

KEY FINE MOTOR MILESTONES	
Age	**Milestone**
4 months	Bidextrous reach (reaching out for objects with both hands)
6 months	Unidextrous reach (reaching out for objects one hand); transfers objects
9 months	Immature pincer grasp; probes with forefinger
12 months	Pincer grasp mature
15 months	Imitates scribbling; tower of 2 blocks
18 months	Scribbles tower of 3 blocks
2 years	Tower of 6 blocks; vertical and circular stroke
3 years	Tower of 9 blocks; copies circle
4 years	**Copies cross; bridge with blocks**
5 years	Copies triangle ; gate with blocks

3. Specific sign of kwashiorkor is:

a) Pitting edema
b) Weight loss
c) Flag sign
d) Muscle wasting

www.medicalcoupons.in

Kwashiorkor	Marasmus
• Because of protein deficiency	• Because of both protein and calorie deficiency
• Mc in 6 months to 3 yrs. of age	• Mc in infancy
• Subcutaneous fat preserved	• Not preserved
• Edema present	• Edema Absent
• Ribs are not very prominent	• Ribs are very prominent
• Lethargic	• Alert and irritable
• Mild or absent muscle wasting	• Sever muscle wasting
• Poor appetite	• Voracious feeder
• Fatty liver is common	• Fatty liver is uncommon
• Skin and hair changes are more common	• Skin and hair changes are Less common

4. Low osmolarity ORS has (as compared to WHO ORS):

a) Low glucose and high sodium
b) Low glucose and low sodium
c) Low glucose and low potassium
d) Low glucose and high potassium

Composition	ReSoMal (mmol/L)	Standard ORS (mmol/L)	Reduced osmolarity ORS
Glucose	125	111	75
Sodium	45	90	75
Potassium	40	20	20
Chloride	70	80	65
Citrate	7	10	10
Magnesium	3	-	-
Zinc	0.3	-	-
Copper	0.045	-	-
Osmolarity (mOsm/L)	300	311	245

5. Jaundice at birth or within 24 hours of birth is due to:

a) Erythroblastosis fetalis
b) Congenital hyperbilirubinemia
c) Biliary atresia
d) Physiological jaundice of newborn

www.medicalcoupons.in

- **Erythroblastosis fetalis** is **hemolytic anemia in the fetus** (or neonate, as erythroblastosis neonatorum) caused by transplacental transmission of maternal antibodies to fetal red blood cells. The disorder usually results from incompatibility between maternal and fetal blood groups, often Rho(D) antigens.
- Destruction of the red blood cells (hemolysis) can be rapid in a fetus. As a result, the fetus will not receive enough oxygen, which may lead to anemia, other illnesses, or even death.
- As hemolysis continues, the fetus will rapidly attempt to produce more red blood cells. However, these cells new red blood cells are often immature and are unable to function fully.
- Because the body produces red blood cells in the liver and spleen, this overproduction can sometimes cause these organs to enlarge.
- When a newborn has this condition, it is known as hemolytic disease of the newborn.
- As the immature red blood cells continue to break down, bilirubin, which is a by-product of the breakdown of red blood cells, builds up. The excess amounts of bilirubin circulating in the newborn's body will lead to jaundice, where the skin and eye whites of the infant turn yellow.

6. 1 month old newborn presents with conjugated hyperbilirubinemia, intrahepatic cholestasis and high alkaline phosphatase. Eosinophilic, PAS positive cytoplasmic granules were seen in liver biopsy specimen. These are suggestive Alpha 1 AT deficiency and usually presents as neonatal cholestasis. What is the MC cause of jaundice in this newborn?

a) Hypoplasia of biliary tract
b) Neonatal hepatitis
c) Choledochal cyst
d) Physiological

- MC cause of cholestatic jaundice in newborn is Biliary atresia > Neonatal hepatitis.
- MCC of jaundice in 1st 24 hrs of life - Hemolytic disease of newborn.
- MCC of jaundice in 24 hrs to 72 hrs of life - Physiological jaundice.

7. MC congenital defect in Down's syndrome:

a) PDA
b) ASD
c) Atrioventricular septal defect
d) VSD

- **MC congenital heart defect in Down's syndrome is Atrioventricular septal defect/ Endocardial cushion defect.**
- MC congenital heart defect in turner syndrome is Bicuspid aortic valve followed by coarctation of aorta.
- MC congenital heart defect in DM is transposition of great arteries. Most characteristics in DM is sacral agenesis.

www.medicalcoupons.in

- MC congenital heart defect in Noonan syndrome is pulmonary stenosis.
- MC congenital heart defect in child born to mothers with lithium intake is Ebstein anomaly.
- MC congenital heart defect in Rubella is Patent ductus arteriosus.
- Overall MC congenital heart defect is VSD
- MC ASD is foramen secundum.
- MC ASD in Down's syndrome is foremen primum.

8. **MC malignancy in children is**:

a) Retinoblastoma
b) Leukemia
c) Wilms tumor
d) Neuroblastoma

- Leukemia is the most common malignant neoplasm in childhood
- MC malignancy in children is ALL > Brain tumor > neuroblastoma > Wilms tumor.
- B cell ALL is more common than T- Cell ALL

9. A malignant tumour of childhood that frequently metastasizes to bone marrow most often is:

a) Wilms tumor
b) Neuroblastoma
c) Adrenal gland tumors
d) Granulosa cell tumor of ovary

- Neuroblastoma is the most common intra-abdominal and extracranial solid tumor in children, accounting for 7-8% of all cancers.
- Metastasis present in 60-70% children, are usually to the skeleton (facial bones, skull), bone marrow and lymph nodes.

10. A 2 year old male child was brought by the parents to the pediatric department with loose stools from last 3 days. On examination she has signs of severe dehydration. The intern on duty was asked by the JR to calculate the volume of fluid to be given in the first 3 hrs if her weight is 10 kg.

a) **1000 ml**
b) 100 ml
c) 200 ml
d) 500 ml

www.medicalcoupons.in

In a 2 year old child with severe dehydration, fluid given over 1st 30 minutes is 30 mL/kg, over next 2.5 hrs is 70mL/kg: So fluid given over 3 hrs is 100mL/kg. In this case, as weight of child is 10 kg, amount of fluid to be given is 10x100=1000mL.

11. A 1-year-old male child present to the hospital with a fever associated with neck stiffness. You are suspecting meningitis. Which of the following is the most common cause of meningitis in this age group?

a) Klebsiella
b) Streptococcus pneumoniae
c) Staphylococcus aureus
d) Mycobacterium tuberculosis

IN INDIA :

Age group	Organism
Neonate	Acinetobacter, E.coli
1 month to 2 years	Strep. pneumoniae
Older children	Strep. Pneumoniae, N. meningitidis

IN WORLD :

Neonate	Group B strep, E.coli, Listeria
Infant and children	Strep. Pneumoniae, N. meningitidis, H. influenza
Adolescent and young adults	Strep Pneumoniae, N. meningitidis

12. Best method to diagnose HIV in infancy is:

a) Western blot
b) ELISA
c) PCR
d) All

WHO recommendation for HIV testing:

<18 months	HIV DNA or RNA PCR or HIV culture
>18 months	IgG antibody to HIV by ELISA and western blot test.

www.medicalcoupons.in

OBG JUNE 2018

1. Size of uterus (in inch):

a) 4x2x1
b) 3x2x1
c) 4x3x1
d) 5x4x2

UTERUS :

- **8 cm x 6 cm wide x 3-4 cm**, weight 50-80 g.
- It Consists of:
  - → body
  - → isthmus
  - → cervix

2. Pearl index is used to calculate:

a) Accidental pregnancies
b) Population
c) Fertility rate
d) Abortions

The Pearl Index is defined as **the number of contraceptive failures(accidental pregnancies) per 100 women-years of exposure**, and uses as the denominator the total months or cycles of exposure from the initiation of the product to the end of the study or the discontinuation of the product.

3. Progesterone of choice in emergency contraception:

a) Micronized progesterone
b) Norgesterone
c) Levonorgesterol
d) DMPA

Only the **progestin levonorgestrel** has been studied for use as emergency contraception. It is marketed as Plan B.

www.medicalcoupons.in

4. **Most likely to be associated with vaginal pH of 4**:-

a) Atrophic vaginitis
b) Trichomanas vaginatis
c) Candidal vaginitis
d) Gardenella vaginitis

Vaginal pH can be tested using a narrow-range pH paper. A pH above 4.5 suggests infections such as bacterial vaginosis or trichomoniasis (pH 5-6) and helps to exclude vulvovaginal candidiasis (**pH 4-4.5**).

5. Which one of the following in NOT a characteristic feature of bacterial vaginosis?

a) Presence of clue cells
b) Thick white discharge
c) Vaginal pH > 4.5
d) Fishy odour

Bacterial Vaginosis :

- Characterized by a pungent vaginal **"fish-like" odor** and a thin, **off-white/gray**, homogenous, discharge that coats the vaginal wall. Does NOT usually cause dysuria, dyspareunia, pruritus, or vaginal vulvar inflammation.
- The best initial test is microscopic examination. Diagnosis is supported by the presence of **clue cells on wet mount**.
- Other supportive findings include a **vaginal pH >4.5** and a positive whiff-amine test.
- Treated with either metronidazole or clindamycin for seven days.
- Routine screening and treatment of all pregnant women with asymptomatic bacterial vaginosis is NOT recommended.

6. Increased LH: FSH ratio is seen in

a) Premature menopause
b) Sheehan syndrome
c) PCOD
d) Turner's syndrome

- Premature menopause → LH↓ & FSH ↓
- Sheehan syndrome → < 1 LH & FSH because of pituitary necrosis
- PCOD → LH: FSH ratio is ↑↑ (i.e 2:1) or (3:1)
- Turner's syndrome → ovarian dysgenesis LH ↑ & FSH ↑
-

7. Which is not classified as primary amenorrhea:

a) Mullerian agenesis
b) Turner syndrome
c) Sheehan's syndrome
d) Turner's syndrome

Sheehan's syndrome causes secondary amenorrhea.

CAUSES OF PRIMARY AMENORRHEA :

- Mullerian agenesis/hypoplasia
- Imperforate hymen
- Vaginal agenesis
- Cervico vaginal atresia
- Endocrinological causes
- Hypogonadotropic hypogonadism
- Primary ovarian failure
- Congenital adrenal hyperplasia
- Constitutional delay
- Hypothyroidism
- Genetic causes
- Turner syndrome
- Pure gonadal agenesis
- Androgen insensitivity syndrome

8. A 23 year old woman presents with irregular cycles. Her cycles come at intervals of 35-45 days, last for 3 - 4 days with mild to moderate flow, no passage of clots and no dysmenorrhea. Which of the following describes this condition?

a) Amenorrhoea
b) Oligomenorrhoea
c) Polymenorrhoea
d) Menorrhagia

Oligomenorrhea is **a condition in which you have infrequent menstrual periods**. It occurs in women of childbearing age. Some variation in menstruation is normal, but a woman who regularly goes **more than 35 days without menstruating** may be diagnosed with oligomenorrhea.

www.medicalcoupons.in

9. A 50 year old woman diagnosed with an ovarian malignancy most likely has?

a) Granulosa cell tumour
b) Dermoid cyst
c) Mucinous cystadenoma
d) Serous cystadenoma

Ovarian tumors:

Epithelial cell tumor	Germ cell tumor	Sex cord tumor
M/C tumor (90%)	5-8%	3%
Types: • Serous (75-80%) (M/C) • Endometrial (8-10%) • Mucinous (5%) • Clear cell • Brenner tumors	• Teratoma • Yolk sac tumor • Endodermal sinus tumor • Choriocarcinoma • Dysgerminoma • Dermoid cyst	Estrogen Secreting tumors → Granulosa cell → Fibroma → Thecoma • Androgen secreting Tumor → Sertoli cell tumor → Leydig cell → Hilus cell tumor

10. Which is not associated with fibroid:

a) Pelvic mass
b) Amenorrhoea
c) Infertility
d) Menstrual irregularity

Symptoms of fibroid

→ M/C presentation in a fibroid is asymptomatic
→ M/C symptom is menstrual disturbances i.e progressive menorrhagia
→ Pressure symptom
→ Infertility
→ Pain
→ Polycythemia
→ Hypoglycemia & hypokalemia.

11. Condyloma acuminate is caused by:

a) HSV
b) HBV
c) HPV
d) HIV

www.medicalcoupons.in

- **Condyloma acuminate (genital warts) → HPV 6,11**
- Cancer cervix HPV 16, 18
- Prevention → Gardasil vaccine prevents HPV types 6,11,16,18 (Quadrivalent)
- RX → Podophyllin resin + Cryotherapy

12. Ferning pattern of drying cervical mucus suggests the action o**f**

a) Estrogen
b) Progesterone
c) Prolactin
d) 17-ketosteroids

Ferning occurs due to the presence of sodium chloride in mucus under **estrogen effect**. When high levels of estrogen are present, just before ovulation, the cervical mucus forms fern-like patterns due to crystallization of sodium chloride on mucus fibers. This pattern is known as **arborization** or 'ferning'.

13. Which of the following is treatment of choice for Trichomonas vaginalis?

a) Amikacin
b) Metronidazole
c) Chloramphenicol
d) Ampicillin

Infections with the sexually transmitted protozoan Trichomonas vaginalis are usually treated with **metronidazole**, a 5-nitroimidazole drug derived from the antibiotic azomycin. Metronidazole treatment is generally efficient in eliminating T. vaginalis infection and has a low risk of serious side effects.

14. A liquid media used to suppress commensal bacteria while allowing the pathogen to remain viable and grow. Which of the following is an enrichment media?

a) Alkaline peptone water
b) Monsour's taurocholate Tellurite peptone water
c) Selenite F broth
d) All of the above

Enrichment media

- LIQUID culture media
- These media are used to suppress commensal bacteria while allowing the pathogen to remain viable and grow.

www.medicalcoupons.in

- example-

→ Alkaline peptone water - V. cholerae
→ Selenite F-broth - Salmonella, shigella
→ Monsour's taurocholate tellurite peptone water - V. cholerae

BIO CHEM JUNE 2018

1. Km increases, but Vmax remains same. This is what Inhibition:

a) Competitive
b) Non-competitive
c) Irreversible
d) Uncompetitive

Effect on Km and Vmax for different type of inhibitors:

Type of inhibitor	Km	Vmax
Competitive	Increased	Same
Non Competitive	Same	Decreased
Uncompetitive	Decreased	Decreased

2. Cofactor for xanthine oxidase:

a) Zinc
b) Copper
c) Molybdenum
d) Selenium

The most important function of **Molybdenum** is as a cofactor for **xanthine dehydrogenase/oxidase, aldehyde oxidase, and sulfite oxidase**.

3. Which of the following correctly defines Isoenzymes?

a) Forms of the same enzyme that catalyze different reaction
b) Forms of the same enzyme that catalyze same reaction
c) Forms of the different enzymes that catalyze different reaction
d) Forms of the different enzymes that catalyze same reaction

www.medicalcoupons.in

- **Isozymes/Isoenzymes** are different forms of same enzyme, catalyzing same reaction in same species.
- Example:- There are five isozymes of Lactate Dehydrogenase (LDH-1 to LDH-5), All convert Pyruvate to Lactate.

4. **Ammonia is detoxified in brain by**:

a) Creatinine
b) Uric acid
c) Glutamine
d) Urea

- **The transport form of Ammonia in blood and from brain is Glutamine**.
- The answer is not glutamate because free ammonia is trapped by glutamine. Glutamate is formed via transamination of amino acids, not from free ammonia.

5. Atherosclerosis involves the formation of lipid rich plaques in the intima of arteries which can narrow the coronary arteries, leading to decreased blood flow and hypoxia of cardiac cells. Which of the following amino acid is known to be associated with Atherosclerosis?

a) Methionine
b) Cysteine
c) Homocysteine
d) Leucine

HOMOCYSTINURIA :

Homocysteine accumulation lead to stroke, atherosclerosis, MI, Pulmonary embolism. Homocysteine binds with the endothelium of the blood vessels and activates signaling pathways which leads to the release of pro-inflammatory molecules. This could result in the following:

→ Extensive atheroma formation at a young age affecting many arteries but not the coronary arteries
→ Intravascular thrombosis

6. Blood glucose levels cannot be augmented by mobilization of muscle glycogen due to lack of:

a) G-6-P dehydrogenase
b) G-6-phosphatase
c) Aldolase
d) Glucokinase

www.medicalcoupons.in

- Muscles lack the enzyme **Glucose-6-phosphatase**, that's why muscles cannot maintain blood glucose levels.
- Liver cells have this enzyme. And therefore liver can only maintain blood glucose.

7. Glucose can be synthesized from all except

a) Amino acids
b) Glycerol
c) Acetoacetate
d) Lactic acid

Gluconeogenesis (literally, "formation of new sugar") is the metabolic process by which glucose is formed from noncarbohydrate sources, **such as lactate, amino acids, and glycerol**.

8. Which is branching enzyme?

a) Glycogen synthetase
b) Glucose-6 phosphatase
c) Amylo-1, 4-1, 6-transglycolase
d) Glycogen Phosphorylase

- Branching enzyme - alpha (1-6) extend with alpha (1-4)
- Another name amylo (1-4) (1-6) transglycosylase.

9. Most Common enzyme deficient in galactosemia:

a) Galactose-1-phosphate uridyl transferase/GALT
b) Galactosidase
c) UDP galactose epimerase
d) Galactokinase

Classic galactosemia (type 1) - the most common and severe type, caused by mutations in the GALT gene, and characterized by a complete deficiency of an enzyme called **galactose-1-phosphate uridyl transferase (GALT)**.

www.medicalcoupons.in

10. Essential fatty acid:

a) Citric acid
b) Linoleic acid
c) Stearic acid
d) Palmitic acid

Essential Fatty Acid :

- Omega-3 Fatty Acid.
- Alpha-Linolenic Acid.
- Linoleic Acid.
- Eicosapentaenoic Acid.
- Docosahexaenoic Acid.
- Polyunsaturated Fatty Acid.
- Lipids.
- Fatty Acids.

11. Myocardium normally utilizes:

a) Glucose
b) Lactose
c) Fatty acid
d) Glycogen

	FED	FASTING	STARVATION
BRAIN	GLUCOSE	GLUCOSE	KETONE BODIES
HEART	FATTY ACIDS	FATTY ACIDS	KETONE BODIES
LIVER	GLUCOSE	FATTY ACIDS	AMINO ACIDS
MUSCLE	GLUCOSE	FATTY ACIDS	FATTY ACIDS & KETONE BODIES
ADIPOSE TISSUE	GLUCOSE	FATTY ACIDS	FATTY ACIDS
RBC	GLUCOSE	GLUCOSE	GLUCOSE

12. Apoprotein for chylomicron remnant:-

a) Apo E
b) Apo C1
c) Apo A2
d) Apo A1

www.medicalcoupons.in

APOPROTEIN - PROTEIN PRESENT IN LIPOPROTEIN

Lipoprotein	Protein
Chylomicron	APO B48
Chylomicron remnant	APO B48 + APO E
VLDL	APO B100
IDL	APO B100 APO E
LDL	APO B 100 + APO
HDL	APO A, C, E

13. Energy reserve of the body is:

a) Carbohydrate
b) Triglyceride
c) Proteins
d) Fibres

The major energy reserve in the body is found stored as **fat (triglyceride) in the adipose tissue**.

14. Chaperones are:

a) Purine metabolism mediators
b) Antigen presenting cells
c) Mediators of post-translational assembly of protein complexes
d) None of the above

CHAPERONES-FOLDING/SHAPE /ASSEMBLY OF PROTEIN

- POST-TRANSITIONAL CHANGE
- They facilitate and favour the interactions on the polypeptide surfaces to finally give the specific conformation of a protein.

15. Vitamin K is required for:

a) Hydroxylation
b) Oxidation
c) Transamination
d) Carboxylation

www.medicalcoupons.in

Vitamin K :

- Act as coenzyme in Carboxylation of clotting factor 2,7,9,10
- Only fat-soluble vitamin with coenzyme function.

ENT JUNE 2018

1. Most common manifestation of nasopharyngeal carcinoma

a) Epistaxis
b) Headache
c) Nasal obs
d) Cervical lymphadenopathy

- Nasopharyngeal Carcinoma m/c site is Fossa of Rosenmuller
- **Ca nasopharynx m/c presentation - neck mass (cervical lymph node metastasis) 75% of patients.**
- It causes U/L serous otitis media - U/L conductive hearing loss.

2. A 14 year old Male presented with history of fever since 2 days, unable to swallow the food with muffled voice. On examination it is noted right tonsil is shifted to midline. What is the diagnosis:

a) Quincy
b) Acute tonsillitis
c) Para pharyngeal abscess
d) Acute retropharyngeal abscess

If tonsil is enlarged / hypertrophied and reaching midline and soft palate is moved on other side it is a case of quincy but, if tonsil itself is pushed to the midline it is a case of parapharyngeal abscess.

3. Adam's apple seen in boys is because of:

a) Hyoid bone
b) Tracheal rings
c) Thyroid cartilage
d) Cricoid cartilage

- **Prominence of Thyroid cartilage (because of sharp angle) in males is called Adam's apple.**

www.medicalcoupons.in

- Thyroid cartilage with right and left lamina both meet al midline and form an angle of 90 in males, 120 in females.
- At rest it lies at level of C3.

4. All of the following are seen in Meniere's Disease except:

a) Tinnitus
b) Vertigo
c) Fullness of ear
d) Ear Pain

- Meniere's disease also called endolymphatic hydrops is a disorder of the inner ear where the endolymphatic system is distended with endolymph.
- It is characterized by
  → Vertigo
  → Sensorineural hearing loss
  → Tinnitus
  → Aural fullness

5. 34 yrs old female presents with tinnitus, vertigo and fullness in ear. Likely Diagnosis:

a) Meniere's Disease
b) Otosclerosis
c) Ototoxicity
d) Noise Induced Hearing Loss

- Meniere's disease also called endolymphatic hydrops is a disorder of the inner ear where the endolymphatic system is distended with endolymph.
- It is characterized by
  → Vertigo
  → Sensorineural hearing loss
  → Tinnitus
  → Aural fullness

6. Most common cause of SNHL

a) Labyrinthitis
b) Presbycusis
c) Meniere's disease
d) Vestibular Schwannoma

www.medicalcoupons.in

Presbycusis :

- Most common cause of SNHL.
- Associated with physiological aging process in the ear
- Bilateral slopping curve
- Usually manifests at the age of 65 years, (old age) but may do so early if there is a hereditary predisposition, chronic noise exposure or generalized vascular disease.

7. Most common site for laryngeal web is

a) Supraglottic
b) Subglottic
c) Anterior Glottis
d) Posterior Glottis

Laryngeal web

- It is due to incomplete recanalization of larynx -most site is anterior glottis
- Rx- Excision or Co2 laser excision. (depends upon thickness of the web).

8. Lifesaving muscle of vocal cords is:

a) Posterior cricoarytenoid
b) Lateral cricoarytenoid
c) Transverse arytenoid
d) Thyroarytenoid

- **Safety muscle/lifesaving muscle of vocal cord is Posterior cricoarytenoid because this muscle is the only abductor.**
- Tensor of vocal cord (1 - cricothyroid, 2-vocalis).

9. All of the following is seen in superior laryngeal nerve palsy EXCEPT:

a) Aspiration
b) Stridor
c) Bowed vocal cords
d) Loss of pitch

Superior laryngeal nerve palsy causes paralysis of cricothyroid muscle, and ipsilateral anesthesia of the larynx above the vocal cords.

www.medicalcoupons.in

Clinical features-

- Voice is weak and pitch cannot be raised with decreased ability to sing.
- Anaesthesia of the larynx on one side may pass unnoticed or cause occasional aspiration.
- Presence of both paralysis and bilateral anaesthesia. Causes inhalation of food and pharyngeal secretions gives rise to cough and choking fits. Voice is weak and husky. –
- **Stridor** is not seen In superior laryngeal nerve palsy **is seen in B/L recurrent laryngeal nerve palsy.**

10. Most common cause of facial nerve palsy:

a) Idiopathic Bell's palsy
b) Herpes zoster oticus
c) Mastoid surgery
d) Chronic suppurative Otitis media

Bell's palsy is the most common cause of facial paralysis, although its exact cause is unknown. Generally, Bell's palsy affects only one side of the face; however, in rare cases, it can affect both sides.

11. Rhinolalia clausa occurs in all except

a) Nasopharyngeal polyp
b) Nasopharyngeal carcinoma
c) Cleft palate
d) Nasopharyngeal angiofibroma

Rhinolalia clausa

- Abnormal speech due to obstruction in nose
- Hyponasal speech
- Causes:
  → Nasopharyngeal polyp
  → Nasopharyngeal cancer
  → Nasopharyngeal angiofibroma, adenoids

12. Cold water is not used for ear cleaning because ?

a) It will make the wax hard
b) Damage to tympanic membrane
c) Caloric stimulation caused by cold water
d) It will cause infection

www.medicalcoupons.in

Temperature for ear syringing is 37°c , because if you decrease or increase the temperature we can stimulate lateral semicircular canal in the middle ear and patient will have vertigo and nystagmus **(caloric stimulation).**

13. Otomycosis is most commonly caused by:

a) Actinomycetes
b) Aspergillus niger
c) Mucor
d) Candida albicans

Otomycosis / Singapore ear, fungal common causative organism **Aspergillus niger**

Other cause :

- Aspergillus fumigatus
- Candida albicans

14. A 3 yrs old child presents with c/o fever, barking cough and stridor for 2days, what is the diagnosis

a) Acute tonsillitis
b) Acute Epiglottitis
c) Croup
d) Adenoiditis

Croup - Acute laryngo tracheo bronchitis.

Causative - Parainfluenza virus type 1 and 2

Child - 3 month to 3 years

C/F - fever, cough (barking cough - Croup)

Stridor - expiratory

X-ray - steeple sign Rx symptomatic

15. CSF Otorrhea is due to involvement / trauma of ?

a) Cribriform plate
b) Petrous temporal bone
c) Parietal bone
d) Tympanic membrane

www.medicalcoupons.in

- CSF otorrhea - CSF coming from ear due to trauma to petrous temporal bone.
- CSF Rhinorrhea - CSF watery discharge from nose

16. Eustachian tube develops from:

a) 2nd and 3rd pharyngeal pouch
b) 1st pharyngeal pouch
c) 2nd pharyngeal pouch
d) 3rd pharyngeal pouch

Pharyngeal Pouch	Derivative
1	Tympanic (middle ear) cavity, auditory (eustachian) tube
2	Palatine tonsils, tonsillar fossa
3	Inferior parathyroid gland, thymus
4	Superior parathyroid gland, ultimobranchial body (parafollicular C cells of thyroid)

17. Recruitment phenomenon is seen in:

a) Otitis media with effusion
b) Acoustic nerve schwannoma
c) Meniere's disease
d) Otosclerosis

- Recruitment is phenomenon of abnormal growth of loudness.
- Recruitment is typically seen in lesions of the cochlea (e.g. Ménière's disease and presbycusis) and this helps to differentiate a cochlear from a retrocochlear sensorineural hearing loss.

18. Tobey-Ayer test is done for:

a) Acantholysis
b) Hemoglobinuria
c) Ketosis
d) Lateral sinus thrombosis

The Tobey-Ayer test is used for **lateral sinus thrombosis by monitoring cerebrospinal fluid pressure** during a lumbar puncture.

www.medicalcoupons.in

19. Laryngocele arises from ?

a) Subglottis
b) Anterior commissure
c) Saccule
d) True glottis

Laryngocele is basically a herniation of the mucosa of the laryngeal ventricle (Morgagni's ventricle) arising usually from **the saccular region.**

20. Angiofibroma bleeds profusely because:

a) It has multiple sites of origin
b) It lacks a capsule
c) Vessels lack a contractile component
d) None of the above

- Angiofibroma is made of vascular and fibrous tissues
- Vessel are endothelial lined spaces with no elastic or muscle coat
- Vessels lose ability to contract , thus bleed profusely.

21. Young's operation is done for.

a) Atrophic rhinitis
b) Vasomotor rhinitis
c) Antrachonal polyp
d) Allergic rhinitis

Young's operation is a surgery designed for the **treatment of atrophic rhinitis**, first described by Anderten Young in 1967.

22. Mikulicz cells are seen in:

a) Rhinoscleroma
b) Rhinosporidiosis
c) Rhinophyma
d) Rhinitis

The Mikulicz cell is a large macrophage with clear cytoplasm that contains the bacilli; this cell is specific to the lesions in **rhinoscleroma**.

www.medicalcoupons.in

23. MC malignancy of maxillary antrum:

a) Muco-epidermoid Carcinoma
b) Adeno-cystic Carcinoma
c) Adeno Carcinoma
d) Squamous cell Carcinoma

M/c malignancy of paranasal sinus and nasal cavity → Squamous cell carcinoma

Most common Paranasal sinuses involved → maxillary sinus

Associated with occupation:

Furniture industry → Adeno ca.

Nickel industry → Squamous cell ca.

MEDICINE JUNE 2018

1. A 44 year old female presenting to the opd with complaints of fatigue, weakness, and shortness of breath with minimal activity. Her friends and family have told her she appears pale, and combined with her recent symptoms she has decided to get checked out. She also states that she has noticed her hair and fingernails becoming extremely thin and brittle, causing even more concern. The patient first started noticing these symptoms a few months ago and they have been getting progressively worse. Upon initial assessment, her mucosal membranes and conjunctivae are pale. Considering the above scenario the most sensitive and specific test to diagnose this particular type of deficiency anemia is:

a. Serum iron levels
b. Serum ferritin levels
c. Serum transferring receptor population
d. Transferrin saturation

- **This is the classic presentation of Iron deficiency anemia**
- Serum ferritin levels is the most sensitive and specific test to diagnose iron deficiency anemia
- Under steady-state conditions, the serum ferritin level correlates with total body iron stores; thus, the serum ferritin level is the most convenient laboratory test to estimate iron stores.

www.medicalcoupons.in

2. True for von-willebrand disease

a. Normal partial thromboplastin time
b. Decreased platelets
c. Normal prothrombin time
d. Normal bleeding time

The prothrombin time **(PT) is normal in VWD (von-willebrand disease)**

3. Which Is NOT true about iron deficiency anemia?

a. Hyper-segmented neutrophils
b. Microcytosis precedes hypochromia
c. MCHC < 50%
d. Commonest cause of anemia in India

- Hyper segmented neutrophils (more than 5 lobes of neutrophils) is seen in vitamin B12 deficiency.
- Iron deficiency anemia is the most common cause of anemia in India.
- It is characterized by microcytic hypochromic peripheral smear.
- MCHC < 50% Low MCHC values occur if you have anemia due to iron deficiency.

4. Severity of mitral stenosis is determined by:

a. Intensity of S1 heart sound
b. Diastolic murmur duration
c. Opening snap
d. Intensity of diastolic murmur

- Since there is mitral stenosis, more time for blood to enter LV.
- Therefore, increased transient time, so increase duration of murmur
- **Intensity of S1 sound can be loud or soft. If sound is soft it tells us about calcified mitral valves but not of severity.

www.medicalcoupons.in

5. Malignant hypertension can lead to all of the following EXCEPT

a. Hypertensive retinopathy
b. Respiratory failure
c. Renal failure
d. Stroke

Emergencies include

- **CNS INVOLVEMENT** hypertensive encephalopathy (headache, irritability, confusion, and altered mental status due to cerebrovascular spasm), intracranial hemorrhage.
- **RENAL INVOLVEMENT** hypertensive nephropathy (hematuria, proteinuria, and acute kidney injury due to arteriolar necrosis and intimal hyperplasia of the interlobular arteries),
- **MALIGNANT HYPERTENSION** Encephalopathy or nephropathy accompanying hypertensive retinopathy has historically been termed malignant hypertension.
- **CARDIOVASCULAR** aortic dissection, preeclampsia-eclampsia, pulmonary edema (acute LVF) unstable angina, or myocardial infarction.

6. Which is the most common arrhythmia:

a. Inappropriate sinus tachycardia
b. Junctional premature complexes
c. Atrial fibrillation
d. Atrial flutter

Atrial fibrillation is the most common sustained arrhythmia, increases with age, and presents with a wide spectrum of symptoms and severity Paroxysmal, persistent, and permanent forms require very individualized approaches to management.

7. Preferred biochemical marker(s) in patients presenting with myocardial infarction:

a. Myoglobin
b. Cardiac specific Troponins
c. CK-MB
d. All of the above

Cardiac troponin I and T is the preferred markers of myocardial injury. It is highly specific to cardiac tissue and accurately diagnoses myocardial infarction with a history of ischemic pain or ECG changes reflecting ischaemia.

www.medicalcoupons.in

8. On ECG, ST segment elevation is seen in all of the following conditions EXCEPT:

a. Left ventricular aneurysm
b. Acute pericarditis
c. Myocardial infarction
d. Hypocalcemia

Cardiac	ST Elevation Myocardial Infarction (STEMI) Pericarditis Ventricular Aneurysm Brugada Syndrome Left Bundle Branch Block
Intracranial	Stroke Raised Intracranial Pressure Intracranial Hemorrhage
Abdominal	Peritonitis
Drugs	Digoxin Isoprenaline Quinidine Procainamide
Metabolic	Hypothermia Hyperventilation Hyperkalaemia
Other	Spinal Cord injury Pulmonary Embolism

9. Most common cause of death in patient on chronic hemodialysis:-

a. Uremia
b. Infection
c. Cardiovascular disease
d. Malnutrition

Cardiovascular disease is the leading cause of death in dialysis patients and sudden death (SD) represents a significant proportion of overall mortality in both hemodialysis (HD) and peritoneal dialysis (PD) patients.

www.medicalcoupons.in

10. True statements regarding Gluten sensitive enteropathy is/ are -

1. Associated with HLA B27

2. Due to auto antibody against brush border epithelium

3. Anti-tissue transglutaminase antibody is Screening test

4. Large intestinal biopsy is Confirmatory test

a. 2, 3 and 4 are true
b. 1, 2 , 3 and 4 are true
c. 2 and 3 are true
d. 1, 2 and 3 are true

Gluten enteropathy/Celiac Sprue

- It is associated with HLA-DQ2
- Due to auto Ab against brush border epithelium which is present in the gut mucosa
- Clinical Feature - Osmotic diarrhea
- Auto ab: Anti tissue transglutaminase Ab - Screening test
- Small intestinal mucosal Biopsy - Confirmatory test
- Treatment:
  → No 'BROW (Barley, Rye, Oats, Wheat)
  → Only maize is allowed.

11. In gastric outlet obstruction in a peptic ulcer patient, the site of obstruction is most likely to be:

a. Antrum
b. Duodenum
c. Pylorus
d. Fundus

- Gastric ulcer leads to gastric outlet obstruction causing pyloric stenosis showing HOUR GLASS STOMACH.

12. Gold standard method of diagnosing celiac disease is

a. Small bowel biopsy
b. Anti-endomysial antibodies
c. Blood picture
d. Biochemical test

www.medicalcoupons.in

- Best screening test: Anti TTG antibody
- Confirmatory / Gold standard: Small Bowel biopsy.

13. In plasmodium vivax malaria, relapse in caused by:

a. Sporozoite
b. Schizont
c. Hypnozoite
d. Gametocyte

Plasmodium vivax malaria is characterized by the presence of dormant liver-stage parasites, **called hypnozoites**, which can cause malaria relapses after an initial attack.

14. A 39 year old male presents with fever, cough and anorexia. He is sputum positive for AFB. In order to consider this as a case of XDR-TB, which of the following should he be resistant to ?

a. Resistance to at least INH & rifampicin + another drugs
b. Resistance to any of the 3 first line drugs
c. Resistance to all 1st line drugs & any 3 second line injectable drugs
d. Resistance to at least INH & rifampicin + any quinolones + at least 1 injectable 2nd line drug


XDR -TB - resistant to :

- INH & rifampicin
- Any fluoroquinolone
- 1 of 3 injectable drug (Amikacin, kanamycin or Capreomycin)
- MDR-TB - Resistance to INH & Rifampicin.

15. Acromegaly is associated with all of the following EXCEPT:

a. Protruding supraorbital ridges
b. Increased heel pad thickness
c. Diabetes mellitus
d. Muscle hypertrophy

www.medicalcoupons.in



16. Vitamin B12 deficiency can lead to all EXCEPT:

a. Myopathy
b. Peripheral neuropathy
c. Optic atrophy
d. Myelopathy

Vitamin B12 deficiency causes :

- Peripheral neuropathy
- Optic atrophy
  Myelopathy But Not Myopathy
- SACD spinal cord Sub-acute combine degeneration of spinal cord).

www.medicalcoupons.in

17. You are an intern, posted in the department of Internal Medicine ward. You are seeing the reports of a 47-year-old female who had history of AGE & having neurological manifestations. Investigations revealed significant elevation of protein & other normal values in CSF. This characteristic dissociation in CSF is a feature of ?

a. Multiple sclerosis
b. GB syndrome
c. TB meningitis
d. Multiple myeloma

- Guillain-Barré syndrome, sometimes known as GBS, is a **rare but serious autoimmune disorder** in which the immune system attacks healthy nerve cells in the peripheral nervous system (PNS). This leads to weakness, numbness, and tingling. It can eventually result in paralysis.
- Patients with GBS have an elevated CSF protein level (**>400 mg/L**), with normal CSF cell counts. Elevated or rising protein levels on serial lumbar punctures and 10 or fewer mononuclear cells/mm^3 strongly support the diagnosis.

18. A 34-year-old woman presented with complaints of sudden onset headache which she described as a characteristic 'thunderclap headache'. She has no significant medical history. On examination, her pupils were dilated bilaterally. These findings are consistent with the diagnosis of?

a. Meningitis
b. Brain stem encephalitis
c. Acute ischemia of midbrain
d. Acute aneurismal hemorrhage

Thunderclap headache (TCH) is a term initially introduced to describe the **apoplectic onset of a headache that begins suddenly**, without warning, and peaks with severe intensity within seconds.

- SAH → Thunder clap headache
- Cause → Rupture of berry's aneurysm
- If post. communicating artery rupture → Involves oculomotor Nerve → So Dilated pupils.

www.medicalcoupons.in

19. A 44 year old man presents to emergency room with Progressive atrophy and weakness of hands. On further examination he is found to have brisk jaw jerk Fasciculations, spasticity of lower leg and Bulbar palsy. In the given clinical scenario Upper motor neuron lesion is characterized by which of the following?

a. Weakness and spasticity
b. Fasciculation's
c. Rigidity
d. Muscle atrophy

- The most likely diagnosis in the given clinical scenario isAmyotrophic lateral sclerosis (ALS).
- Amyotrophic lateral sclerosis (ALS)is the Most common form of motor neuron disease.
- The diagnosis is based on clinical criteria in which there is bothUpper and lower motor neuron signs with progressive disease.
- Spasticity is the velocity-dependent increase in muscle tone due to the exaggeration of stretch reflex. It is only one of the several components of the **upper motor neuron syndrome.**

20. In restrictive lung disease:

a. FVC is high
b. FEV1 is high
c. FEV1/FVC is high
d. All of the above

The FEV1/FVC ratio can be **higher than normal**, This is because it is easy for a person with a restricted lung (e.g. fibrosis) to breathe out quickly, because of the high elastic recoil of the stiff lungs.

21. A 49 year old man, presented with 10 days history of progressive worsening dyspnea on exertion, A chest radiograph confirmed a large left pleural effusion. Which of the following feature and finding you won't see in this case ?

a. Horizontal fluid level
b. Low lung volume
c. Muffled heart sound
d. Decreased chest movements

Horizontal fluid level is characteristic feature of Hydropneumothorax.

www.medicalcoupons.in

Features of Pleural Effusion -

- Blunting of costophrenic angle
- Low lung volumes
- ↓ chest movement
- Dullness on percussion
- Ellis curve
- Muffled Heart sound is characteristic feature of → Pericardial effusion but can be associated with pleural effusion also.

22. Deficiency of which vitamin causes subacute combined degeneration of spinal cord?

a. Vitamin B1
b. Vitamin B6
c. Vitamin B2
d. Vitamin B12

Subacute combined degeneration of the spinal cord is a **neurological complication of vitamin B12 (cobalamin) deficiency**. A deficiency of vitamin B12 can occur as a result of nutritional deficiency, reduced absorption due to altered gastrointestinal anatomy or function, or due to the intake of certain drugs

23. Obstructive lung function is seen in which of the following condition?

a. Obesity
b. Kyphoscoliosis
c. Pleural effusion
d. Asthma

Obstructive lung function is seen in

- COPD
- Asthma
- Bronchiectasis
- α_1-antitrypsin deficiency.


www.medicalcoupons.in

24. In which of the following conditions there is an increase in lung diffusion capacity?

a. Emphysema
b. Idiopathic pulmonary fibrosis
c. Alveolar haemorrhage
d. Pulmonary oedema

DLCO(diffusing capacity of the lungs for **carbon monoxide**)is increased in alveolar hemorrhage whereas it decreases in pulmonary edema, idiopathic pulmonary fibrosis, emphysema.

25. In the given ECG widespread concave ST elevation and PR depression is present throughout the precordial (V2-6) and limb leads (I, II, aVL, aVF).There is reciprocal ST depression and PR elevation in aVR. What is the diagnosis ?



a. Acute myocardial infarction
b. Pericarditis
c. Digoxin toxicity
d. Hypertrophic obstructive cardiomyopathy

- A concave ST segment elevation is the classical pattern of ECG pericarditis.
- A convex ST segment elevation is the classical pattern of ECG MI.

www.medicalcoupons.in

26. A 59-year-old woman presented with a six-week history of left facial weakness, left hearing loss, and left facial numbness, and one-week history of headaches, gait disturbance, and dizziness. She has no significant past medical history. MRI revealed contrast-enhancing dural-based left middle cranial fossa mass with extension into the internal auditory canal and cerebral edema which leads to the suspicion of intracranial metastasis The most common site of intracranial metastasis is from primary carcinoma of ?

a. Breast
b. Lungs
c. Stomach
d. Testes

MC brain tumor metastasis, is arising from

- Oat cell Ca lung/small cell Ca lung
- Ca Breast
- M. Melanoma

MC primary Brain tumor → Glioma

MC benign brain tumor → Meningioma

SURGERY JUNE 2018

1. Which of the following is not seen in massive blood transfusion?

a. DIC
b. Hypothermia
c. Hypercalcemia
d. Thrombocytopenia

- Massive blood transfusion can cause DIC, Hypothermia, Thrombocytopenia
- Because of the use of chelating agents in blood that is transfused hypocalcemia is caused and not Hypercalcemia.

2. In traumatic cases, shock is most likely due to:-

a. Injury to intra-abdominal solid organ
b. Head injury
c. Septicemia
d. Cardiac failure

www.medicalcoupons.in

5 Causes of massive bleeding or shock:

- Injury on the floor (external bleed)
- Injury to Thorax
- Injury to Abdomen
- Injury to Retroperitoneal organs
- Fracture Of long bone

Out of these most common cause is injury to intra-abdominal solid organ.

3. Most common site of peripheral aneurysm?

a. Brachial artery
b. Popliteal artery
c. Radial artery
d. Femoral artery

- MC site of peripheral aneurysm: Popliteal aneurysm
- Most common site of splanchnic aneurysm is splenic artery
- Most common site of visceral arterial aneurysm is splenic artery
- MC vessel involved in aneurysm: Circle of Willis
- MC location of extra - cranial aneurysm: Aorta >Iliac > Popliteal > Femoral (**AIPE**)
- MC site of extra- cranial arterial aneurysm is infrarenal aorta
- Degenerative aneurysms (caused by atherosclerosis) are MC AAA (**abdominal aortic aneurysm**) (90%) .

4. Best way to localize extra-adrenal pheochromocytoma:-

a. X-ray
b. Nucleotide scan
c. VMA excretion
d. Clinical examination

IOC for extra adrenal pheochromocytoma is MRI > MIBG scan(aka-nucleotide scan)

5. Grey Turner's sign in seen in:-

a. Acute appendicitis
b. Acute pancreatitis
c. Acute cholecystitis
d. Acute hepatitis

In acute pancreatitis, bleeding into fascial planes can produce bluish discoloration of the flanks (known as Grey turner's sign) or umbilicus (Cullen sign)

6. All of the following are features of Zollinger-Ellison syndrome except:

a. Intractable peptic ulcers
b. Severe diarrhea
c. Arise from beta cell tumors of pancreas
d. Very high acid output

- ZES(Zollinger-Ellison syndrome) aka Gastrinoma - arises from G- cells/ Gastrin producing cells
- Has increased gastrin production causing increased acid production- leads to peptic ulcer disease
- Causes of diarrhea:
  → Acid produced reaches duodenum and inactivates pancreatic enzymes(works only in alkaline ph) causing Malabsorption
  → Acid reaches proximal jejunum and causes irritation- leads to jejunal secretion.

7. Most common type of gastric sarcoma:-

a. Lipoma
b. Glomus tumor
c. Leiomyosarcoma
d. Leioblastoma

- Leiomyosarcoma is the most common type of gastric sarcoma

8. Most common type of intussusception:-

a. Colocolic
b. Ileoileal
c. Ileocolic
d. Ileal

→ Most common type of intussusception is ileocolic
→ Most common type of intussusception in adults is colocolic.

www.medicalcoupons.in

9. A patient after heavy drinking of alcohol presents with multiple episodes of vomiting and hematemesis. Most likely diagnosis could be:-

a. Mallory-Weiss syndrome
b. Esophageal carcinoma
c. Achalasia cardia
d. Boerhaave's syndrome

- **Mallory- Weiss syndrome** characterized by longitudinal partial tear (mucosa+submucosa) Most common site is cardia and source of bleeding for hematemesis is left gastric artery.
- **Seen in alcoholic males due to forceful and repeated vomiting or Retching**

10. The treatment of choice for inguinal hernia in infants is:-

a. Herniotomy
b. Herniorrhaphy
c. Truss
d. Hernioplasty

- **TOC for congenital hydrocele and inguinal hernia in infant is Herniotomy**
- Mesh is not used in infants because there will be growth in the body of infants while the mesh wont grow.

11. Serum acid phosphate is raised in:-

a. Osteosarcoma
b. Prostatic carcinoma
c. Paget's disease
d. Hyperparathyroidism

Elevated serum levels of acid phosphatase are seen in **patients with carcinoma of the prostate that has extended beyond the prostatic capsule**. Patients with prostatic carcinoma still confined within the capsule usually have a normal serum acid phosphatase level.

www.medicalcoupons.in

12. A 61 year old patient undergoes TURP, after 3 days patient develops altered sensitiveness and drowsiness. Most probable diagnosis is:-

a. Hypernatremia
b. Hyponatremia
c. Stroke
d. Meningitis due to spinal anesthesia

TURP syndrome (Dilutional hyponatremia or water intoxication)

- TURP syndrome (Dilutional hyponatremia or water intoxication) resulting from a hypervolemic, **hyponatremic state due to absorption of the hypotonic irrigating solution.**
- **Clinical Features:** Nausea, Vomiting, confusion, hypertension, bradycardia, & visual disturbances.

13. 54 year old female known case of multinodular goitre now complaints of sudden increase in size of swelling and mild pain over the swelling. There is no history of dysphagia, dyspnea. The patient complaints of mild hoarseness of voice. No lymph nodes palpable. Malignancy in this patient is most often:-

a. Follicular carcinoma
b. Papillary carcinoma
c. Anaplastic carcinoma
d. Medullary carcinoma

Follicular ca: -

- Seen in iodine deficient areas
- M/C malignancy in long standing goiter
- M/C malignancy in multinodular goiter.

14. Paget's disease of the nipple is:-

a. Infection
b. Dermatitis
c. Neoplasia
d. Hypopigmentation

Paget disease (PD) is defined as the presence of **neoplastic cells of glandular differentiation**, interspersed between keratinocytes of nipple epidermis, and constitutes approximately 1% of breast cancer cases.

www.medicalcoupons.in

15. Most common parotid tumor:

a. Metastasis
b. Adenoid cystic carcinoma
c. Warthin's tumor
d. Pleomorphic tumor

- **The most common tumor of the parotid gland is the pleomorphic adenoma, which represents about 60% of all parotid neoplasms.**
- MC neoplasm of salivary gland in children: Hemangioma
- MC neoplasm of salivary gland: Pleomorphic adenoma
- MC malignant tumor of salivary gland: Mucoepidermoid carcinoma
- MC malignant tumor of salivary gland in children: Mucoepidermoid carcinoma
- MC malignant tumor of minor salivary glands: Adenoid cystic carcinoma
- Best diagnostic modality for parotid swelling: FNAC
- Open incisional biopsy is contraindicated due to tumor cell implantation and formation of parotid fistula
- Best imaging investigation for salivary gland neoplasms: MRI

16. Commonest site of branchial cyst:-

a. Lower 1/3 sternomastoid on anterior border
b. Lower 1/3rd sternomastoid on posterior border
c. Upper 1/3rd sternomastoid on anterior border
d. Upper 1/3rd sternomastoid on posterior border

A branchial cleft cyst (BCC) commonly presents as a solitary, painless mass in the neck of a child or young adult. They are most commonly located **along the anterior border and the upper third of the sternocleidomastoid muscle in the anterior triangle of the neck**.

17. Lucid interval is classically seen in:-

a. Intracerebral hematoma
b. Acute subdural hemorrhage
c. Chronic subdural hemorrhage
d. Epidural hemorrhage

THE LUCID interval following head trauma and unconsciousness is described classically in **epidural hematomas**. The historic emphasis placed on the lucid interval in cases of extradural hematoma has made this one of the best-remembered signs of the syndrome.

www.medicalcoupons.in

18. Marjolin's ulcer is:-

a. Squamous cell carcinoma from scar
b. Adenoma of scar
c. Tubercular ulcer
d. Amoebic ulcer

A Marjolin ulcer is a **rare and aggressive type of skin cancer** that grows from burns, scars, or poorly healing wounds. It grows slowly, but over time it can spread to other parts of your body, including your brain, liver, lungs, or kidneys. **Most Marjolin ulcers are cancerous and form squamous cancer cells** in the upper layers of your skin. Some Marjolin ulcers may also form as basal cell tumors which form in deeper layers of your skin.

19. In treatment of hand injuries, the greatest priority is:-

a. Repair to tendons
b. Repair of skin cover
c. Repair of nerves
d. All

Sequence of Repair in Hand Injuries (LE FAN OF VEINS)

→ **B**one shortening of stabilization/ fixation
→ **E**xtensor tendon repair
→ **F**lexor tendon repair
→ **A**rterial anastomoses
→ **N**erve repair
→ **V**enous anastomosis
→ **S**kin/ wound closure

20. A boil is due to staphylococcal infection of:-

a. Hair follicle
b. Sweat gland
c. Subcutaneous tissue
d. Epidermis

The most common type of staph infection is the boil, a pocket of pus that develops in a hair follicle or oil gland. The skin over the infected area usually becomes red and swollen.

www.medicalcoupons.in

21. Pendred syndrome is due to a defect in:-

a. Chromosome 7p
b. Chromosome 7q
c. Chromosome 8p
d. Chromosome 8q

Pendred syndrome can be caused by changes, or mutations, **in a gene called SLC26A4 (formerly known as the PDS gene) on chromosome 7**. Because it is a recessive trait, a child needs to inherit two mutated SLC26A4 genes—one from each parent—to have Pendred syndrome.

22. Cock's peculiar tumor is:-

a. Papilloma
b. Infected sebaceous cyst of scalp
c. Cylindroma
d. Squamous cell carcinoma

Cocks Peculiar Tumour

- Infected or ulcerated sebaceous cyst of the scalp
- Resembles fungating epithelioma


23. Mondor's disease is:-

a. Lymphangitis of mammary lymphatic's
b. Multiple breast cysts
c. Eczema by nipple and areola
d. Thrombophlebitis of superficial veins of breast

Mondor's disease is a **condition characterized by thrombophlebitis of the superficial veins of the breast and anterior chest wall**. Initially this condition presents with rapid development of a subcutaneous red cord-like lesion that causes pain at an early stage and subsequently becomes a painless fibrous band.

www.medicalcoupons.in

24. What is the procedure shown in the image?



a. Mile's procedure
b. Frey's procedure
c. Scanlon's modification surgery
d. Anderson-Hynes operation

The Anderson-Hynes pyeloplasty (AHP) is **a surgical technique used in the management of pelvi-ureteric junction obstruction**. This operation has been performed for over six decades and has etched its name in the annals of urology.

25. Oliguria is defined as:-

a. Absence of urine production
b. More than 900 ml of urine excreted in a day
c. 600 ml to 700 ml of urine excreted in a day
d. Less than 300 ml of urine excreted in a day

Oliguria is defined as **urinary output less than 400 ml per day or less than 20 ml per hour** and is one of the earliest signs of impaired renal function.

www.medicalcoupons.in

FMGE DEC 2018
ANATOMY

1. A 55-year-old male is brought to the Neurology OPD by his wife. She complains that something is wrong with her husband and they were referred to this center by a local doctor. The doctor's note read:- Patient unable to repeat words spoken, though is able to understand what is told & nods head accordingly. When asked to name people and write, the patient found difficulty. Referred higher center for CT brain. You being the intern on duty, would inform the CMO it could be a lesion in which area of the brain?

a) Arcuate fasciculus
b) Anterior commissure
c) Corpus callosum
d) Fornix

- Arcuate fasciculus connects Wernicke's area present in the posterior superior temporal gyrus of the dominant hemisphere (sensory area of speech) to Broca's speech area present in the prefrontal & premotor facial region of the cerebral cortex to integrate language understanding & skilled motor function.
- Damage to Wernicke's area = Wernicke's aphasia (lack meaning)
- Damage to Broca's area = motor aphasia (vocal system couldn't emit words)
- Damage to arcuate fasciculus = conduction aphasia

2. Mammillary body is a structure on the undersurface of the brain, developing from the diencephalon, that forms a part of the limbic system. They receive afferent fibers via the:-

a) Corpus callosum
b) Thalamus
c) Pituitary gland
d) Fornix

- Mammillary body receives afferent fibers via Fornix
- C-shaped bundle of nerve fibers which acts as major output tract of the hippocampus
- They are association fibers(connects one brain center to another)
- It is a part of limbic system (Papez circuit)
- Papez circuit for memory consolidation, emotions & behavior

3. The parotid gland is a paired salivary gland contributing serous secretions. Its duct opens at the level of the:.

a) First mandibular molar
b) Second mandibular molar
c) First maxillary pre molar
d) **Second maxillary molar**

www.medicalcoupons.in

Parotid duct/Stenson's duct passes through the buccinator muscle and pierces buccal fa , opening into the vestibule of the mouth at the parotid papilla, which lies opposite to the 2nd maxillary (upper) molar tooth.

4. Trismus is due to spasm of which of the following muscle (FMGE Dec 2018)

a) Buccinator
b) Lateral pterygoid
c) Medial pterygoid
d) Mentalis

- **Trismus → spasm of muscles of mastication esp ally medial pterygoid muscle. [classical example of tetanus]**
- Lateral pterygoid + depressor of mandible pull mandible antero-inferiorly and open the mouth
  → Antagonize (MTM muscles)
- Tetanus neonatorum → spasm of masseter
- **MTM- elevators of mandible**

→ M - Medial pterygoid → inserted to ramus of mandible from inner side
→ T Temporalis → inserted into coronoid process of mandible
→ M -Masseter → inserted to ramus of mandible from outside

5. Which branch of facial nerve supplies muscles of lower lip (FMGE Dec 2018)

a) Temporal
b) Cervical
c) Buccal
d) Mandibular

The marginal mandibular branch of the facial nerve passes forward beneath the platysma and depressor anguli oris, **supplying the muscles of the lower lip** and chin, and communicating with the mental branch of the inferior alveolar nerve.

→ **TEMPORAL BRANCH-**Supply upper part of orbicularis oculi and frontalis
→ **ZYGOMATIC BRANCH-**Supply lower part of orbicularis oculi [helps to shut eyelid tightly]
→ **BUCCAL BRANCH**-Supply upper orbicularis oris, zygomaticus minor/major

( ) Mediastinal surface of the right lung is related to:-

a) The Azygos vein
b) The Right Lymphatic Duct
c) The Aorta
d) The Trachea

Azygos vein > > Trachea (because heart & its blood vessels should be preferred first as an answer as they cause direct impression on lungs).

The Mediastinal surface of

Right lung contains	Left lung contains
• Heart (right atrium) • Superior vena cava • Inferior vena cava • Azygos vein • Esophagus • Right subclavian artery & vein arch over and are related to the superior lobe of the right lung.	• Heart (left ventricle) • Aortic arch • Thoracic aorta • Esophagus • Left subclavian artery & vein arch over related to the superior lobe of the left lung.

7. Sternum is attached to scapula via (FMGE Dec 2018)

a) Manubrium
b) Clavicle
c) First rib
d) Second rib

- **By forming stenoclavicular joint & Acromioclavcular joint clavicle connects sternum (Axial skeleton) to scapula (Appendicular skeleton).**
- Clavicle transmits the weight / force of appendicular skeleton to axial skeleton. acromioclavicular ligament is important in this.
- That's why clavicle is most commonly fractured bone in body [mc = middle third]

8. Retraction of scapula at sternoclavicular joint is done by (FMGE Dec 2018)

a) Serratus anterior
b) Trapezius
c) Subscapularis
d) Supraspinatus

Retractors of scapula:

- Trapezius
- Rhomboid major
- Rhomboid minor

www.medicalcoupons.in

Protraction of scapula:

- Pectoralis minor
- Serratus anterior (damage causes winging of scapula)

9. Winging of scapula is due to paralysis of (FMGE Dec 2018)

a) Trapezius
b) Serratus anterior
c) Latissimus dorsi
d) Pectroralis major

The most common cause of scapular winging is **paralysis of the serratus anterior muscle due to the injury of the long thoracic nerve**.



10. During the embryological development, the midgut rotates a total of _____in the intrauterine life.

a) 270 degree clockwise
b) 270 degree anticlockwise
c) 360 degree clock wise
d) 360 degree anticlockwise

www.medicalcoupons.in

During development, midgut undergoes rapid phase of growth in which loops of midgut herniate outside the abdominal cavity of fetus & protrude into umbilical cord

This is PHYSIOLOGICAL UMBILICAL HERNIA (occurs at 6 weeks of IUL)

At week 10, it comes back to abdominal cavity.

When midgut s Herniating into umbilicus, it follows superior mesenteric artery Thus, superior mesenteric artery becomes axis of rotation.

While entering umbilical cord there is:

- **High point: duodenojejunal junction (small intestine).**
- **Low point: ileocecal junction (large intestine)**

→ Which rotates 1st 90 degree anti- clock wise resulting in small intestine to right side & large intestine to left side.

→ While coming back to abdominal cavity, there would be another 180 degree anti - clock wise rotation.

→ **Total of 270 degree anti-clock wise rotation occurs during development of midgut in IUL**

→ Duodenum become C shaped and have 4 parts and colon will also have 4 parts (Ascending, transverse & descending & sigmoid colon).

11. Vasa brevia is the other name for which of the following arteries carried by the gastrosplenic ligament?

a) Long gastric arteries
b) Short gastric arteries
c) Iliea vasa recta
d) Jejunal vasa recta

- Splenic artery (branch of celiac trunk) supply greater curvature of stomach by it following branches:
  → **Short gastric arteries which are called as vasa brevia. (carried by Splenogastric ligament)**
  → Long gastric artery called left gastroepiploic artery.
- Jejunal vasa recta (long arteries) & ileal vasa recta (short) are straight arteries coming from arcades in mesentery of jejunum & ileum.

12. **In a healthy adult female, the tail of pancreas lies in lienorenal ligament derived from peritoneum. This is another name for which of the following ligaments?**

a) Hapato-gastric
b) Hepato- duodenal
c) Gastro- splenic
d) Spleno- renal

www.medicalcoupons.in

- **Splenorenal ligament**, also known as the lienorenal ligament is a peritoneal ligament. It represents the dorsal most part of dorsal mesentery and forms part of the lateral border of the lesser sac. It is continuous with the gastrosplenic and phrenicocolic ligaments
- Tail of pancreas lies in Spleno renal ligament.
- Gastro splenic short gastro and left gastro-omental vessels.

13. Bleeding vessels in hemorrhoids is (FMGE Dec 2018)

a) Superior rectal artery
b) Superior rectal vein
c) Middle rectal artery
d) Middle rectal vein

External hemorrhoids:

- Hemorrhoids occurring at anal verge (distal boundary of anal canal below pectinate line) Normally have low tendency to bleed on straining at stool.
- **Bleeding comes from inferior rectal vein.**

Internal hemorrhoids:

- Occur inside the rectum (above pectinate line) have high tendency to bleed.
- **Bleeding from "superior rectal vein"**
- Prolapsed hemorrhoids are internal hemorrhoids that pass outside anal canal & form lump, which may undergo thrombosis & become painful.

14. Foot eversion is caused by (FMGE Dec 2018)

a) Extensor digitorum
b) Peroneus longus
c) Tibialis anterior
d) Tibialis posterior

- In human anatomy, the peroneus longus (also known as fibularis longus) is **a superficial muscle in the lateral compartment of the leg**, and acts to evert and plantarflex the ankle.
- **Eversion of the Foot (tilting of the sole of the foot away from the midline): Performed by the fibularis brevis and fibularis longus.**
- Inversion of the Foot (tilting of the sole of the foot inwards towards the midline): Performed by the tibialis posterior and tibialis anterior.

www.medicalcoupons.in

15. Tibial and common peroneal nerve supplies the muscle (FMGE Dec 2018)

a) Gracilis
b) Biceps femoris
c) Adductor longus
d) Adductor magnus

Biceps femoris is a muscle of the posterior compartment of the thigh and lies in the posterolateral aspect. It arises proximally by two 'heads', termed the 'long head' (superficial) and the 'short head' (deep).

Nerve Supply

- Long head: tibial division of the sciatic nerve
- Short head: common peroneal division of the sciatic nerve

PHYSIO DEC 2018

1. The ICF volume represents the fluid contained within the body's cells. This volume cannot be measured directly but is calculated as the difference between the measured TBW and the measured ECF volume. Potassium provides the osmotic skeleton for the ICF and because water is freely diffusible into and out of the cell, changes in the tonicity of the ECF are rapidly reflected by similar changes in ICF tonicity. Which of the following has the least concentration in intracellular fluid?

a) Calcium
b) Magnesium
c) Potassium
d) Protein

- Calcium is kept very LOW in cytoplasm, hence intracellular fluid doesn't contain much Ca2+
- Calcium is mostly stored in organelles inside cell

E.g. In sarcoplasmic reticulum of muscle cell etc.

2 Resistance against passive stretch is called?

a) Tone
b) Spasticity
c) Rigidity
d) Paratonia

Muscle tone is defined as the continuous and passive-partial contraction of the muscle or the muscle's resistance to passive stretch during the resting state.

Spasticity:

Condition in which certain muscles are continuously contracted

This contraction causes stiffness/tightness of muscle & interfere with movement

Occurs in lesions of upper motor neurons.

Rigidity:

increase in muscle tone leading to resistance to passive movement throughout range of motion

E.g. Cogwheel rigidity in parkinsonism

Paratonia:

Inability to relax muscles during muscle tone assessment.

3. The formation of the active enzyme peptidases from their inactive precursors occurs only when they have reached their site of action, secondary to the action of the brush border hydrolase enterokinase. The powerful protein-splitting enzymes of the pancreatic juice are secreted as inactive proenzymes. Enterokinase is an activator of?

a) Trypsinogen
b) Trypsin
c) Chymotrypsin
d) Antitrypsin

- When first synthesized in pancreatic cells, the proteolytic digestive enzymes like trypsinogen, chymotrypinogen & pro-carboxypeptia are in inactive forms(to prevent auto proteolysis of pancreas)
- After secreted into intestine, they are activated:
  → Trypsinogen is activated by enterokinase (secreted by intestinal mucosa)
  → Trypsinogen → can also be autocatalytically activated by trypsin
  → Chymotrypsinogen is activated by trypsin to form chymotrypsin.

www.medicalcoupons.in

4. Which structure connects the Broca's area and Wernicke area?

a) Arcuate fasciculus
b) Anterior commissure
c) Corpus callosum
d) Fornix

- **Arcuate fasciculus connects Wernicke's area present in the posterior superior temporal gyrus of the dominant hemisphere (sensory area of speech) to Broca's speech area** present in the prefrontal & premotor facial region of the cerebral cortex to integrate language understanding & skilled motor function.
- Damage to Wernicke's area = Wernicke's aphasia (lack meaning)
- Damage to Broca's area = motor aphasia (vocal system can't emit words)
- Damage to arcuate fasciculus = conduction aphasia

5. Calcium binding proteins are proteins that participate in calcium cell signaling pathways by binding to Ca2+, which plays an important role in many cellular processes. Calcium-binding proteins have specific domains that bind to Ca2+ and are known to be heterogeneous. One of the functions of calcium-binding proteins is to regulate the amount of free (unbound) Ca2+ in the cytosol of the cell. The cellular regulation of calcium is known as Calcium homeostasis. Which of the following is not a calcium-binding protein?

a) Calbindin
b) Calmodulin
c) Troponin
d) Clathrin

Clathrin (does not bind Ca2+)

- On inside of cell membrane beneath coated pits, latticework of fibrillar protein is present called clathrin
- Helps in pinocytosis

Calbindin

- Present in brush border of intestinal epithelial cells & transports Cat2into cell cytoplasm
- Amount of calcium absorption is directly proportional to quantity of calcium-binding protein

Troponin

- Troponin C - binds to Ca+2 ions = Strong affinity of troponin to calcium initiate contraction process.

www.medicalcoupons.in

Calmodulin

- In place of troponin, smooth muscles have calmodulin
- Ca2+ inside the cell binds reversibly with calmodulin & activates myosin light chain kinase resulting in a chain of events that cause muscle contraction

6. Histamine is secreted by:

a) Enterochromaffin cell
b) Parietal cell
c) Oxyntic cell
d) Chief cell

"Enterochromaffin-like (ECL) cells" are a population of cells that are found in the **gastric pits of the stomach luminal epithelium** and secrete histamine. In response to gastrin released by neighbouring G-cells, secreted histamine from ECL cells acts on parietal cells to stimulate the release of gastric acid.

7. Reward centre in the brain is?

a) Amygdala
b) Thalamus
c) Hippocampus
d) Ventral tegmental area

- Reward center contains ventral tegmental area connected with nucleus accumbens (through dopamine NT)
- Previously major reward centers are thought to be located along course of "medial forebrain bundle"

8. Glucose is absorbed by?

a) Upper part of PCT
b) Ascending limb of LOH
c) Cortical Collecting duct
d) Medullary collecting duct

Substances like glucose, AA, Na+2, H,0 is absorbed maximum from upper part of PCT

- As proximal tubule epithelial cells have:
  → Large number of mitochondria to support powerful active transport processes (highly metabolic)
  → Have brush borders & provide extensive surface area for absorption

www.medicalcoupons.in

→ **As glucose is reabsorbed by secondary active transport, it is absorbed almost all by proximal convoluted tubule.**

9. Sperm remains in epididymis for_____hours?

a) 24
b) 48
c) 72
d) 96

- Sperms after forming in seminiferous tubules enter epididymis
- Sperms in seminiferous tubules & early portions of epididymis → non-motile & cannot fertilize ovum
- After sperm had been for (18-24 hrs) in epididymis sperms attain capability of motility.

10. What mediates Inverse stretch reflex

a) Golgi tendon
b) Muscle spindle
c) Unmyelinated C fibres
d) Dorsal Column

The Inverse Myotatic Reflex involves **Sensors of Muscle Force in the Tendon**. **Stretch receptors called Golgi tendon organs** are found within the collagen fibers of tendons and within joint capsules. They are generally located in series with the muscle rather than the parallel arrangement of the extrafusal muscle fibers.

11. Organelle having DNA is_____?

a) Mitochondria
b) Golgi complex
c) SER
d) RER

- **The three organelles that contain DNA are the nucleus, mitochondria and chloroplasts.**
- Organelles are membrane-bound subunits within a cell -- analogous to organs in the body -- that perform specific functions.
- The nucleus is the control center of the cell, and houses genetic information.
- The mitochondria and chloroplasts both produce energy, in animal and plant cells, respectively.

www.medicalcoupons.in

12. All are Glucogenic hormones except?

a) ADH
b) Glucagon
c) Thyroxine
d) Glucocorticoids

Anti - diuretic hormone (ADH) / vasopressin :

- It is secreted from posterior pituitary
- Controls rate of water excretion into urine thus controlling concentration of water in body fluids
- No effect on glucose

Glucogenic hormones are those which are capable of increasing blood glucose levels

1.Glucagon

2. Thyroxine

3. Glucocorticoids

4. Catecholamines,

5. Growth hormone etc.

13. Delivery of stimulus above threshold intensity leads to a constant amplitude of AP and is known as:

a) All or none law
b) Electronic potential
c) Absolute refractory period
d) Relative refractory period

All or none principle -

The all-or-none law is a principle that states that **the strength of a response of a nerve cell or muscle fiber is not dependent upon the strength of the stimulus**. If a stimulus is above a certain threshold, a nerve or muscle fiber will fire.

www.medicalcoupons.in

14. Which part of thalamus is related to motor control?

a) Ventrolateral thalamus
b) Ventral posteromedial
c) Ventral posterolateral
d) Superior and inferior colliculi

Thalamus is sensory relay station for all sensations except olfaction

It is related to motor control also by :

→ Ventrolateral thalamus
→ Ventroanterior thalamus
→ Centro median thalamus

15. Active form of Vitamin D is?

a) Calcitriol
b) Calciferol
c) Calcidiol
d) Ergocalciferol


Calcitriol is the active form of vitamin D, normally made in the kidney. It is also known as **1,25-dihydroxycholecalciferol**. It is a hormone which binds to and activates the vitamin D receptor in the nucleus of the cell, which then increases the expression of many genes. Calcitriol increases blood calcium (Ca^{2+}) mainly by increasing the uptake of calcium from the intestines.

16. Tyrosine kinase receptors are membrane-spanning proteins with large amino-terminal extracellular domains bearing the ligand binding site, a juxtamembrane domain, a protein kinase catalytic domain, and carboxyl-terminal tail. Which of the hormone has 4 subunits and 2 units for tyrosine kinase receptor binding?

a) Insulin
b) Glucagon
c) TSH
d) ADH

www.medicalcoupons.in

The insulin receptor is an enzyme-linked receptor.



Insulin binds to α-subunits → due to linkage, portions of β-subunits become auto-phosphorylated → activates [Tyrosine kinase & IRS]

17. In case a small stimulus causes more pain, it is called:

a) Hypersensitivity
b) Causalgia
c) Hyperalgesia
d) Allodynia

yperalgesia:

An increased sensitivity to feeling pain and an extreme response to pain. Hyperalgesia may occur when there is damage to the nerves or chemical changes to the nerve pathways involved in sensing pain.

- **Hypersensitivity** (also called hypersensitivity reaction or intolerance) refers **to undesirable reactions produced by the normal immune system**, including allergies and autoimmunity.
- **Allodynia** is **a type of neuropathic pain (nerve pain)**. People with allodynia are extremely sensitive to touch. Things that don't usually cause pain can be very painful. These may include cold temperatures, brushing hair or wearing a cotton t-shirt.
- **Causalgia** is burning of type pain caused crushing, by injuries to nerves. (neuropathic pain)

www.medicalcoupons.in

BIO-CHEM DEC 2018

1. Urea cycle occurs in:

a) Mitochondria
b) Cytoplasm
c) Both
d) None

In both (first in mitochondria then in cytoplasm)- **HUG**

- **H**eme synthesis
- **U**rea cycle
- **G**luconeogenesis

2. Start codon is:

a) UAA
b) UAG
c) UGA
d) AUG

- Start codon - AUG - synthesize methionine in eukaryote and formyl methionine in prokaryote
- Stop codon -UAA ,UAG, UGA

3. Melanin is synthesized from :

a) Tyrosine
b) Tryptophan
c) Phenyl alanine
d) Threonine

Melanin is a highly irregular heteropolymer consisting of monomeric units derived from the enzymatic oxidation of the **amino acid tyrosine**. The process of melanin formation takes place in specialized acidic organelles (melanosomes) in melanocytes.

4. Which of the following helps in wound healing?

a) Keratan sulfate
b) Dermatan sulfate
c) Hyaluronic acid
d) Chondroitin sulfate

www.medicalcoupons.in

Hyaluronic acid (hyaluronan) is a key component of the extracellular matrix and is known to be involved in several mechanisms of the wound healing process. It has been known to **improve and accelerate the healing process of chronic wounds**.

5. Oxidative phosphorylation is inhibited by all EXCEPT:

a) CO
b) Antimycin A
c) Malonate
d) Thermogenin

Inhibitor of ETC - inhibit oxidation phosphorylation

- Complex 1- Rotenone, phenobarbitone, amobarbitone, piericidine A
- Complex 2- Malonate, TIFA (Thionyl tri fluoroacetate)
- Complex 3 - Antimycin A
- Complex 4 - CO, CN, H2S, sodium azide

Natural uncouplers:

- Thermogenin - responsible for non- shivering thermogenesis
- Thyroxin

Synthetic uncouplers - 2,4 DNP

6. Use of zinc in diarrhea is because it:

a) Reduces the risk, duration and severity of diarrheal episodes
b) Enhances immune response
c) Regulates intestinal transport & absorption of water and electrolytes
d) All

Zinc is required for:

→ Protein synthesis
→ Immune response
→ Cell growth & differentiation
→ Regulation intestinal transport & absorption of water and electrolytes
→ Improve intestinal epithelium lining and increase enzyme in brush border epithelium
→ Cell membrane transport and function
→ For polyribosome
→ More than 100 metalloenzyme require zinc example carbonic anhydrase alkaline phosphates , lactate dehydrogenase
→ Deficiency lead to acrodermatitis enteropathica - skin affected around cheek elbow knee anus and mouth

→ In diarrhea zinc is given along with ORS to reduce the risk , duration and severity.

www.medicalcoupons.in

7. Which is cardio protective?

a) HDL
b) LDL
c) IDL
d) VLDL

High-Density Lipoprotein :

- HDL is synthesized and secreted from both liver and intestine
- HDL transport cholesterol from peripheral tissue to liver this is know as reverse cholesterol transport that why HDL is cardio protective.
- HDL concentrations are inversely related to the incidence of atherosclerosis
- HDL have APO A, CAND E
- HDL have LCAT (Lecithin cholesterol acyl transferase) which convert cholesterol into cholesterol ester and then it's taken up by HDL
- LCAT is activate by APOA1
- HDL IS smallest in size max density , max phospholipid (lecithin),min triglyceride.

8. Adenosine receptor stability is because of extensive disulfide bonds formed between:

a) Cysteine
b) Methionine
c) Arginine
d) Alanine

- Cysteine is unique among coded amino acids **because it contains a reactive sulph-hydryl group**. Therefore, two cysteine residues may form a cystine (disulfide link) between various parts of the same protein or between two separate polypeptide chains.
- No other amino acid given as options can form disulfide bond due to absence of -SH group

9. Steps of PCR in sequence are?
a) Extend DNA, anneal primers, Denture DNA
b) Anneal primers, extend DNA, Denature DNA
c) Denature DNA, anneal primers, Extends DNA
d) Denature DNA, Extend DNA, Anneal primers

PCR is based on three simple steps required for any DNA synthesis reaction:

(1) **denaturation of the template into single strands**;

(2) annealing of primers to each original strand for new strand synthesis; and

(3) extension of the new DNA strands from the primers.

www.medicalcoupons.in

10. Genetic material is transferred from one bacterial to another by all EXCEPT

a) Conjugation
b) Transduction
c) Transformation
d) Transfection

Transfection - DNA transfer into eukaryotic cell

- **Conjugation** -a natural microbial recombination process. During conjugation a donor and a recipient come together, join by cytoplasmic bridges and transfer single stranded DNA. Inside the recipient the DNA will combine with chromosome which is rare or it will be remain free in case of plasmid
- **Transduction** - transfer of DNA from one bacteria to another bacteria via virus like bacteriophage
- **Transformation** - transfer of DNA from dead bacteria or environment to another live bacteria
- **Lipofection** - liposome mediated gene transfer. liposome is lipid molecule that can carry nucleic acids

11. Glucose is reabsorbed in which part?

a) Early PCT
b) Henle loop
c) Collecting duct
d) Distal convoluted tubule

Part	Reabsorption	Secretion
PCT	• Glucose 100% • Amino acid 100% • Na along with glucose by SGLT 2 • Chloride (manly in late part of PCT) • Bicarbonate	• Bile salts • Hydrogen ion • Creatinine • Uric acid • Drugs
LOOP OF HENLE	• Descending limb - water • Ascending limb - Na k, Cl by 1NA 2Cl 1k • Cotransporter - Slight reabsorption of cation such as Mg Ca also	• Hydrogen ion by Na H PUMP

www.medicalcoupons.in

DISTAL TUBULE	• 5% sodium and chloride by sodium chloride cotransport	
LATE DISTAL TUBE & COLLECTING DUCT	• Principle cell - reabsorb sodium and water. • Intercalated cell - reabsorb potassium ions	• Principle cell - secrete potassium ions Intercalated cell- hydrogen ions

12. Low glycemic index food is:

a) Easily digestible
b) Increases plasma glucose
c) Has slower absorption
d) Increase glycogen deposits

GLYCEMIC INDEX -glycemic index which measures the time course of postprandial glucose concentrations from a graph

→ **Glycemic index of complex carbohydrate is low because of slow digestion and slow absorption**
→ Glycemic index of carbohydrate is low when its combine with fat , protein and fiber
→ Foods that have a low glycemic index are considered to be more beneficial since they cause less fluctuation in insulin secretion.

PHARMA DEC 2018

1. Which of the following drugs is considered as first line therapy for the prophylaxis of motion sickness in adults?

a) Promethazine
b) Ondansetron
c) Metoclopramide
d) Domperidone

As hyoscine is not given in the options, the answer is first generation Antihistaminics

Best drug/ DOC for prophylaxis of motion sickness - Hyoscine (scopolamine).

First generation Antihistamines can cross the blood brain barrier and also have strong Anti-cholinergic properties.

Example of a **first generation Antihistamines – Promethazine.**

www.medicalcoupons.in

2. Exenatide is a new drug used in diabetes mellitus. Mechanism of action of this drug is:

a) Inhibition of DPP - 4
b) Release of insulin via acting as agonist of GLP-1 receptors
c) Inhibiting intestinal absorption of carbohydrates
d) Stimulation of PPAR-gamma

Mechanism of Action:

Exenatide is a GLP-1 receptor agonist released from the gut and **acts to increase glucose-dependent insulin secretion from pancreatic beta cells**, suppress glucagon secretion, delay gastric emptying, and reduce food intake. The binding of the drug to pancreatic GLP-1 receptors mediates these actions.

3. The compound prostaglandin has been indicated in multiple conditions is actually a derivative of which of the following

a) Stearic acid
b) Arachidonic acid
c) Linoleic acid
d) Linolenic acid

The prostaglandins (PG) are a group of physiologically active lipid compounds called eicosanoids having diverse hormone-like effects in animals. Prostaglandins have been found in almost every tissue in humans and other animals. **They are derived enzymatically from the fatty acid arachidonic acid.**

4. Which of the following drug is contraindicated in a pregnant female?

a) Erythromycin
b) Amoxicillin
c) Ceftriaxone
d) Streptomycin

Some antibiotics are known to be teratogenic and should be avoided entirely during pregnancy. These include **streptomycin and kanamycin** (which may cause hearing loss) and tetracycline (which can lead to weakening, hypoplasia, and discoloration of long bones and teeth).

www.medicalcoupons.in

5. Which of the following is the mechanism of action of methyl dopa?

a) Blockade of beta adrenergic receptors
b) Agonism of beta 2 receptors
c) Agonism of alpha 2 receptors
d) Blockade of alpha 2 receptors

Mechanism of Action

Alpha-methyldopa is **converted to methyl norepinephrine centrally to decrease the adrenergic outflow by alpha-2 agonistic action** from the central nervous system, leading to reduced total peripheral resistance and decreased systemic blood pressure.

6. A 12 months old baby is brought to your clinic. If you have to give a cyclopegic agent for this pediatric patient, which of the following drugs would you prescribe?

a) Atropine oral
b) Atropine eye ointment
c) Tropicamide eye drops
d) Phenylephrine eye drops

In children the tone of ciliary muscle is very high, so we have to give drug which is a strong cycloplegic drug-Atropine.

Atropine is given in the form of eye ointment because it would then have only localised effect in the eyes. If we give it in form of eye drops, it can be absorbed through nasolacrimal duct, and there will be risk of hyperthermia.

Cycloplegic DOC in 1yr old child - Atropine (longest acting cycloplegic)

Cycloplegic DOC in Adults- Tropicamide (shortest acting cycloplegic)

7. Pilocarpine produces:-

a) Active miosis
b) Active mydriasis
c) Passive miosis
d) Passive mydriasis

www.medicalcoupons.in

Pilocarpine is a drug that acts as a **muscarinic receptor agonist**. It acts on a subtype of muscarinic receptor (M_3) found on the iris sphincter muscle, causing the muscle to contract - resulting in pupil constriction (miosis). Pilocarpine also acts on the ciliary muscle and causes it to contract.

8. Which of the following drugs is a direct inhibitor of clotting factor Xa?

a) Apixaban
b) Argatroban
c) Fondaparinux
d) Aspirin

The factor Xa inhibitors that are currently commercially available include **rivaroxaban, apixaban, betrixaban, and edoxaban**. These drugs bind to factor Xa and prevent the formation of thrombin by interrupting the extrinsic and intrinsic coagulation cascades.

9. Which of the following drug is a protein synthesis inhibitor?

a) Linezolide
b) Nevirapine
c) Indinavir
d) Ceftriaxone

Following category of drugs are protein synthesis inhibitors (in Bacteria):-

- Tetracyclines - doxycycline, minocycline
- Chloramphenicol
- Macrolides - Erythromycin, Azithromycin, etc.
- Linezolid
- Clindamycin
- Aminoglycoside - streptomycin, Gentamicin, Kanamycin, etc.
- Ceftriaxone → Cephalosporin → inhibits cell wall synthesis
- Nevirapine → NNRTI → inhibits reverse transcriptase enzyme
- Indinavir → Protease inhibitor used in highly Active anti- retroviral therapy to treat HIV/AIDS.

10. Drug of choice for prophylaxis of meningococcal meningitis in a pregnant female after contact with a case is:-

a) Ciprofloxacin
b) Ceftriaxone
c) Cefalexin
d) Rifabutin

www.medicalcoupons.in

Important points to remember about Meningococcal Meningitis

→ Doc for Empirical Treatment → ceftriaxone
→ Doc for Definitive Treatment → Penicillin G
→ Doc for Prophylaxis → ciprofloxacin
→ Doc for Prophylaxis in pregnancy → ceftriaxone

11. Dose of oseltamivir for chemoprophylaxis in an infant is:-

a) 3 mg / kg once daily
b) 20 mg once a day
c) 20 mg twice day
d) 20 mg thrice day

	Age	Weight	Dose	Frequency	Duration
Oral oseltamivir for treatment of influenza	<1yr		3mg/kg	Twice daily	5 days
	>1yr	<15kg	30mg		
		15-23kg	45mg		
		23-40kg	60mg		
		>40kg	75mg		
Oral oseltamivir for Chemoprophylaxis	3month-1yr		3mg/kg	Once daily	7 days
		<15kg	30mg		
		15-23kg	45mg		
		23-40kg	60mg		
		>40kg	75mg		

12. Purpose of adding zinc to insulin in commercial preparations for treatment of diabetes mellitus is:-

a) To make it long acting
b) To make it short acting
c) To make is fast acting
d) Addition of zinc make the insulin effective orally

Insulin exists in two forms:-

1. Hexamer

2. Monomer

www.medicalcoupons.in

→ Monomer is the active form of Insulin, so it will start producing action - > decreasing blood sugar immediately and in no time the action will be finished.
→ We want Insulin to stay as Hexamer and gradually it is released into the monomer forms so that duration of action can be increased.
→ This is achieved by **adding zinc to the Insulin** (on adding zinc, the insulin starts getting crystallized and those crystals are deposited and act as depot preparation and slowly-slowly from that site, monomers are released and insulin keep on acting) Hence, **zinc is added to make insulin long acting.**

13. Which of the following is the shortest acting muscle relaxant and is considered for rapid sequence induction?

a) Tubocurarine
b) Succinylcholine
c) Pancuronium
d) Atracurium

Muscle Relaxants are of two types:-

MR

- Depolarizing

- Non Depolarizing

- Shortest Acting Depolarizing muscle relaxant → succinyl choline.
- Overall shortest acting muscle relaxant →succinyl choline.
- Action of succinyl choline **last for < 5 mins.**
- Shortest Acting Non Depolarizing Muscle relaxant → Mivacurium.

14. The drug that causes fall in elderly patients with postural hypotension is:

a) Metformin
b) Prazosin
c) Acarbose
d) Nor-adrenaline

The side effect postural hypotension AkA "First Dose Hypotension" is observed with drug ending with zosin - Prazosin, Terazosin, Doxazosin.
These drugs cause blocking of a1 receptors, leading to vasodilation, that can Cause Postural Hypotension.

www.medicalcoupons.in

15. A 23 year old female presented with burning pain during micturition. On investigations, she is found to be suffering from co-infection with chlamydia and gonococcus. Drug of choice for treatment of this female is:.

a) Azithromycin
b) Ceftriaxone
c) Fluconazole
d) Clindamycin

Doc for gonococcal urethritis → ceftriaxone

Doc for gonococcal and non-gonococal urethritis → Azithromycin

Doc for candidiasis (fungal infection) → fluconazole

Anaerobic bacterial infections → Clindamycin

16. A patient was given anti-cancer therapy and developed neutropenia. Which of the following drug can be used to prevent neutropenia in next cycle of chemotherapy?

a) Filgrastim
b) Prednisolone
c) Vitamin B12
d) Folic acid

CSFs help your body make more white blood cells. This lowers your risk for febrile neutropenia. CSFs include **Neupogen (filgrastim), Neulasta (pegfilgrastim)**, and Leukine and Prokine (sargramostim). They are usually given as shots 24 hours after a chemotherapy treatment.

There are 3 types of cells in the body.

(1) RBC

(2) WBC

(3) Platelets

On bone marrow suppression due to anticancer therapy it will lead to

↓ RBC → Anemia

↓ WBC → Neutropenia

↓ Platelets → thrombocytopenia

www.medicalcoupons.in

www.medicalcoupons.in

17. Bacillus Anthrax is:

a) Gram positive cocci in cluster
b) Gram positive rods with square ends
c) Gram positive bacilli with spherical ends
d) Gram negative cocci in cluster

Bamboo Sticks appearance is seen in Anthrax

Gram positive cocci in cluster → staph aureus

Gram positive bacilli with spherical ends → Drumstick app seen in cl. Tetani.

18. Mark the incorrect statement about Corona virus :

a) Crown shaped peplomer like spike giving appearance of solar corona
b) Linear segmented positive sense single stranded RNA
c) Mostly infect animals except gamma coronavirus
d) Human corona virus 229E is alpha type

Corona virus

- Enveloped, carrying petal like crown shaped peplomer like spike giving appearance of solar corona
- 120-160 nm helical symmetry
- Linear positive sense single stranded RNA (Non segmented genome)
- Mostly infect animals except gamma coronavirus
- There are 6 types of coronavirus to infect humans
  → Human coronavirus 229E alpha
  → Human corona virus NL 63 alpha
  → Human coronavirus OC 43- beta
  → Human corona virus HK U1- beta
  → SARS Cov- beta
  → MERS- beta
  → SARS CoV-2 (COVID-19)

19. DNA transfer in bacteria via phage is:

a) Conjugation
b) Transduction
c) Transformation
d) Translation

www.medicalcoupons.in

Transduction is the process by which a virus transfers genetic material from one bacterium to another. ... Later, when one of these bacteriophages infects a new host cell, this piece of bacterial DNA may be incorporated into the genome of the new host. There are two types of transduction: generalized and specialized.

20. Sporulation occurs in this phase of bacterial growth curve:

a) Stationary phase
b) Lag phase
c) Log phase
d) Decline phase


1.**Lag phase**- Adaptation phase to culture media decreased Accumulate nutrient size

2. **Log Phase** - Exponential increases in number

3. **Stationary phase** - Start of Accumulation of toxic metabolites = Dead

Live -Sporulation occur in stationary phase (S-S)

4. **Decline -** Complete accumulation of Toxin metabolites



www.medicalcoupons.in

21. Which of the following is the most likely cause in a case of granuloma?
a) Cat scratch disease
b) Trench fever
c) Leprosy
d) Syphilis
Cat scratch disease is the most likely cause in a case of granuloma.

Various granulomatous diseases and their causative agents.

Cat-scratch disease - Bartonella henselae (Gram-negative bacillus): The necrotic foci enlarge and coalesce to form **necrotizing granulomas** called stellate micro abscesses.

22. HUS in children is caused by:
a) E. coli O57:H7
b) Malaria
c) Parvovirus B19
d) Bartonella henselae

EHEC

- Entero hemorrhagic E.coli

-Cause HUS (Hemolytic Uremic Syndrome)

-Shiga like toxin or Verotoxin (VT-1 & VT-2)

Aka verotoxigenic E.coli

Culture media is Sorbitol MacConkey agar (SMAC) and Rainbow agar used

- Diarrhea causing E coli

1. EPEC -Entero Pathogenic Attaching/Effacing lesions to villi

2. EIEC - Entero Invasive resemble shigellosis & sereny test +ve 3.

3. ETEC-> Traveler's diarrhea CFA (colonization factor antigen 4.

4. EAEC -Aggregative persistent type of diarrhea (EAST-1 Toxin)

Parvo Virus B19: Single standard DNA Virus cause Aplastic Crisis

23. N. meningitis can be due to deficiency of this complement system:

a) C1, C4 deficiency
b) C5-C9 deficiency
c) C3 deficiency
d) C2 deficiency

- C5-C9, MAC (membrane attack complex) deficiency cause Recurrent Neisseria infection.
- C3 deficiency is also associated with pyogenic infections and rheumatologic disorders.

24. Which of the following activates classical complement?

a) C1
b) C3 Convertase
c) IgA
d) Ag - Ab complex

The complement pathway. Complement can be activated through three pathways: classical, lectin, and alternative. **The classical pathway is activated when C1q binds to antibody attached to antigen**, activating C1r and C1s, which cleave C4 and C2.

FORENSIC DEC 2018

1. Gastric lavage, also commonly called stomach pumping or gastric irrigation, is the process of cleaning out the contents of the stomach. Since its first recorded use in the early 19th century, it has become one of the most routine means of eliminating poisons from the stomach. Such devices are normally used on a person who has ingested a poison or overdosed on a drug such as ethanol. They may also be used before surgery, to clear the contents of the digestive tract before it is opened. In which one of the following conditions is gastric lavage contraindicated?

a) Organo-Phosphosphate poisoning
b) Hydrocarbons
c) Bicarbonate
d) PCM toxicity

Contraindications of gastric lavage:

Absolute: Corrosives to prevent further damage from perforation (except carbolic acid - causes leathery mucosa of stomach - no risk of perforation).

Relative:

1. Convulsant - may induce more convulsions

2. Comatose - ↑ change of aspiration.

3. Varices - can cause bleeding

4. Volatile - hydrocarbons / kerosene: risk of aspiration reading to chemical pneumonitis.

2. Forensic experts investigated the crime scene. what confirmatory test will be used to prove a bloodstain?

www.medicalcoupons.in

a) Spectroscopic test
b) Kastle meyer test
c) Benzidine test
d) Orthotolidine test

SCREENING TEST FOR BLOOD STAINS	
I. Color test:	• Benzidine test / Adler's test: BLUE + • Phenolphthalein test : PINK + • Orthotolidine test: BLUE +
II. Luminescent test:	• Luminol spray • UV light

CONFIRMATORY TEST FOR BLOOD STAINS	
I.MICROSCOPY:	• To identify human RBC
II. ABSORPTION spectroscopy:	• Definitive confirmatory test • Most reliable • Used for recent / old stains.
III. Microchemical test	• Teichmann test / haemin crystal test [brown rhombic] • Takayama test / hemochromogen crystal [pink feathery]

3. The charred body has been recovered from the dense vegetation close to the highway. The police were led to the spot by a sniffer dog that was tracking a 25-year-old IT employee who had been missing since the previous evening. The team has to collect sample to identify the victim. A sample of which tissue from a burnt body, is best for identification?

a) Hair
b) Bone
c) Teeth
d) Blood

Identification of burnt bodies:

Hair: singed

Blood: burned

Bone: difficult to extract DNA from charred bones

Teeth: DNA can be extracted from pulp of the teeth even in a burnt body.

www.medicalcoupons.in

4. Death registration in India done within?

a) 21 days
b) 28 days
c) 30 days
d) 40 days

According to 'registration of birth & death act 1969 every birth / still birth / death to be registered with the covered state / UT within 21 days.

Registration of vital events

→ Birth - 21 days
→ Deaths - 21 days
→ Marriages - 30/60/90 days

- Birth registration is the responsibility of hospital
- Child born to NRI couple abroad , birth registration done within 60 days of arrival

5. Estimation of age at death and determination of sex of the victim or remains are important factors in the identification of an individual in forensic odontology. Teeth are among the most reliable tools in the process of identification of age. Teeth are one of the most durable parts of our body, which can withstand more assaults than any other part of the body. This is particularly useful in the identification of bodies in mass disasters and natural calamities. The first scientific technique for age estimation in adults was presented by Gustafson {1950}. It was based on longitudinal sections of teeth cut through the central area. The technique consisted of attributing scores from 0-3 for the presence and amount of age-related changes such as attrition, periodontal ligament retractions, secondary dentin formations, root translucency, and root resorption. The most reliable criteria among Gustafson's criteria is?

a) Root resorption
b) Paradentosis
c) Translucency of root
d) Attrition

Gustafson's criteria: age determination using secondary changes in teeth.

- A. Attrition
  P. Paradentosis
- S. Secondary dentin 2nd most reliable
  R. Root resorption
- T. Transparency of root most reliable
  Cementum opposition

www.medicalcoupons.in

SPM DEC 2018

1. **Mosquito having spotted wings is**

a) Anopheles
b) Culex
c) Aedes
d) Mansonia

	Anopheles	Aedes	Culex
			
Adult	• Sit at 45° spotted wings	• Hunchback posture • White stripes on body (Tiger Mosquito)	• Hunchback posture
Egg	• Boat shaped • Lateral floats	• Single • Cigar shape	• Small • Clusters
Larvae	Rest parallel to water surface (No siphon tube)	Rest at an angle to water surface (Short Siphon tube present)	Rest at an angle to water surface (Slender Siphon tube present)

2. **Population covered by one PHC in plains is**
a) 3000
b) 5000
c) 20000
d) 30000

www.medicalcoupons.in

Population covered by	PLAINS	HILLY / TRIBAL
Subcenter	1/5000	1/3000 (MPW)
PHC	1/30000	1/20000 (Health Assistant)
CHC	1/12000	1/80000
AWC	1/400 - 800	1/300 - 800

3. Treatment of AIDS (ART) is started at CD4 cut off

a) 200
b) 350
c) 500
d) No cut-off

Treatment of AIDS is ART

ART initiation -irrespective of CD_4 count

→ Any age
→ Any stage
→ Any group

Start ART

Normally CD4 count is 900 - 1400 & if it starts going below 500 opportunistic infections start.

Most common opportunistic infection HIV world → Pneumocystis carinii Pneumonia

Most common opportunistic infection HIV India → TB

4. Normal Waist to hip-ratio

a) 0.8 in males
b) 0.85 in females
c) 0.90 males
d) < 0.95 males

Waist to hip ratio used as indirect marker of obesity and direct marker of cardiovascular risk factor the body.

Cutoffs :

www.medicalcoupons.in

Male ≤ 0.90

Female ≤ 0.85

According to new WHO guidelines waist hip ratio has been replaced by a new indicator that is "Waist Height Ratio"

WHO SAYS, this is now the best indicator for cardiovascular risk.

Age and sex independent

Single cutoff for al age and sex → 0.5

5. Rapid diagnosis of Rabies in a Rabid living dog is by

a) Fluorescent antibody test
b) Corneal impression
c) Inoculation in mouse
d) None of the above

- Flourescent Antibody Test : This is a highly reliable and the best single test currently available for the Rapid diagnosis of Rabies viral Antigen in Infected specimen.
- Corneal impression & Mouse inoculations are also used for diagnosis.

6. 1 year old with Keratomalacia, treatment is

a) 1 Lac IU once
b) 1 Lac IU three times
c) 2 Lac IU once
d) 2 Lac IU three times

Keratomalacia is manifestation of vit. A deficiency syndrome known as Xerophthalmia

Xerophthalmia Treatment (oral vit. A)		
	≤ 1 year age	**>1 year age**
Day 0	1 Lac IU	2 Lac IU
Day 1	1 Lac IU	2 Lac IU
Day 14	1 Lac IU	2 Lac IU

IM vit. A dose will be exactly half of oral dose.

www.medicalcoupons.in

7. A 3-5 months old child with fever, nasal congestion, sore throat and chills presented to the OPD with his mother. Nasopharyngeal swab confirmed H1N1 as the causative agent. Which of the following is the correct treatment dose of Oseltamivir?

a) 20 mg OD X 5 days
b) 12 mg BD X 5 days
c) 25 mg BD X 5 days
d) 20 mg BD X 5 days

Oseltamivir (Tamiflu) is the drug of choice for swine flu H1N1. But it's dose is given according to the age and weight of the person.

Age	>3months to 12 months → mg/kg BD x 5 Days
	>1 month to 3 months → 2.5 mg/kg BD x 5 Days
	0 to 1 month → 2 mg / kg BD x 5 Days

Weight	<15 kg →g 30 mg BD x 5 Days
	15-23 kg → 45 mg BD x 5 Days
	24-40 kg → 60 mg BD x 5 Days
	>40 kg → 75 mg BD x 5 Days

8. Beri-Beri seen in

a) Sorghum eaters
b) Wheat esters
c) Polished rice eaters
d) Maize eaters

Vitamin B1 Deficiency (Thiamine)

→ **Beri-Beri** → Seen in Polished rice eaters
→ **Wernickes korsakoff psychosis** → seen in Alcoholics

9. Most common cause of LBW in India

a) Smoking during pregnancy
b) Maternal tetanus
c) Intra-uterine growth retardation
d) Premature birth

www.medicalcoupons.in

The primary cause is premature birth, being born before 37 weeks gestation, a baby born early has less time in the mother's uterus to grow and gain weight, and much of a fetus's weight is gained during the latter part of the mother's pregnancy. **Another cause of low birthweight is intrauterine growth restriction.**

10. The proposal to control malaria in towns named as Urban Malaria Scheme. According to Urban Malaria Scheme, Which of the following is not true?

a) Utilization of anti-larva measures
b) Slide positivity rate more than 10%
c) It is done in town/city with minimum 50,000 population
d) Introduction of active surveillance

Urban malaria has three specific criteria-

→ Utilization of Anti-Larval measures.
→ It is done in town/city with minimum 50,000 population.
→ Slide positivity rate more than 5% [active surveillance].

11. A 30 year old male truck driver visited to the clinic with fever, skin rash, swollen glands and fatigue. The physician referred him to suraksha clinic. Which among the following is included in provision of services a Suraksha Clinic ?

a) Immunization services
b) Blood transfusion safety services
c) RTI/STI services
d) Diarrhea control services

Services expected at suraksha clinic (STD clinic) are related to RTI/STI. It includes :

→ Diagnostic services
→ Treatment care
→ Preventive services



www.medicalcoupons.in

12. Ujjawala scheme is for prevention of

a) Child abuse
b) Child trafficking
c) Child labour
d) Child marriage

UJJWALA	Providing LPG connections to women from Below Poverty Line (BPL) households.
UJJAWALA	Prevention of trafficking and Rescue, Rehabilitation and Re-integration of Victims of Trafficking and Commercial Sexual Exploitation
UJALA	Unnat Jyoti by Affordable LEDs for All Promote efficient lighting, enhance awareness on using efficient equipment which reduce electricity bills and help preserve environment.

13. Peripheral blood smear sample for malaria is done by

a) Female MPW
b) ANM
c) Medical office
d) Male MPW

Multi Purpose Worker (MPW)	
MPW Male	**MPW Female**
• Field work • Peripheral blood smear	• Reproductive and Child Health (***RCH***) • Family planning • Record

→ Located at subcenter
→ Education - 12th pass
→ 1st introduced by Kartar singh committee
→ Training - 12 months


14. A government school in rural village was checking all students for their health conditions. Under Vision 2020, to check visual acuity, teacher will refer school children's to which of the following centre?

a) Service centre
b) Training centre
c) Centre for excellence
d) Vision centre

The goal of Vision 2020 is to eliminate needless blindness by the year 2020. The role played by Vision Centre in achieving this target by providing primary eye care in the community therefore assumes significant importance.

www.medicalcoupons.in

Vision2020 has developed certain policies for Vision Centre:



15. A school has a small outbreak were children are showing symptoms like runny nose, cough, fever and skin rash, The doctors identified it as measles outbreak. Which of the following is the correct time to give post exposure measles vaccine?

a) 3-4 days
b) 7-10
c) 10-14 days
d) 1-6 months

- IP of measles →10 days
- IP of vaccine induced measles → 7 days
- For providing protection vaccine requires only 6-7 days.
- So, vaccine should be used within 3-4 days of exposure.

Cholera period of Isolation

3 days after starting Tetracycline till 48 hours of antibiotics

7 days after starting Tetracycline

-14 days

Until 3 weeks of effective antibiotic therapy

www.medicalcoupons.in

Cholera:

- Caused by - vibrio cholera
- Field microscopy test - hanging drop
- IP - 1-2 days
- Rice watery stool

16. Additional energy requirement during Pregnancy (Kcalories/day)

a) +300 kcal
b) +600 kcal
c) +350 kcal
d) +520 kcal

Additional Energy Requirements	
2nd and 3rd Trimester	1st Trimester → 150
	2nd & 3rd Trimester → 350
Lactation	0 – 6 months → + 600
	6 – 12 months → + 520

17. Highest Contraceptive failure is reported in

a) Spermicidal methods
b) Implant
c) Oral contraceptive pills
d) IUD

Spermicidal methods has pearl index of 10-25 / HWY(Highest).

Contain chemical - Non oxynol 9

MOA - By rupture of plasma membrane of Acrosomal cap.

18. While launching a new national health programme and policy population attributable risk is calculated. Which of the following best defines population attributable risk?

a) Incidence of disease among exposed is so many times higher as compared to that among non-exposed
b) So much disease can be attributed to exposure
c) If risk factor is modified or eliminated, there will be so much annual reduction in incidence disease in the given population.
d) None of the above

www.medicalcoupons.in

Population Attributable Risk (PAR)

- It is the incidence of disease (or death) in total population minus incidence of disease (or death) among those who were not exposed to suspected causative factor.
- It is useful as it provides an estimation of amount by which incidence of disease could be reduced in that population if the suspected factor was eliminated or modified.
- PAR is useful for National health programmes and policies.

19. Human anatomical wastes are disposed in ___ BMW (Bio Medical Waste) category

a) Red
b) Yellow
c) White
d) Blue

- **Yellow category**
  → Human anatomical waste
  → Animal waste
  → Soiled waste
  → Chemical waste
  → Discarded medicine
  → Liquid chemical waste
  → Microbiological waste
- **Red category:** Solid contaminated (Recyclable) (plastic/rubber)
- **White category:** Wasted sharp
- **Blue category:**
  → Glassware
  → Metallic body implants

20. A disaster is a sudden, calamitous event that seriously disrupts the functioning of a community or society and causes human, material, and economic or environmental losses. Which among the following covers Disaster management?

a) Ministry of Health
b) Ministry of HRD
c) Ministry of Home
d) Ministry of Agriculture

A disaster is a sudden, calamitous event that seriously disrupts the functioning of a community or society and causes human, material, and economic or environmental losses.


www.medicalcoupons.in

Disaster management

→ Nodal ministry - ministry of home affairs
→ Nodal centre – District

21. **Phase(s) of Disaster cycle before disaster impact include**

a)Disaster response
b)Medial relief and rehabilitation
c)Mitigation and preparedness
d)Rehabilitation



22. At a PHC, vaccine storage is done in

a) Vaccine carrier
b) ILR
c) Walk in Cold rooms
d) Cold box

At the PHC (Primary health centre) level, all vaccines are kept in **the ILR** for a period of one month at temperature of +2°C to +8°C

www.medicalcoupons.in

23. A specific package of intervention and services which are given to population those who have either high risk of HIV or those who have reported high risk of HIV IS known as targeted intervention. Which of the following group is not involved in this intervention?

a) Migrant labourer's
b) Street children
c) Long distance truckers
d) Doctors and nursing professionals

Targeted intervention :

It is a specific package of intervention & services which are given to population those who have either high risk of HIV or those who have reported high risk of HIV.

Groups with high risk of HIV are:

→ Commercial sex workers
→ Injecting Drug Users
→ Truck Drivers
→ Migrant Labour
→ Street children
→ Transgender
→ Adolescents

24. A patient from rural village was given ORS + Zinc. What does ORS plus Zinc do?

a) Reduces infection
b) Antispasmodic
c) Reduces duration of diarrhea
d) Enhance sodium absorption

Zinc supplementation with ORS therapy reduce

→ Duration of diarrhea
→ Severity of acute and persistent diarrhea
→ Volume of diarrhea

www.medicalcoupons.in

ENT DEC 2018

1. **Mainstay of treatment of glue ear:-**

a) Radical Mastoidectomy
b) Myringotomy + aeration to middle ear
c) Temporal bone resection
d) Tonsillectomy & adenoidectomy

Mainstay treatment of glue ear should be treat the cause, like if its adenoids or tumor so removing it would be the treatment. But the cause is not mentioned in the question. **So the mainstay treatment should be myringotomy in the antero-inferior quadrant with grommet tube insertion.**

2. **A patient presented with 2 days history of fever. On examination there was swelling in the neck and one side tonsil was pushed to midline. What is the most likely diagnosis:-**

a) Quinsy
b) Parapharyngeal abscess
c) Retropharyngeal abscess
d) Tonsillitis

In the history of the patient it is given that the tonsil is pushed to the midline and there is a swelling in the neck, this is the typical clinical feature of parapharyngeal abscess.

3. **Most common symptom of nasopharyngeal carcinoma:-**

a) Airway obstruction
b) Headache
c) Neck swelling
d) Epistaxis

Symptoms of nasopharyngeal cancer may include:

- Lump in neck (most common)
- Blurry or double vision.
- Ear infections that recur.
- Face pain or numbness.
- Headache.
- Hearing loss, ringing in the ears, or a feeling of fullness in the ear.
- Difficulty opening mouth.

www.medicalcoupons.in

- Nosebleeds.

4. Picket fence fever is seen in:-

a) Lateral sinus thrombophlebitis
b) Quincy
c) Laryngitis sicca
d) Croup

- Picket fence fever:- fever never comes back to normal baseline. Also known as remittent fever. (intermittent fever is malaria).
- This is the typical presentation of lateral sinus thrombosis/sigmoidal sinus thrombosis.

5. Cold water is not used for syringing because:-

a) Cold water leads to wax impaction
b) Cold water leads to foreign body impaction
c) Can cause tympanic membrane rupture
d) It causes vertigo by caloric stimulation

Syringing

- Simpsons aural syringe (50ml) is used
- Syringing is done in the posterior superior quadrant.
- The temperature of the water should be ideally 37º C.
- If we use cold water it stimulates labyrinth and the patient has vertigo.
- Cold water 30^0 C & 4^0 C is warm water used in caloric testing to check if labyrinth is working.

6. Which of the following muscles originates from the first pharyngeal arch:-

a) Stapedius muscle
b) Anterior belly of digastric muscle
c) Posterior belly of digastric muscle
d) Stylopharyngeus

The first arch gives rise to the muscles of mastication and the **mylohyoid, anterior belly of digastric, tensor veli palatani and tensor tympani** – all of which are innervated by the branches of the trigeminal nerve.

www.medicalcoupons.in

7. Dip at 4000hz in pure Tone audiometry indicates:-

a) Otosclerosis
b) Meniere's disease
c) Noise induced hearing loss
d) Age related hearing loss

2000Hz / Carhart's Notch: seen in otosclerosis (CHL)

4000Hz Dip/boiler's notch :seen in noise induced hearing loss (SNHL)

Both are seen in bone conduction.

8. Which of the following structures is used as a landmark for facial nerve in parotid gland surgery?

a) Tragal pointer
b) Posterior border of mylohyoid
c) Tip of helix
d) Styloid process

Surgical landmarks:

- The tympanomastoid suture is one of the landmarks that's used.
- We also use the tragus cartilage/tragal pointer. It's the most commonly used landmark in identifying the facial nerve.
- Styloid process can also be used but there are high chances of nerve damage
- Posterior belly of digastric is also a landmark.

9. Continuous watery discharge from nose after trauma is most likely a feature of::

a) CSF otorrhea
b) Common colds
c) CSF rhinorrhea
d) Anterior epistaxis

Watery discharge from nose always implies CSF.

CSF Rhinorrhea:

MC cause Head injury/trauma due to fracture of cribriform plate of ethmoid.

www.medicalcoupons.in

10. Which of the following is not a premalignant lesion / condition:-

a) Chronic hypertrophic candidiasis
b) Oral submucosal fibrosis
c) Lichen planus
d) Leukoplakia

Chronic hypertrophic candidiasis is a long term infection of candida seen in the middle aged male patients, smokers. There is a membrane formation, But its not a premalignant condition.

Premalignant lesion:

→ Oral submucous fibrosis
→ Lichen planus
→ Leukoplakia (MC premalignant lesion seen in smokers & tobacco chewers)

11. Hypoglossal nerve supplies all of these muscle except:-

a) Palatoglossus
b) Styloglossus
c) Genioglossus
d) Hyoglossus

The hypoglossal nerve supplies:

- All of the intrinsic muscles of the tongue.
- Three extrinsic muscles of the tongue: the genioglossus, hyoglossus and styloglossus muscles.
- As well as the geniohyoid muscle (a suprahyoid muscle)

12. After head trauma a patient in unable to smell coffee but can smell ammonia. Which of the following statements is true about the patient?

a) Ammonia is used for testing olfactory nerve
b) The olfaction system is not damaged
c) Ammonia irritates trigeminal nerve
d) All of the above

www.medicalcoupons.in

Ammonia has pungent smell so, there is orthonasal olfaction & the odour goes directly to cribriform plate and directly stimulates the CN - IX, V & X. so it can curse smell even when the olfactory Nerve is not working.

13. Which of the following is an accessory masticatory muscle?

a) Buccinator
b) Masseter
c) Medial pterygoid
d) Lateral pterygoid

Masticatory muscles are

- Masseter
- Temporalis
- Pterygoids (Lateral & Medial)

These are Supplied by the mandibular nerve


→ Buccinator is a secondary masticatory muscle,
→ Its main function is facial expression.
→ Hence, it's called an accessory masticatory muscle.

OPTHAL DEC 2018

1. True about pterygium

a) Inflammation of cornea
b) Fatty deposition of sclera
c) Conjunctival invasion making a flap over cornea
d) Dead epithelial debris accumulation

Pterygium:



→ Wing shapes fold of conjunctiva encroaching upon the cornea from the either side

Etiology-

→ Hot climates
→ Degenerative & hyperplastic condition of conjunctiva

Symptoms

→ Visual disturbance (astigmatism)
→ Cosmetic intolerance

Sign -

→ Triangular fold of conjunctiva mainly on nasal side
→ Stocker line - Iron deposition on corneal epithelium

Rx-

→ Surgical excision-recurrence chance is 30-50 %
→ Best RX- Surgical excision with free conjunctival autograft taken from the same eye Of other

2. For laser iridotomy which laser is used?

a) Argon laser
b) Nd YAG laser
c) Excimer laser
d) Co2 Laser

LASER	EFFECTS	CLINICAL USE
ARGON LASER	Photocoagulation	Diabetic Retinopathy
ND YAG LASER (1064nm)	Photodisruption	CAPSULOTOMY, IRIDOTOMY
EXCIMER (193nm)	Photoablation	LASIK, Phototherapeutic Keratectomy
CO2 LASER	Incisional device Resurfacing device	Oculoplasty

3. A newborn is hypermetropic by

a) +2.5 D
b) + 5 D
c) + 10 D
d) +1D

NORMAL - REFRACTIVE error at birth 2-3 D hypermetropic

www.medicalcoupons.in

→ By 5-7 years of age the eye is emmetropic and remains so till the age of about 50 years
→ After 50 years of age there is tendency to develop hypermetropia again which gradually increase the extreme of life by which the eye has the same +2 TO +3 with which it started. this senile hypermetropia is due to change in the lens.

4. Corneal endothelium is derived from:

a) Neuroectoderm
b) Neural crest cells
c) Mesoderm
d) Surface ectoderm

SURFACE ECTODERM	NEURAL ECTODERM	MESODERM	NEURAL CREST
• LENS • Epithelium of the all structures except ciliary body & iris • Glands	• Epithelium of ciliary body & iris • Retinal pigment epithelium • Optic nerve • Sphincter and dilator pupillae muscles ▪ Smooth muscles of iris	• Extraocular Muscles • Endothelium of all the orbit & stroma of iris & ciliary body • Sclera (temporal part)	• Stroma of cornea & choroid • Corneal endothelium • Sclera (except temporal part) • Melanocytes • Corneal keratocytes • Ciliary muscles • Ciliary ganglion • Stroma of iris • Choroid Trabecular meshwork

5. True about retinoscopy with a plane mirror

a) In hypermetropia the red glow moves in opposite direction
b) In myopia the red glow moves in same direction
c) In emmetropia the red glow moves in opposite direction
d) Performed at 1 meter away from patient

Retinoscopy

→ Differentiate b/w type of refractive errors
→ Optical error & organic error
→ Performed at 1 meter away

www.medicalcoupons.in

6. Rosette cataract is seen after

a) Blunt trauma to eye
b) Copper foreign body in eye
c) Penetrating injury to eye
d) Infection

Cataracts caused by blunt trauma classically form stellate- or rosette-shaped posterior axial opacities that may be stable or progressive, whereas penetrating trauma with disruption of the lens capsule forms cortical changes that may remain focal if small or may progress rapidly to total cortical opacification.

7. Patient with history of fall on symmetrical surface presents with enophthalmos diplopia on upward gaze and loss of sensitivity over cheek. True statement about this is :

a) Maxillary fracture
b) It is a blowout fracture
c) Zygomatic bone is most likely injured
d) Frontal bone fracture

Blow out fracture -

- MC fracture of orbit
- Inferior wall M/C involved
- Entrapment of inferior rectus - inferior wall fracture - not able to look up

Features

→ Enophthalmos
→ Diplopia
→ Infra orbital nerve anesthesia

Hanging drop or tear drop sign on imaging

8. Blow out fracture can be due to:

a) Tennis ball injury
b) Chisel and hammer injury
c) Sudden fall
d) Punch at the chin from below

- Blow out orbital fracture result from trauma to orbit by a relatively large often rounded object such as tennis ball, cricket ball, human fist, or part of automobile.
- The force of blow causes a backward displacement of the eyeball and increase in the intraorbital pressure.

www.medicalcoupons.in

9. Sarcoidosis is associated with

Cataract

Ectopia lentis

Anterior uveitis

Keratitis

Sarcoidosis -Can cause

→ Anterior uveitis (M /C)
→ Posterior uveitis
→ Intermediate uveitis
→ Pan uveitis
→ **M/C presentation** - B/L Lymphadenopathy

10. 10 years old child brought to your clinic with developmental delay associated with extrapyramidal symptoms. On further examination, dislocation of the lens towards the inferonasal area is revealed. which of the following is the most probable diagnosis?


a) Homocystinuria
b) Refsum's disease
c) Wilson's disease
d) Sarcoidosis

Congenital ectopia lentis is seen in :

→ Marfan's syndrome - supratemporal
→ Homocystinuria - inferonasal
→ WEILL-MARCHESANI SYNDROME -Forward subluxation of lens
→ Ehlers Danlos syndrome -subluxation of lens blue sclera.

www.medicalcoupons.in

11. Obstacles in the condition shown in the image is:



a) Motor obstacles
b) Sensory obstacles
c) Central obstacles
d) All the above

The given image shows a Concomitant squint

CONCOMITANT SQUINT - a type of manifest squint which the amount of deviation in the squinting eye remains constant in all directions of gaze, with no limitation of ocular movements. Any obstacles in the development of these processes result in concomitant squint.

Sensory obstacles-hinder the formation of a clear image in one eye.

E.g.

- Refractive error
- Prolonged use of incorrect spectacles
- Anisometropia
- Media opacities - Corneal/Lenticular opacities

Motor obstacles - hinder the maintenance of eyes in the correct positional relationship in primary gaze or during ocular movements.

E.g.

- Congenital abnormalities of shape/size of the orbit
- Abnormalities of extraocular muscles

Central obstacles - Deficient development of fusion faculty.

www.medicalcoupons.in

Abnormalities of cortical control of ocular movements as occurs in mental Trauma and hyperexcitability of CNS during teething.

OBG DEC 2018

1. **A woman at 26 weeks gets ultrasound done for fetal well-being and is diagnosed with hydrops fetalis. Which of the following fetal infections could have led to this condition?**

a) Ebstein barr virus
b) Human papilloma virus
c) Parvovirus B19
d) Influenza Virus

Causes of hydrops fetalis -

- Cardio vascular cause → congenital heart block (Most common)
- Chromosomal cause → turner syndrome (45 XO), triploidy , trisomy's 21,18 and 13

Hematological → α 4 - thalassemia (Hemoglobin Bart's), fetomaternal hemorrhage

- **Infections → Parvovirus B19(mc)** , syphilis, cytomegalovirus, toxoplasmosis Rubella, HSV.

Hydrops fetalis -

- Excessive destruction of the fetal red cells leads to severe anemia, tissue anoxemia and metabolic acidosis
- It is accumulation of fluid in the body of the fetus in two out of three compartments: ascites, pleural effusion, pericardial effusion
- Most common cause of immune hydrops fetalis - Rh incompatibility.

2. **Childbirth is the single most common cause of uterovaginal prolapse in our country. All the events during child birth may account for risk of uterine prolapse in future, EXCEPT?**

a) Precipitated labour
b) Prolonged 2nd stage
c) Premature bearing down
d) Episiotomy

<u>Episiotomy will not cause prolapse</u>

Main cause of prolapse is abnormal conduct of labor***

www.medicalcoupons.in

Abnormal conduct of labor is by :

→ Prolonged 2nd stage
→ Faulty instrumentation
→ Early bearing down
→ Cervical lacerations & muscle lacerations

3. Conservative treatment for pelvic organ prolapse is

a) Perineorrhaphy
b) Shirodkar Procedure
c) Pessary
d) Le Fort Repair

Surgical treatment of prolapse	→ Shirodkar procedure → Perineorrhaphy → temporary Rx → Le fort repair

Conservative treatment of prolapse	→ Pessary (should be replaced every 3 months)

Management of Genital Prolapse	
Nulliparous	Abdominal sling operations
Pregnancy	Ring pessary up to 16 weeks
Postnatal	→ Ring pessary and pelvic floor exercises for 3-6 months. → Surgery if required thereafter.
Young woman <40 years	Conservative vaginal surgery (fertility sparing surgery) → Cystocele, rectocele repair → Manchester repair → Sling operation
Woman beyond 40 years and multipara	Vaginal hysterectomy and pelvic floor repair

4. Cut off point for 50 gram glucose challenge test at 1 hour mg%

a) 100
b) 140
c) 180
d) 200

www.medicalcoupons.in

Glucose challenge test :

50 grams of glucose to non-fasting women :

After 1 hour

→ **After 1 hour** → Value: < 140 normal
→ >140 → May be gestational diabetes mellitus.

- We do the diagnostic test i.e. glucose tolerance test
- >200 - gestational diabetes mellitus

Glucose tolerance test

100 grams of glucose to fasting women

- Fasting value → < 105

Value:	
1 hr - < 190 2 hr - < 165 3 hr - < 145	any 2 abnormal values then it is diagnostic of gestation diabetes mellitus.

5. 28 year old lady (G2 P1) who does not want to conceive after abortion, which is the preferred mode of contraception

a) OCP
b) IUD
c) Barrier
d) Implant

Contraception → IUD > OCP > Barrier

6. A 30 year old primigravida delivers a full term baby. However she chooses to formula feed her newborn. After how much duration can she start using combined oral contraceptives for avoiding pregnancy?

a) 2 weeks
b) 6 weeks
c) 12 weeks
d) Immediately after delivery

Post-partum use of combined oral contraceptives

www.medicalcoupons.in

- COC are contraindicated in breast feeding mothers (causes breast milk suppression) and can be started after **6 weeks** post-partum.
- COC can be used in non-breast feeding mothers after **3 weeks** if there are no persisting risk factors for VTE such as smoking, history of DVT, known thrombogenic mutations, peripartum cardiomyopathy
- COC can be safely started after **6 weeks** in non-breast feeding mothers
- Here in this question, since there is no history suggestive of any risk factors for VTE, the better answer would have been 3 weeks, but since it is not in the options, the answer would be 6 weeks.

Please note that **COC should never be started immediately post-partum** in any scenario since post-partum period in itself is a high risk state for thromboembolism, maximum risk being in the first week. Progesterone only pills can be safely used in lactation, and can be started as soon as contraception is required.

7. A 43 year old woman, from low socio-economic status, a chronic smoker, hypertensive, P3L3, tests positive for type 16 HPV on cervical cancer screening. She now uses IUCD for contraception. All of the following factors in her history predispose to her acquiring cervical cancer, EXCEPT?

a) HPV 16, smoking
b) Hypertension and IUCD use
c) Multiparity and low socioeconomic status
d) Multiparity and smoking

Risk factors for development of cervical cancer

- HPV
  → 16: Most common
  → 18: most malignant
- HSV 1 and.
- HIV 1 and 2
- Commercial sex worker
- Women with many sex partners
- Partner with STD
- Early intercourse (< 16 yrs)
- Low Socio Economic status
- Smokers

(Expected Date of Delivery) is calculated by:

a) Young's rule
b) Naegele's formula
c) Cardiff Formula
d) Hadlock Formula

www.medicalcoupons.in

Expected date of delivery is calculated by **Naegele's formula**

From the 1st day of last menstrual cycle → 9 months + 7 days.

9. If a zygote divides 2-3 days after fertilization which of the following type of twining is seen:

a) Monochorionic, Monoamniotic
b) Monochorionic, Diamniotic
c) Diamniotic, Dichorionic
d) Dichorionic, Monoamniotic


If a zygote divides :	Type of twining
Before 3rd day of life	→ Dichorionic Diamniotic
4th to 8th day of life	→ Monochorionic Diamniotic
After 8th day of life	→ Monochorionic Monoamniotic
After 14th day of life	→ Conjoined twins / Siamese twins

10. Chromosome pattern in complete mole:

a) 46 XY both maternal origin
b) 46 XX both maternal origin
c) 46 XX both paternal origin
d) 46 YY both paternal origin

Complete mole

A complete mole may be formed if. 23 , - X bearing haploid sperm penetrates a empty ova paternal chromosome then duplicate to create 46 X. diploid complement Solely of paternal origin



www.medicalcoupons.in

11. A woman is diagnosed with an unruptured tubal ectopic. She is admitted for medical management with methotrexate while a strict watch is kept for signs of tubal rupture. An ectopic sac in which of the following parts of the tube is likely to rupture the earliest in case of failed medical management?

a) Interstitial
b) Isthmus
c) Ampulla
d) Infundibulum

Fate of Ectopic pregnancy -Tubal abortion

→ M/C site s ampulla
→ 2nd M/C site S infundibulum

Rupture of the tube:

→ M/C site is Isthmus (4-6 weeks)
→ Longest survival → interstitium (12 weeks)

12. A pregnant woman with a known history of fibroid uterus is being planned for an elective cesarean section for breech presentation. She insists her fibroid be removed with the surgery, however the gynaecologist counsels her the regarding disadvantages of myomectomy at the time of cesarean and tells her that he would remove it only if it is ?

a) Pedunculated subserosal
b) In the broad ligament
c) Posterior in location
d) Small in size

A myomectomy should not be performed during a caesarean section as the uterus is extremely vascular at this time and any attempt at a myomectomy can lead to torrential bleeding. However, pedunculated fibroid may be removed by clamping and ligating the pedicle, if easily accessible.

13. After 28 weeks POG major amount of amniotic fluid is secreted from:

a) Fetal skin
b) Fetal urine
c) Placental cell membrane
d) Plasma

www.medicalcoupons.in

In early pregnancy	Ultrafiltrate of maternal plasma
Beginning of 2nd trimester	Extra cellular fluid which diffuses through fetal skin
After 20 weeks	Fetal urine > fetal lung

14. Which of the following infections can commonly lead to infertility:

a) Chlamydia
b) Trichomoniasis
c) Candidiasis
d) Bacterial vaginosis

Major infectious cause of infertility is chlamydia

Chlamydia is an indolent PID (Quit / symptomless)

15. A young couple presents to fertility clinic with inability to conceive since years. They are living together,have been married for 4 years and not using any contraception at present. The initial tests that should be advised to this couple are?

a) Semen analysis, CXR, mantoux
b) Semen analysis, Ovulation test, Tubal patency test
c) Ovulation, tubal patency, Mono-dux test
d) Testicular biopsy, USG, Sperm penetration test

Order for initial work - up for infertility

1. Semen analysis

2. Ovulation test

3. Tubal patency

Note: Tubal patency test is done by 2 ways

1 . HSG (initial test)

2. Laparoscopy with Chromopertubation (Confirmatory)

16. A G4P2 lady presented with history of two abortions at 16 weeks and 20 weeks POG. Which of the following could be the most likely reason for these abortions?

www.medicalcoupons.in

a) Chromosomal abnormality
b) Cervical incompetence
c) Placenta previa
d) Thyroid abnormality

Abortions- MC causes

→ 1st trimester - chromosomal causes
→ 2nd trimester - Anatomical cause

17. True about endometriosis:

a) Presence of endometrial gland in deep myometrium
b) Presence of endometrium at ectopic locations
c) Treated preferably with hysterectomy
d) Seen in multiparous women

Endometriosis

→ **Presence of endometrial tissue at ectopic location**
→ Etiology : Retrograde menstruation
→ Proposed by Sampson (Sampson's) Implantation theory
→ 70-80% of all women have retrograde menses
→ Among them 5-10% have poor immunity and ↑ estrogenicity and develops Endometriosis

NOTE: Option A, C, D. are related to Adenomyosis.

18. A 29 year old female presented with infertility. There is history of abdominal pain, dyspareunia, dysmenorrhea, menorrhagia. Most likely cause:

a) Adenomyomatosis
b) Endometriosis
c) Myomas
d) Cervicitis

Symptoms of Endometriosis :

- Dysmenorrhea
- Abdominal pain
- Dyspareunia
- Infertility
- Menorrhagia
- Chronic pelvic pain

www.medicalcoupons.in

19. A 32 year old pregnant woman presents with 36 week pregnancy with complains of 01 pain abdomen and decreased fetal movements. Upon examination PR= 96/min, BP = 156/100 mm Hg, FHR = 128 bpm. On per-vaginum examination there is altered blood seen and cervix is soft 1 cm dilated. What is the preferred management?

a) Tocolytics
b) Induce labour
c) Immediate LSCS
d) Wait and watch

This appears to be a case of abruption with a normal fetal heart rate.

The patient should be induced to expedite delivery with a strict watch over maternal vitals and fetal heart rate.

20. A 30 year old woman presents to the clinician with excessive growth of facial hair associated with hairgrowth on her upper back, arms and thighs. Which of the following is the most likely cause?

a) Drug induced
b) PCOS
c) Endometriosis
d) Adenomyosis

Excessive hair growth in male pattern in women is known as **hirsutism**

Hirsutism

- M/C cause : PCOS
- Due to Hyperandrogenism
- Increase Androgen → increase Serum Testosterone & Androstenedione →decrease Sex hormone binding globulin → increase Free Androgen → Hirsutism.

21. 52 year old lady presents to the physician with complains of episodes of feeling of intense warmth followed by profuse sweating. She stopped having her periods since the past 3 years. Which of the following is correct regarding her condition?

a) Increased FSH, increased estrogen
b) Decreased FSH, decreased estrogen
c) Decreased FSH, increased estrogen
d) Increased FSH, decreased estrogen

The patient has presented with post-menopausal hot flashes.

www.medicalcoupons.in

Due to exhaustion of ovarian function after menopause, the estrogen levels declin[illegible]s a result of low estrogen, the feedback inhibition of gonadotropins IS lost, thus increasing levels of FSH and LH.

- FSH > 10 is suggestive of menopause
- FSH > 40 is diagnostic of menopause

22. A primigravida presents at 9 weeks of amenorrhea with a p[illegible]tive pregnancy test. She complains of nausea and vomiting which gets better by ev[illegible]ning. The doctor advices her folic acid and dietary modifications to help w[illegible]h he[illegible] vomiting. He explains that the cause of such presentation is a horm[illegible]ne. [illegible]e source of this hormone is?

a) Yolk sac
b) Liver
c) Syncytiotrophoblast
d) Umbilical cord

- Pregnancy **nausea is directly linked t[illegible] the pregnancy hormone hCG**, Early in pregnancy, when morning sickness [illegible] typically at its worst, hCG levels rise rapidly to help grow and nourish the fertiliz[illegible]d [illegible]gg
- **Human chorionic gonadotrophin (hCG)** is a glycoprotein hormone synthesized in the rough ER of **syncytiotrophoblast** immediately after implantation of the blastocyst into the uterine w[illegible]ll.

PEDIA DEC 2018

1. Continuous murm[illegible]s heard in?

a) Patent ductus arte[illegible]sus
b) Ventricular s[illegible]pta[illegible]defect
c) Atrial sept[illegible]efect
d) Tetralogy of fa[illegible]t

PDA → Continuous murmur

VSD → P[illegible]n systolic murmur

A[illegible]D → Silent, wide fixed split of S2 is seen

T[illegible]E → Ejection Systolic Murmur

www.medicalcoupons.in

2. Menke's disease is due to defect in the metabolism of:-

a) Zinc
b) Copper
c) Selenium
d) Iron

Menke's disease: aka kinky hair disease

- X- linked recessive disorder caused by mutations in genes coding for the copper transport protein ATP7A.
- Copper level is low in liver & brain, high in enterocytes & fibroblasts

Clinical features:-

→ Characterized by sparse hypopigmented kinky hairs
→ On Microscopic examination of the hair: Trichorrhexis nodosa (fractures along the hair shaft) and pili torti (twisted hair) seen
→ Growth retardation
→ CNS involvement- Intellectual disability, delayed milestones, seizures, hypotonia
→ Hypothermia
→ Patients with classic form of Menke's disease usually die before 3 years of age.

3. Pellagra like rash is seen in:

a) Gaucher disease
b) Alkaptonuria
c) Hartnup disease
d) Phenylketonuria

Hartnup disorder:

→ Defect in transport of neutral amino acids by intestinal mucosa & renal tubules:
→ Autosomal recessive inheritance
→ Defect in SLC6A19 gene on chr 5p15
→ Most children remain asymptomatic
→ Cutaneous photosensitivity and pellagra like rash.

4. A pediatrics patient present to the hospital with complaints of multiple pathological fractures and the X-ray reveals the below-depicted image. Blood picture presents Low phosphorus and calcium levels with normal Vitamin D levels and elevated PTH. Which of the following is the most probable diagnosis?



a) Osteopetrosis
b) Osteomalacia
c) Osteosclerosis
d) Histiocytosis X

Osteopetrosis/ marble bone disease:

- Bone within bone appearance on X-ray (increased bone density)
- Due to mutation in CLCN7 gene → Leads to defective bone
- C/F: Neurological problems present, Deafness, Macrocephaly, hepatosplenomegaly. Pancytopenia
- Investigation: Low phosphorus and calcium levels with normal Vitamin D levels and elevated PTH.

5. **All of the following are manifestations of congenital syphilis except**:-

a) Gumma
b) Hutchinson's teeth
c) Olympian brow
d) Interstitial keratitis

→ **GUMMA is seen in TERTIARY SYPHILIS.**

Manifestations of early congenital syphilis include:

- Syphilitic rhinitis
- Maculopapular rash
- Condyloma lata

Late signs of Congenital syphilis:

Symptom/sign	Descriptions
Olympian brow	Bony prominence of the forehead caused by persistent or recurrent periostitis.
Higoumenakis sign	Unilateral or bilateral thickening of the sternoclavicular third of the clavicle
Saber shins	Anterior bowing of the midportion of tibia
Hutchinson teeth	Peg-shaped upper central incisors with a notch along the biting surface
Mulberry molars	Abnormal 1st lower molars with excessive number of cusps
Saddle nose	Depression of the nasal root
Rhagades	Linear scars in a spoke-like pattern from mucocutaneous fissures of mouth, genitalia.

www.medicalcoupons.in

Hutchinson triad	Hutchinson teeth, interstitial keratitis, and 8th nerve deafness.
Clutton joint	Unilateral or bilateral painless joint swelling

6. Abnormality involving which chromosome is seen in hunter disease

a) X chromosome
b) Y chromosome
c) Chromosome 21
d) Chromosome 22

All mucopolysaccharidoses are Autosomal Recessive except Hunter's disease which is inherited as X-linked Recessive

Mucopolysaccharidoses:

Type	Name	Enzyme deficien	Clinical Features
I	Hurler	**alpha-L-iduroni das**	Coarse face+ Corneal clouding + MR + HSM + bony changes
II	Hunter	**iduronate-2-sulfatase (I2S)**	Same as above but NO corneal clouding.
IV	Morquio	**N-acetylgalactosamine-6-sulfate sulfatase (GALNS)**	Bony abnormalities severe & corneal clouding may be present

7. Mental retardation is seen in?

a) Phenylketonuria
b) Alkaptonuria
c) Albinism
d) Von Gierke disease

Mental retardation is feature of PKU due to elevated level of phenylalanine hydroxylase in brain.

8. A 6 year child presents with hepatosplenomegaly & generalized lymphadenopathy along with fever The child developed a rash, after being administered ampicillin. What could be the possible diagnosis?

www.medicalcoupons.in

a) Infectious mononucleosis
b) Scarlet fever
c) Kawasaki disease
d) HIV infection

Infectious mononucleosis:

→ Aka glandular fever
→ Etiological agents: Epstein Barr virus (EBV)
→ Mode of transmission: Blood transfusion, Direct contact
→ C/F: Fever, headache, sore throat, generalized lymphadenopathy, splenomegaly, hepatomegaly

Patients with infectious mononucleosis treated with ampicillin or amoxicillin may experience "**Ampicillin rash**" which may occur with other β- lactam antibiotics.

→ This morbilliform, vasculitis "**ampicillin rash**" is probably immune mediated and resolves without specific treatment.

9. In an eye camp doctor's try to make people aware of the causes of blindness. Vitamin A is the leading cause for preventable childhood blindness. A parent brings her 18 month old child to the doctor for routine eye check up. Doctor notices thinned out cornea and from medical records it was seen that the child also has iron deficiency anemia. Weight of the child is 10kg, what should be the dose of vit. A to be given in this case if doctor marks it as keratomalacia.

a) 50,000 IU
b) 1,00,000 IU
c) 2,00,000 IU
d) 5,00,000 IU

Vitamin A DEFICIENCY

It is characterized by night blindness, conjunctival xerosis. Bitot spots, Keratomalacia and fundus changes in severe cases

Treatment:

- <6 months of age Vitamin A 50,000 IU orally/dose
- 6-12 months of age Vitamin A 100,000 IU orally/dose
- >12 months of age Vitamin A 2,00,000 1U orally/dose
- The same dose is repeated next day & weeks later.
- Clouding of cornea in a child with Vit A deficiency is emergency → requires parenteral administration of 50,000 - 100,000 IU Retinol
- In keratomalacia → Local treatment with antibiotic drops & ointment, padding of eye → enhances healing

www.medicalcoupons.in

10. A new born loses maximum heat from:-

a) Head
b) Abdomen
c) Palms & soles
d) Neck

A new born loses maximum heat from 'Head' because of greater surface area.

11. Recommended daily dose of Iodine in child is:-


a) 120-200 µg
b) 90-120 µg
c) 30-60 µg
d) 500 µg

Recommended dietary allowance of iodine:.

- For Infants- 30 µg /kg/day
- For children → 90-120 µg /day
- For adolescents and adults →150 µg /day
- Additional 25 µg & 50 µg during pregnancy & lactation respectively.

PATHO DEC 2018


1. Most characteristic feature of diabetic nephropathy which is show below is:



a) Kimmelstiel Wilson change
b) Armani Ebstein change
c) Focal segmental glomerulosclerosis
d) Membranoproliferative glomerulonephritis

www.medicalcoupons.in

Features of diabetic nephropathy are:



Glomerular lesions like :-

- → Capillary basement membrane thickening → Earliest & most common change in DM
- → Diffuse mesangial sclerosis → presents as nephrotic syndrome in severe cases
- → Nodular glomerulosclerosi aka Kimmelstiel-Wilson lesion → Pathognomonic of diabetes

Renal vascular lesions, principally arteriolosclerosis

Pyelonephritis, including necrotizing papillitis.

2. Xenograft is:

a) Graft across species
b) Graft from same species
c) Graft from same individual
d) Graft from same organ

Autograft	graft taken from one part of the body of an individual and transplanted onto another site in the same individual, e.g., skin graft.
Isograft	graft taken from one individual and placed on another individual of the same genetic constitution, e.g., grafts between identical twins.
Allograft	graft taken from one individual placed on genetically non-identical member of the same species.
Xenograft	graft taken from one individual placed on an individual belonging to another species, e.g., animal to man.

3. Which of the following is most abundant glycoprotein in basement membranes?

a) Fibronectin
b) Laminin
c) Integrin
d) Syndecan

- Laminin is the most abundant glycoprotein in basement membranes.
- Connects cells to underlying ECM components such as type IV collagen and heparan sulfate.

www.medicalcoupons.in

- Laminin can also modulate cell proliferation, differentiation, and motility.

4. Which of the following is true about Anaplasia?

a) Loss of cohesion between cells
b) Loss of differentiation
c) Change of epithelium type
d) Benign and fully reversible

Lack of cellular differentiation or Anaplasia is considered a hallmark of malignant transformation. The term anaplasia means "to **form backward**," which implies dedifferentiation (or loss of the structural and functional differentiation) of normal cells during tumorigenesis.

5. Cell that can form the maximum number of other cells in the body is called?

a) Totipotent
b) Multipotent
c) Pluripotent
d) Lineage stem cells

Totipotent cells- can form maximum no of other cells in body

e.g. cells of zygote [MCQ]

Note: Totipotent cells give rise to teratoma.

6. Which of the following disease has X linked recessive pattern of inheritance?

a) Hurler Syndrome
b) Thalassemia
c) Hereditary spherocytosis
d) G6PD deficiency

X-LINKED RECESSIVE DISORDERS: Oblivious **F**emale **W**ill **G**ive **H**er **B**oys **H**er X-**L**inked **D**isorders

- → **O**cular albinism
- → **F**abry disease
- → **W**iskott-Aldrich syndrome
- → **G**6PD deficiency
- → **H**unter syndrome
- → **B**ruton agammaglobulinemia
- → **H**emophilia A & B

→ **L**esch - Nyhan syndrome
→ **D**uchenne muscular dystrophy

7. Which patter of inheritance of disease is associated with consanguinity?

a) Autosomal recessive
b) Autosomal dominant
c) X - linked recessive
d) X - linked dominant

- CONSANGUINITY condition in which closely related individuals are going to have the marriage.
- This can lead to expression of even those genes which are recessive in nature.
- Therefore, it is a risk factor for expression of Autosomal Recessive disorder.

8. Which organ is involved in Goodpasture syndrome?

a) Liver
b) Adrenals
c) Kidney
d) Brain

Good pasture syndrome

- Type II Hypersensitivity reaction.
- Auto Ab against glomerular basement membrane type IV collagen.
- **Affects kidneys & lung causing hematuria & hemoptysis.**
- Characterized by linear deposits.

9. A 57-year-old male came with complaints of swelling over the scalp region which gradually developed over 6 months. Now complaints of pus discharge several times in the past month associated with pain. On examination, swelling sized 3x2 cm, Firm, Fixed; Tenderness(+), Warmth(+), Punctum(+), Discharge(+). Incision & Drainage was done, the wound was left open to heal. Which of the following is a mechanism for healing in this case?

a) Granuloma formation
b) Scab formation
c) Granulation tissue
d) Neovascularization

- Healing intention: Wounds by secondary are left open and allowed to heal by granulation, contraction and epithelialization.
- In healing by second intention, large amounts tissue are formed of granulation to fill a bigger gap caused by the larger area of deficit.

10. A 47 years old male, active smoker with a known case of Bronchial Asthma now presents with difficulty in breathing with the breath sounds being audible externally. Which of the following is not a feature you will expect to find in this patient?

a) Thickening of bronchial wall
b) Increase in number of airway goblet cells
c) Fibroblastic foci
d) Sub-basement membrane fibrosis

Fibroblastic foci (exuberant proliferation of fibroblasts) is a feature of idiopathic pulmonary fibrosis (IPF). Bronchial asthma is an example of Type I Hypersensitivity reaction and is characterized by airway remodeling which includes:

- Thickening of bronchial wall with hypertrophy and/or hyperplasia of bronchial smooth muscles
- Increase in size of submucosal glands and number of goblet cells
- Increased vascularity
- Sub-basement membrane fibrosis (due to deposition of type I and III collagens)

Key points - Sputum findings in bronchial asthma:

- Curshmann spirals - impacted mucin plugs.
- Charcot Leyden crystals - Eosinophil crushing artefact.
- Creola bodies - damaged respiratory epithelial cells.

Mneumonic	Type 2 HSR is seen in following condition:
My :	Myasthenia gravis
Blood :	Blood transfusion (mismatch)
Group :	Graves, good pasture
Is:	Immune hemolytic anemia
[illegible] :	Rheumatic heart disease
Positive :	Pernicious anemia/ pemphigous vulgaris

11. Acute graft rejection occurs within?

a) Seconds to minutes
b) Days to weeks
c) Weeks to months
d) Months to years

Hyperacute rejection: type 2 HSR, happens over span of few minutes to hours. Eg: kidney transplant

Acute rejection: Few weeks to months(< 6 months)

Chronic rejection: After 6 months.

www.medicalcoupons.in

Graft vs host disease (Runt Disease):

→ Acute GVH: < 100 days
→ Chronic GVH: > 100 days

12. All are reduced in iron deficiency anemia EXCEPT:

a) Total iron binding capacity
b) Percentage saturation of Transferrin
c) Hemoglobin level
d) Serum ferritin

Iron deficiency typically causes low serum iron, high iron-binding capacity, and **low serum ferritin levels.** Anemia occurs when you have a decreased level of **hemoglobin** in your red blood cells (RBCs).

13. Pheochromocytoma produces all except?

a) Nor-epinephrine
b) Secretin
c) Vasoactive intestinal polypeptide
d) Somatostatin

Pheochromocytoma - Tumor of adrenal medulla

Follows rule of 10 -

10% are extra adrenal

10% are bilateral

10% are children

10% are w/o hypertension

Tx: Surgical Resection

c/s β blocker

www.medicalcoupons.in

SURGERY DEC 2018

1. Most common immediate complication of splenctomy:

a) Hemorrhage
b) Fistula
c) Bleeing from gastric mucosa
d) Pancreatitis

Complications of splenectomy:

➢ **Early:**
- Haemorrhage
- Injury to stomach, splenic flexure, pancreas

➢ **Delayed:**
- Fistula from stomach, pancreas
- Subdiaphragmatic collection
- Left basal atelectasis and pleural effusion
- Thrombocytosis- thrombotic complications

2. All are premalignant conditions of oral cavity except:

a) Chronic hyperplastic canidiasis
b) Oral submucosal fibrosis
c) Oral lichen planus
d) Leucoplakia

All the given options increase the risk of carcinoma of oral cavity but, oral Lichen planus has the least risk as it is Low-risk or Equivocal lesion.

3. A 6 year old child is brought with high fever with rigors for 5 days with pain examination patient s anicteric and tenderness is noted in right upper quadrant. What is best investigation for this case?

a) USG
b) Serology
c) SGOT/LFT
d) CECT

- The patient is having pain in the right hypochondrium which suggests the involvement of the liver.
- An abscess can be suspected as the patient is suffering from fever accompanied with rigors & pain and tenderness in the right upper outer quadrant

www.medicalcoupons.in

- Abscess be of two types:
  → Pyogenic liver abscess
  → Amoebic liver abscess
- On examination the patient is anicteric. So, the patient could be suffering from Amoebic liver abscess.
- In developing countries like India, Amoebic liver abscess is more common because of the feco-oral and poor sanitation barrier
- Initial investigation for the diagnosis of amoebic liver abscess can be done by USG and CT-Scan.
- **Investigation of choice or the best investigation for this patient is Serology i.e. ELISA which detects the anti-bodies formed by the parasite Entamoebahistolytica.**
- The amoebic liver abscess can be treated by metronidazole empirically.

4. Comment the diagnosis of this 60 years of man:



a) Basal cell carcinoma
b) Plunging Ranula
c) Epulis
d) SCC of lip

- Ulcero proliferative growth or Cauliflower like growth on the lower lip suggestive of squamous cell carcinoma of the lower lip
- In total 95%of carcinomas of lip arise the lower lip and 5% arise in central one-third and commissures.
- Aetiological feature include UV radiation exposure.
- Tumors tend to spread laterally over the mucosal surface.
- Lymph node metastases, usually to submental or submandibular nodes which reflect aggressive disease behavior then approach.

www.medicalcoupons.in

5. Best treatment strategy for carcinoma anal canal:

a) Chemoradiation
b) Surgery
c) Radiation
d) Chemotherapy

- Nowadays the primary treatment of choice is chemoradiotherapy (Combined modality therapy (CMT) Nigro Regimen), which usually includes 5-FU, Mitomycin-C and cisplatin
- Radical surgery is indicated in those with persistent or recurrent disease following CMT or Nigro Regimen.

6. 32 year old female presented to the OPD with complaints of abdominal pain, diarrhea and weight loss. On further evaluation patient was found to have multiple strictures of small bowel with interspersed areas of normal mucosa in between. Which of the following is false regarding the given condition?

a) No recurrence after surgery
b) Aphthous ulcer
c) Skip lesions
d) Fistula formation

Recurrences occur after surgery in Crohn's disease. Fistula formation, skip lesions and aphthous ulcers all seen in Crohn's disease.

7. Most common subtype of Rodent ulcer is:

a) Superficial
b) Cystic
c) Nodular
d) Pigmented

- Rodent ulcer is also known as basal cell carcinoma of the skin, or tear cancer which is a slow growing type of cancer.

Types of Rodent ulcers:

→ Nodular (90%) (MC)
→ Pigmented
→ Cystic
→ Superficial

- The most common type being nodular which has characteristic pearly white nodules in the periphery with central umbilication and telangiectatic vessels.

www.medicalcoupons.in

8. Which of these is the best for management of a 3 cm stone in renal pelvis without evidence of hydronephrosis?

a) ESWL
b) PCNL
c) Antegrae pyeloplasty
d) Retrograde pyeloplasty

- Generally, ESWL is a non-invasive method used for removal of renal calculi but the maximum size of stone that can be removed by ESWL is up to 1.5 cm
- If the size of the stone is more than 2 cm then the preferred treatment option is **Percutaneous nephrolithotomy** (PCNL)
- Other contraindications to ESWL are obese patient, pregnancy, & patients taking oral anti-coagulants.

9. AFP is a tumour marker for which of the following?

a) HCC
b) RCC
c) Oncocytoma
d) Chordoma

Markers of Hepatocellular carcinoma are:

AFP (α -fetoprotein)

PIVKA-2 (Protein induced by vitamin K absence 2)

10. Germ cell tumour not seen in males:

a) Choriocarcinoma
b) Seminoma
c) Sertoli cell tumour
d) Teratoma

WHO classification of testicular tumor:

Germ cell tumors:-

→ Seminoma
→ Non-Seminomatous GCT
→ Yolk sac tumor
→ Teratoma
→ Embryonal carcinoma
→ Choriocarcinoma

www.medicalcoupons.in

11. Dye for Senitnel Lymph Node Biopsy is injected in which of the following sites?

a) Axilla
b) Tail of spence
c) Nipple
d) Areola

- Sentinel lymph node biopsy (SLNB) is done by two techniques:

1. Blue dye technique - 1% lymphazurin (ISO sulfan blue)

2.Radioactive colloid technique-technetium^{99} labelled Sulphur colloid.

- Maximum accuracy is seen with both techniques combined
- **A radioactive dye is injected around the areola subcutaneously in the subdermal plexus.**

12. Correct description abut Paget's disease of breast:

a) Eczema of skin of nipple
b) Eczema of skin of areola
c) Mastitis carcinomatosis
d) Atrophic scirrhous carcinoma

- Paget's disease of the nipple is a superficial manifestation of an underlying breast carcinoma.
- **It presents as eczema-like condition initially around the nipple** and then spreads to areola.
- In Paget's disease of nipple underlying manifestations are:
  → Ductal carcinoma in situ (DCIS).
  → Invasive ductal carcinoma.
- Paget's is not associated with lobular breast cancer.
- It is Estrogen and Progesterone Receptor negative.



www.medicalcoupons.in

13. Cut off duration for diagnosis of Priapism is:

a) 1 hour
b) 2 hours
c) 3 hours
d) 4 hours

Priapism: It is defined as a persistent painful erection lasting longer than 4 hours irrespective of any sexual stimulation & beyond reaching of orgasm.

It is an emergency situation

It is of two types:

1.Ischemic priapism

2.Non-ischemic priapism


14. The image given below indicates which of the following procedures?



a) Percutaneous transhepatic cholangiogram
b) T- tube cholangiogram
c) ERCP
d) HIDA scan

Endoscopic retrograde cholangiopancreatography, or ERCP, is a **procedure to diagnose and treat problems in the liver, gallbladder, bile ducts, and pancreas**. It combines X-ray and the use of an endoscope—a long, flexible, lighted tube.

www.medicalcoupons.in



15. Which of the following is not done in carcinoma esophagus?

a) Biopsy
b) pH - metry/monitoring
c) CT chest
d) PET scan

INVESTIGATIONS of Carcinoma Esophagus :

- 1st investigation → Barium Swallow
- IOC for Dx → Endoscopy with biopsy
- IOC for staging → EUS
- IOC for staging in cases of obstructed lumen/ advanced stage: CECT
- IOC for distant metastasis → PET Scan

16. Bell clapper testis predisposes to:

a) Torsion testis
b) Varicocele
c) Cancer of testis
d) Hydrocele

A bell clapper deformity is a predisposing factor in testicular torsion in which the tunica vaginalis has an abnormally high attachment to the spermatic cord, leaving the testis free to rotate. Bell clapper deformity predisposes to **intravaginal torsion of the testis**.

www.medicalcoupons.in

17. A 29 yr old man with a history of varicose veins since the age of 12 presented with c/o pain, heaviness, and swelling on the right leg. The patient underwent below-knee stripping of the Long saphenous vein after which the symptoms completely resolved along with visible varicose veins. The most common complication that can occur to this patient is:

a) Hemorrhage
b) Thromboembolism
c) Neuralgia
d) Infection

- Ideally, the long saphenous vein should not be stripped below the knee as it causes injury to the saphenous nerve which is the most common iatrogenic complication.
- The incidence of saphenous nerve neuralgia up to 7 % following long saphenous vein stripping to the knee.
- And the, most commonly injured nerve while stripping the short saphenous vein is sural nerve whose incidence of injury may be as high as 20%.

18. Most common benign breast tumour:

a) Fibroadenoma
b) Fibroadenosis
c) DCIS
d) Phyllodes tumour

- **Fibroadenoma is the most common benign breast tumor which is most commonly seen in young female .**
- Fibroadenoma is also known as "Breast Mouse" due to its characteristic high mobility.
- On the other hand fibro adenosis is a benign breast diseaseso it is excluded
- On Mammography (if performed on older female) there is popcorn calcification.
- The name DCIS(Ductal carcinoma in-situ) suggests its malignancy and hence can be excluded.
- Phylloides tumor or Cystosarcoma Phylloides is a malignant tumor.

19. Sebaceous cyst is not seen in:

a) Back
b) Spleen
c) Scalp
d) Scrotum

Sebaceous cyst arises from sebaceous glands, **commonly related to hair follicles**

They usually occur on the scalp, face, ear, back & scrotum

A Sebaceous cyst consists of a punctum usually which discharges on pressing

www.medicalcoupons.in

The punctum is absent in the sebaceous cyst o1 scalp & scrotum.

20. Which of the following is not a Hospital-Acquired infection?

a) Surgical site infection
b) STD
c) UTI
d) Pneumonia

Infection acquired from the environment or the staff following surgery or after admission to the hospital is termed as Hospital-acquired infection (HAI) which is also referred to as Nosocomial infections.

There are four main groups of HAI's

1. Respiratory infections → including Ventilator-associated pneumonia (10-15%)

2. Bacteremia → related to indwelling vascular catheters(10-15%)

3. Urinary tract infections (UTI) Related to urinary catheters(30-40%-MC)

4. Surgical site infection (SSI) (2ndMC)

21. Amount of blood loss in class III circulatory failure/Haemorrhagic shock?

a) <15%
b) 15-30%
c) 30-40%
d) >40%

Class 3: Volume loss from 30% to 40% of total blood volume, from 1500 mL to 2000 mL. A significant drop in blood pressure and changes in mental status occurs. Heart rate and respiratory rate are significantly elevated (more than 120 BPM). Urine output declines. Capillary refill is delayed.

- **Class 1:** Volume loss up to 15% of total blood volume, approximately 750 mL. Heart rate is minimally elevated or normal. Typically, there is no change in blood pressure, pulse pressure, or respiratory rate.
- **Class 2:** Volume loss from 15% to 30% of total blood volume, from 750 mL to 1500 mL. Heart rate and respiratory rate become elevated (100 BPM to 120 BPM, 20 RR to 24 RR). Pulse pressure begins to narrow, but systolic blood pressure may be unchanged to slightly decreased.
- **Class 4:** Volume loss over 40% of total blood volume. Hypotension with narrow pulse pressure (less than 25 mmHg). Tachycardia becomes more

www.medicalcoupons.in

- pronounced (more than 120 BPM), and mental status becomes increasingly altered. Urine output is minimal or absent. Capillary refill is delayed.

22. Comment on the diagnosis of al film shown of a 65-year -old man with acute abdomen:



a) Ileocolic intussusception
b) Sigmoid volvulus
c) Toxic megacolon
d) **Colocolic intussusception**

Notice the significant Clawsign in the given film of a Ba. Enema which is suggestive of intussusception.

INTUSSUSCEPTION

M/C type → "ileocolic"

M/C type in adults → Colocolic

Since in the question the age of the patient was clearly mentioned as 65 years old. The diagnosis will be colocolic intussusception.

23. Breast triple assessment contains all except:

a) Clinical examination
b) Axillary sampling
c) USG
d) FNAC and biopsy

www.medicalcoupons.in

Triple assessment, as the name indicates, includes three modalities, **physical examination, imaging (mammography and/or ultrasound)**, and biopsy (FNAC and core biopsy).

24. Identify the lesion shown below:



a) Wet gangrene
b) Dry gengrene
c) Frost bite
d) Ainhum

From the given image we can figure out the lesion is a gangrene due to the presence of black necrotic areas.

Feature	Dry gangrene	Wet gangrene
Site	Commonly limbs	More common in bowel
Mechanism	Arterial occlusion	Venous obstruction
Macroscopy	Dry, shrunken & black	Moist, soft, swollen, rotten and dark
Putrefaction	Limited due to less blood supply	Marked due to congestion of organ with blood
Line of demarcation	Present	No clear-cut line
Bacteria	Bacteria fail to survive	Numerous bacteria present
Prognosis	Better due to septicemia	Poor due to profound toxemia

www.medicalcoupons.in

25. 30 year old male presents to the OPD with complaints of lower abdominal pain and bleeding per rectum. On further evaluation , patient was found to have multiple superficial ulcers in the rectum and extending upward to involve entire colon. On histology, only mucosa and submucosa were found to be involved. Surgery is indicated for the above disease in all of the following conditions except:

a) Toxic megacolon
b) Colonic polyp
c) Dysplasia
d) Massive colonic bleeding

The given clinical condition is ulcerative colitis

Indication for surgery in a patient of Ulcerative colitis:

- Intractability → refractory to medical management
- Dysplasia
- Carcinoma
- Massive colonic bleeding
- Toxic megacolon

26. Patient complains of painless swelling over the face with difficulty in swallowing. The appearance of face is shown. The probable diagnosis is:



a) Acute parotitis
b) Cancer of parotid gland
c) Angioedema of face
d) Acute sialadenitis

- Painless swelling usually points towards malignancy.
- Among the given options Acute parotitis and Acute sialadenitis can be excluded because they both are painful condition.
- Angioedema of the face is characterized by the presence of swelling around the lips and around the oral cavity.
- Hence the answer is cancer of the parotid gland

www.medicalcoupons.in

- Among malignant tumors of the parotid gland, mucoepidermoid carcinoma is the most common followed by adenocystic carcinoma.

27. A 35-year-old lady the presented with a 6-month painless fluctuant, non-transilluminant swelling with a thin watery discharge. Clinical diagnosis is?



a) Brachial cyst
b) Secondaries
c) TB
d) Lymphoma

Among the given options discharge can only be seen with the TB which has characteristic undermined edges which can be identified easily from the given image.

The branchial fistula may, however have discharge but around the anterior border of SCM and doesn't have undermined edges.

Lymphoma has a lump, anorexia, weight loss but no discharge

28. The following patient has presented after chest trauma. On examination crepitus is felt. The clinical diagnosis is?



a) Subcutaneous emphysema
b) Gas gangrene
c) Acute tubular necrosis
d) SIADH syndrome

www.medicalcoupons.in

- **From the given image, you can notice that the patient has puffy eyes and swelling around the face and crepitus is felt when palpated.**
- **This type of appearance is usually noticed in Sub-cutaneous emphysema**
- It is advised not to confuse between gas gangrene and subcutaneous emphysema as both the conditions have crepitus.
- Gas gangrene is a condition that is caused by Cl. Perfiringens which has a gangrenous type of appearance with crepitus due to clostridial myonecrosis and takes more than 24 hours to appear.

MEDICINE DEC 2018

1. Which of the following is not done in W.P.W?

a) Treadmill test
b) Electrophysiological studies
c) Oral beta blocker
d) Procainamide

C/I of treadmill test:

1) severe Aortic Stenosis

2) DVT

3) Charcot's joint

4) WPW

5) MI in first 48 hours

6) Unstable angina during attack

7) Buerger's disease

8) Bilateral osteoarthritis of knee

2. What is the impact on fetus in case of use of Indomethacin in utero in third trimester?

a) Patent ductus arteriosus
b) Early closure of ductus arteriosus
c) Ventricular septal defect
d) Atrial septal defect

Prostaglandins helps to keep ductus arteriosus open.

www.medicalcoupons.in

Indomethacin inhibits prostaglandins levels → So, closure of ductus arteriosus → DOC for PDA

3. In fever of unknown origin, blood sample should be drawn how many times?

a) 2
b) 3
c) 4
d) 5

Criteria for fever of unknown origin-

1. Fever 101.30 F on 2 separate occasions

2. For > 3 weeks

3. Absence of immunocompromised state

4. **Adequate workup-**

→ Blood culture > 3 times should be normal
→ Urine analysis- sterile
→ CRP normal
→ Antinuclear antibody negative
→ LDH normal

4. Which of the following is correct about cancer developing in Ulcerative colitis?

a) Bad prognosis if it involved only rectum
b) Good prognosis if it involves only rectum and is low grade
c) 2% chance of malignancy if present for 10 years
d) Young age onset has low chances as compared to old age onset

- Cancer developing in ulcerative colitis is directly proportional to duration and extent of disease
- More the duration and extent more chances of cancer to develop
- Incidence of malignancy is 2% if disease persist for 10 years
- Incidence of malignancy is 18% if disease persist for 30 years

5. Refractory Septic shock is defined as?

a) Shock persisting > 1 hour after IVF administration
b) Shock persisting > 1 hour after IVF administration and high dose vasopressors
c) Shock that is refractory to medical treatment
d) Shock that does not improve despite treatment for 24 hours

www.medicalcoupons.in

- **Refractory Septic Shock :**

Septic shock that last for 1 hour and does not respond to fluid or pressor administration.

- **Septic Shock :**

Sepsis with hypotension despite adequate fluid resuscitation for at least 1 hr with perfusion abnormalities. or Need of vasopressors to maintain the blood pressure

6. Waist to hip ratio that increases risk of heart disease is?

a) 0.80 in male
b) 0.80 in female
c) 0.85 in male
d) 0.85 in female

Waist to hip ratio that increases risk of heart disease for male is > 1

Waist to hip ratio that increases risk of heart disease for female is > 0.85

7. A 65-year-old, male with hypertension develops sudden onset weakness on the right side of the face & arm. His attender recalls history of headache, blurred vision in the patient which leads to the suspicion of increased ICP. Which is the most common cranial nerve involved in raised ICP?

a) Abducens nerve
b) Trochlear nerve
c) Trigeminal nerve
d) Facial nerve

- Abducens= Longest subarachnoid course (so whenever ICP will raise abducens nerve will affect first)
- Trochlear= Longest intracranial route, thinnest, only nerve has dorsal origin
- Trigemin[illegible]=Thickest nerve
- Facial= Longest intraosseous route

8. After trauma a patient is unable to smell coffee and asafetida but can smell ammonia. Which of the following statement is true about the patient?

a) Ammonia is used for testing olfactory nerve
b) The olfaction system is not damaged
c) Ammonia activates trigeminal nerve
d) All of the above

www.medicalcoupons.in

Ammonia cannot be used as test to check olfactory because it is an irritant not smell that stimulates trigeminal nerve not olfactory.

9. Deepika a 26 years old female , labourer by occupation presented to you with complain of pain in abdomen, nausea and restlessness since 4 hours. She gives h/o 2 episodes of hematuria. USG showed a 3*3 cm stone in Right renal pelvis. There is no evidence of hydronephrosis . Which of following is the best procedure for management of this patient ?

a) ESWL
b) PCNL
c) Antegrade Pyeloplasty
d) Retrograde pyeloplasty

- ESWL- done for stone less than 2.5 cm size
- **PCNL-if stone is > 2.5 cm will treat by PCNL**
- Most common kidney stone → calcium oxalate
- Etiology=idiopathic hypercalciuria

10. A new vaccine which was launched in 2018 in Thailand with the name CYD-TDV. In which of the following condition it is used?

a) Dengue
b) Malaria
c) Yellow fever
d) Japanese encephalitis

- **CYD-TDV vaccine introduced in Thailand in 2018 (DENGVAXIA) is used for Dengue.**
- Vaccine is tetravalent
- Live attenuated vaccine
- Dengue is a mosquito (Aedes aegypti) borne viral (Arbovirus) infection.

11. A 45 year old male who had a history of domestic animal bite has come for follow up post 90 days after receiving rabies post-exposure prophylaxis. Now what will you use for Rapid diagnosis of rabies in this case ?

a) Skin biopsy with fluorescent antibody testing
b) Rabies virus specific antibodies
c) Inoculation in mouse
d) Corneal impression smear

www.medicalcoupons.in

- Fluorescent Antibody Test : This is a highly reliable and the best single test currently available for the Rapid diagnosis of Rabies viral Antigen in Infected specimen.
- Corneal impression & Mouse inoculations are also used for diagnosis.

12. Common neurological tumour in NF-2?

a) Acoustic neuroma
b) Optic glioma
c) Café-Au-Lait macules
d) Meningioma

ACOUSTIC NEUROMA: It is an autosomal dominant condition it is an eighth nerve tumor. It compresses adjacent cranial nerves therefore multiple palsies develop.

OPTIC GLIOMA: occurs in NF1

NF 1 Characterized by presence of cafe au lait macule or small potato like Lesion all over the Skin.

13. A 7-year-old with fever for days with pain in right hypochondrium. On examination tenderness is noted in Right upper quadrant. What is the next best investigation?

a) USG
b) Serology
c) SGOT/LFT
d) CT abdomen

The next best investigation in this case is to do an USG.

Causes of Right upper quadrant pain in pediatric-

1. Acute viral hepatitis - **in viral hepatitis on USG= starry sky appearance seen*****

2. Acalculous cholecystitis

3. Budd Chiari syndrome

4. Acute heart failure

14. Which of the following is not seen in Horner syndrome?

a) Mydriasis
b) Ptosis
c) Anhidrosis
d) Enophthalmos

www.medicalcoupons.in

Horner syndrome is characterized by damage to sympathetic chain

Features are – "**SAMPLE**"

Sympathetic Nerve Fiber Injury

Anhidrosis

Miosis

Ptosis

Ioss of ciliospinal reflex

Enophthalmos

15. Which of the following is not correct about Defibrillation?

a) Easy for untrained person
b) Decrease in success rate with delay in initiation
c) Improve prognosis
d) minute gap between 2 shocks

Yes, Defibrillation can be given by non-medical also

Basic life support protocol (BLS protocol) = 30:2 in 2-minute gap

16. The following image shows?



a) Achalasia cardia
b) Diffuse oesophageal spasm
c) Carcinoma Oesophagus spasm
d) Schatzki's Ring

Diffuse esophageal spasm - Cork screw appearance

Achalasia cardia - Bird peak appearance

Carcinoma esophagus - Apple core appearance.

17. Which of the following features are true about Allergic Broncho Pulmonary Aspergilloma?

1)Distal bronchiectasis

2)Serum precipitins to Aspergillus

3) Increased IgE Levels

4)Seen in asthmatics

a) 1,2,3 only
b) 2,3,4 only
c) 1,3,4 only
d) All of the above

18. Mark the true statements -

1. MEN 2B is also known as Sipple syndrome

2. MEN 1 is also known as Werner syndrome

3. Marfanoid Habitus is a feature of MEN 2B

4. MEN 4 is associated with CDKN1B

a) 1 and 2 are true
b) 3 and 4 are true
c) 1,2 and 4 are true
d) 2,3 and 4 are true

MEN 2A - SIPPLE Syndrome

MEN1 was originally known as **Wermer syndrome**

19. Which of the following is not a component of child pugh Classification

a) **SGOT**
b) Bilirubin
c) Albumin
d) Prothrombin time

www.medicalcoupons.in

The Child-Pugh (CP) classification is the standard to assess preoperative liver function and is determined by five factors: **serum bilirubin and albumin levels, prothrombin time, ascites, and encephalopathy.**

20. Fulminant diabetes mellitus is seen in?

a) Diabetic ketoacidosis
b) Coxsackie B virus
c) Non ketonic hyperosmolar coma
d) Autoimmune pancreatitis

- **Fulminant diabetes mellitus - caused by Coxsackie B virus (viral myocarditis)**
- **Coxsackie B virus:**
  → Herpangina
  → Hand foot mouth virus.

21. Which is correct about rheumatoid nodules?

Tender, located on extensor surface and seen with arthritis

Non tender, located on extensor surface and seen with arthritis

Non tender, located on flexor surface and seen with arthritis
Tender, located on flexor surface and seen with arthritis

- Seen in extensors such as olecranon process, occiput, tips of spinous process of vertebrae.
- Rheumatoid nodules are MC extra articular manifestation of Rheumatoid arthritis

22. Modified duke criteria are used for diagnosis of?

a) Infective endarteritis
b) Infectious mononucleosis
c) Inflammatory myopathy
d) Infective endocarditis

- **The Modified Duke Criteria for the Clinical Diagnosis of Infective Endocarditis**
- Duke's staging - Cancer of rectum
- Duke's score - Chronic stable angina

www.medicalcoupons.in

23. Which of the following shall be seen with use of a small size BP cuff?

a) False elevation of BP
b) Falsely low value of BP
c) Cancels the effects of calcified arteries
d) Increases trans arm impendance

- **SMALL SIZED CUFF**: It leads to false elevation of BP as Blood vessels are squeezed so there is an increase in the peripheral resistance
- **BIG SIZED CUFF**: It leads to falsely low measure of BP

24. Which is correct about larval stage of taenia solium?

a) Cysticercosis cellulose
b) Larva currens
c) larva migrans
d) Visceral larva migrans

Cysticercosis is a common helminthic infection caused by the larval stage of pork tapeworm, Taenia solium. **The larvae, also called as cysticercosis cellulose**, reside within the muscles and other tissues of pigs that act as their intermediate hosts.

25. On putting an internal jugular vein catheter, a patient has developed sudden onset severe respiratory distress. Clinical diagnosis is?

a) Pneumothorax
b) Sepsis
c) ARDS
d) Infective endarteritis

The catheter penetrated the jugular vein and traumatized the lung parenchyma damaging the lung tissues leading to PNEUMOTHORAX.

26. You are an intern, posted in the neurology ward in the Department of Internal Medicine. A 48-year-old woman was admitted with a history of muscle weakness. Investigations & examination revealed normal nerve conduction but flaccidity is seen. Which of the following would be the possible diagnosis?

a) Myasthenia gravis
b) GBS
c) Transverse myelitis
d) Traumatic neuritis

www.medicalcoupons.in

Myasthenia Gravis- Problem in Neuromuscular junction due to improper functioning of Ach receptor, so nerves are not damaged. neuromuscular junction is affected leading to muscle weakness.

In option “b,c,d” the nerve conduction is always abnormal .

27. Which of the following is the most sensitive test for the diagnosis of Organophosphate poisoing ?

a) Plasma acetylcholinesterase
b) Organophosphate level in blood
c) Organophosphate level in plasma
d) RBC transaminase levels

The most commonly used test to confirm acute organophosphate poisoning is **measurement of plasma cholinesterase activity**. Plasma cholinesterase levels usually decline to less than 50% of the normal value before any symptoms of poisoning are observed.

28. Extensor Reflex on pinching gastrocnemius muscle is called?

a) Gower Sign
b) Homan Sign
c) Oppenheim Sign
d) Gordon Sign

On squeezing or pinching the calf muscles (gastrocnemius), the Big toe goes away - GORDON SIGN

29. Reversible dementia is seen in?

a) SACD
b) AD
c) CJD
d) Pick's disease

Vit B12 deficiency Reversible dementia is seen


SACD Subacute combined demyelination of spinal cord.

www.medicalcoupons.in

Reversible Dementia	Irreversible Dementia
➢ Hypothyroidism	➢ Alzheimer disease (MC cause of senile dementia
➢ Normal pressure Hydrocephalus	➢ CJD (Creutzfeldt-iakob disease) ○ Caused by Prion Disease ○ Prion mediated damage.
➢ Whipple Disease (Bacteria: T. Whippli)	➢ Pick's Disease (Frontal part of brain is affected)
➢ Vit B12 ↓	

RADIO DEC 201

1. Identify the pathology in the shown CT:



a) Intraventricular bleed
b) Massive Epidural Hemorrhage
c) Subdural Hemorrhage
d) Subarachnoid Hemorrhage

Extradural Hemorrhage (EDH):

- Lentiform shaped, Biconvex , Lemon shape
- Middle meningeal artery
- Anterion

www.medicalcoupons.in

2. A 45-year-old female presents with a complaint of abdominal pain and fever for the past 3 days. She localizes the pain to her epigastric area and states that it radiates to her right upper quadrant. . She notes that it became markedly worse after eating dinner last night. Her past medical history is significant for hypertension and hypercholesterolemia. On Abdominal examination, Inspiratory arrest is noted on deep palpation of the right hypochondrium. Which of the following is the Investigation of choice for this condition?

a) H.I.D.A Scan
b) USG
c) CT Scan
d) Biopsy

- IOC for Acute Cholecystitis: USG
- USG is the IOC for most GB pathologies
- Most Accurate Investigation for Acute Cholecystitis: H.I.D.A Scan (H.I.D.A is used for equivocal cases where USG is not conclusive)
- IOC for Abdominal trauma :CECT


3. Which of the following investigations work on the same principle?

a) CT and MRI
b) CT and X-ray
c) USG and HIDA Scan
d) MRI and PET Scan

Principle of:

- **CT** : Linear attenuation of X-rays
- **X-ray**: X-rays (Fluoroscopy also works on the principal of X rays)
- **MRI**: Gyromagnetic property of H+
- **USG:** Pulse Echo Principle with Piezoelectric effect
- **PET scan:** Annihilation and coincidence imagine, gamma rays -511 key
- **HIDA Scan**: Radioisotope scintigraphy, gamma rays.

4. Comment on the diagnosis of the image shown:



a) Pneumoperitoneum
b) Pneumothorax
c) Eventration of diaphrgram
d) Pneumomediastinum

www.medicalcoupons.in

Pneumoperitoneum - "Air below Diaphragm" is the characteristic feature

ANAESTHESIA DEC 2018

1. **Preoperative assessment of oral cavity for endotracheal intubation in an adult is done by:**

a) Mallampati score
b) Cormack lehane
c) ASA grading
d) AHA grading

→ **Mallampati score** - preanesthetic evaluation of airway in which we are evaluating size of tongue with respect to oral cavity.
→ Cormack lehane - laryngoscopic view of glottic opening.
→ ASA grading - preanesthetic evaluation by which we assess risk associated with surgery.
→ AHA grading- done in cardiac patients.

2. **Which of the following Anaesthesia agent is hepatotoxic:**

a) Sevoflurane
b) Isoflurane
c) Halothane
d) Procaine

Drugs causing Hepatotoxicity

ANAESTHETIC AGENTS :

→ Halothane
→ Chloroform
→ Isoflurane, Enflurane & Desflurane
→ Nitrous oxide.

3. **Anaesthesia given to dog:**

a) Ketamine
b) Medetomidine hydrochloride
c) Midazolam
d) All of the above

Any I/M injectable drug which we are using in human can be used in canines as well.

www.medicalcoupons.in

4. Shortest action local anaesthetic:

a) Procaine
b) Cocaine
c) Dibucaine
d) Chloroprocaine

Shortest Acting LA → Chloroprocaine

Longest acting LA → Dibucaine

5. During laryngoscopy, best method for checking whether the intubation is correct or not:

a) Capnography
b) Air entry on auscultation
c) Ultrasonography
d) Chest movement

- Best method to check correct placement of endotracheal tube (ETT) is capnograph
- Put the tube, inflate the cuff, attach a device (capnograph) if tube has Co2, the graphical representation of Co2 comes in the monitor then we confirm tube is rightly placed.

ORTHO DEC 2018

1. Blount Disease is involvement of

a) Proximal tibia
b) Distal tibia
c) Proximal femur
d) Distal femur





Blount's disease is a progressive bow-leg deformity associated with abnormal growth of posterior-medial part of proximal tibia.

www.medicalcoupons.in

2. Involvement of DIP PIP and Sparing of MCP and wrist is seen in:

a) Osteoarthritis
b) Rheumatoid arthritis
c) Psoriasis
d) Hyperparathyroidism

- Sparing of MCP is a feature of OA
- DIP is spared in RA
- In psoriasis all joints are involved but most specifically DIP is involved
- In Hyperparathyroidism all the bones are involved.

3. Tennis player can spontaneously reduce a shoulder dislocation. He can do it again and again himself. He is suffering from?

a) Acute shoulder dislocation
b) Recurrent shoulder dislocation
c) Inferior shoulder dislocation
d) Fracture upper end humerus

- Most common shoulder dislocation - Anterior type of shoulder dislocation which accounts for 95%. Since it was mentioned that the player can reduce the dislocation "again and again", it could be a habitual type of dislocation which is recurrent.
- Acute shoulder dislocations are very painful which cannot be reduced or relocated again and again.
- Inferior type of shoulder dislocation usually presents with a characteristic "salute position" hence can be excluded.

4. Adamantinoma affects which of the following ?

a) Radius
b) Humerus
c) Femur
d) Tibia

- Epithelial tumors are classified into Ameloblastoma and adamantinoma, though both the tumors show same type of histopathology, they are not interchangeable.

www.medicalcoupons.in

	Ameloblastoma	Adamantinoma
Site	Mandible	Tibia

- Most common site for Ameloblastoma is mandible
- Most common type of cancer of mandible is squamous cell carcinoma.

5. A lady had a trauma to the neck. X-ray is attached. What is the diagnosis?



a) Hangman's fracture
b) Jefferson's fracture
c) Clay shoveler fracture
d) Sciwora

- Hangman's fracture is fracture of pars interarticularis of C2 and slip oi C2 over C3 as shown in the image.
- Jefferson's fracture involves Atlas (C1).
- Clay shoveler fracture involves the spinous process of lower cervical spine.
- Sciwora usually occurs in children <8yrs old. It is a spina cord injury without obvious radiological abnormality.

www.medicalcoupons.in

6. Patient had an injury to thumb causing thumb abduction. Which of the following can happen ?

a) Game keepers thumb
b) Bowler's thumb
c) Kaplan injury
d) Mallet finger

- Thumb abduction leads to ulnar collateral ligament damage. It is the most common injury of metacarpal phalangeal joint. Game keeper's thumb occurs due to forceful abduction of thumb as seen in gamekeepers who twist necks of small animals.

7. In fracture of upper 1/3 of forearm, it is immobilized in:

a) Pronation
b) Supination
c) Mid prone
d) Any position

Immobilization of forearm at various fracture sites: 'SMP"

- Upper 1/3rd - Supination
- Middle 1/3rd - Midprone
- Lower 1/3rd – Pronation

8. A 80 yr old lady had slip and fall, the following X-ray was taken. Which of the following is treatment?




a) Hemiarthroplasty
b) Hip spica
c) Total Hip Replacement
d) Nailing

www.medicalcoupons.in

X-ray shows fracture in the neck of Femur. The treatment for neck of femur fracture is based on age:

- < 65 years old → Fixation
- > 65 years old → Hemiarthroplasty
- Total hip replacement is done for arthritis of any age group. X- ray shows no arthritis.
- Nailing is used for shaft of femur fracture.

9. A child comes with limp and limb shortening. X-ray is below, what is given the diagnosis?



a) Perthes
b) SCFE
c) DDH
d) Fracture neck of femur

X-ray show normal right side of hip joint. In left side of hip joint, the epiphysis is smaller, shallow acetabulum and the hip joint is dislocated- indicates Developmental Dysplasia of Hip (DDH).

PSYCHIATRY DEC 2018

1. Management of violent patient in psychiatry is treated with all except:-

a) [illegible]
b) CBT
c) ECT
d) Haloperidol

Usually a violent patient in Psychiatry comes with a condition of mania or schizophrenia.

www.medicalcoupons.in

→ **BZD can be used to decrease agitation**
→ **ECT(Electro convulsive therapy) can bring the symptoms down immediately.**
→ **Haloperidol inj. can be used in the treatment of acute agitation and it is considered as the best treatment for acute agitation**

- CBT (cognitive behavioral therapy)takes longer i.e. months of therapy SO can't be used in emergency management of a violent patient. Cognitive behavioral therapy (CBT) is usually short-term approach, typically with 12 to 20 sessions. CBT is collaborative, structured, and goal-oriented intervention that targets core components of a given disorder.

2. Double depression is:-

a) Major depressive disorder from 2 years
b) Major depression with OCD
c) Depression with dysthymia
d) Depression with anxiety attack

HOW TO REMEMBER: double = D+D = Depression + Dysthymia

Double depression: Patient has symptoms of Dysthymia (mild depression symptoms continuing for 2 years) the patient develops superimposed major depressive episode.

Major depressive disorder from 2 years is called chronic depression.

3. Most commonly psychiatric illness seen in medico-surgical patient -

a) Depression
b) Schizophrenia
c) Delirium
d) Dementia

Patients who have medical comorbidities, surgical comorbidities or the postoperative patients are vulnerable to delirium. Especially if they are elderly.

4. Negative symptoms of schizophrenia are all except.-

a) Over activity
b) Anhedonia
c) Alogia
d) Apathy

www.medicalcoupons.in

Negative symptoms of Schizophrenia :

- → **Avolition**: loss of drive for meaningful activities.
- → **Apathy**: lack of concern.
- → **Anhedonia**: lack of pleasure.
- → **Asociality**: lack of social interaction.
- → **Affective flattening**/Emotional blunting.
- → **Alogia**: decreased verbal communication.

- Do not respond well to medications.**
- Poor prognostic factor.**

5. Most common acute organic mental disorder:-

a) Delirium
b) Dementia
c) Amnesia
d) Anxiety disorder

Delirium: Always has an acute onset, there's disturbance of consciousness, has fluctuating course.

Dementia: Always has an insidious onset and there's no disturbance of consciousness, has a progressing course.

DERMA DEC 2018

1. Identify condition in the shown image:-




a) Impetigo
b) Chickenpox
c) Herpes Simplex
d) Molluscum contagiousm

www.medicalcoupons.in

Impetigo is **a common and highly contagious skin infection** that mainly affects infants and young children. It usually appears as reddish sores on the face, especially around the nose and mouth and on the hands and feet. Over about a week, the sores burst and develop honey-colored crusts.

2. What are these horizontal lines called?



a) Lymphatic development
b) Dermatomes
c) Blaschko lines
d) Vascular development

Blaschko lines are consistently **V-shaped on the upper spine**, S-shaped on the abdomen, inverted U-shaped from the breast area to the upper arm, and perpendicular down the front and back of the lower extremities. They never cross the anterior truncal midline but run along it.

3. Which of the following is used for treatment of vitiligo?

a) Retinoids
b) Systemic steroids
c) Psoralen PUVA
d) Nd-YAG laser

www.medicalcoupons.in


Vitiligo is an autoimmune disease, can only be treated with immunosuppressants.

Treatment of choice PHOTOTHERAPY

Target Phototherapy is used in vitiligo with Excimer laser.

PUVA Given with UV radiation.

4. Tzanck Smear shown in the image shows:




a) Acanthosis
b) Acantholytic cells
c) Smudge cells
d) Anaplasia

On Tzanck smear if we se

- Acanthocytes - Pe nph gus
- Multi nucleated gi n cel s - Herpes

Acanthocyt /Acantholytic cells:



ultinucleated syncytial giant cells of herpes

→ Oval
→ Single cell with nucleus
→ Perinuclear halo

www.medicalcoupons.in

ANATOMY JUN 2019

1. Failure of closure of rostral neuropore at 25th day leads to ?


a) Hydranencephaly
b) Rachischisis
c) Anencephaly
d) Spina bifida

Anencephaly is the absence of a major portion of the brain, skull, and scalp that occurs during embryonic
development. It is a cephalic disorder that results from a neural tube defect that occurs when the rostral (head) end of the neural tube fails to close, usually between the 23rd and 26th day following conception.

2. Killian dehiscence is present in ?

a) Superior constrictor muscle
b) Inferior constrictor muscle
c) Middle constrictor muscle
d) Esophageal fibres


Killian's dehiscence is a potential triangular gap between the oblique fibers of thyropharyngeus and transverse fibers of cricopharyngeus (Thyropharyngeus and cricopharyngeus are 2 parts of the inferior constrictor of pharynx)



www.medicalcoupons.in

3. Which of the following gland is supplied by glossopharyngeal nerve ?

a) Submandibular salivary gland
b) Lacrimal gland
c) Parotid salivary gland
d) Sublingual salivary gland



4. Inferior scapular angle is at which level ?

a) T2
b) T4
c) T6
d) T8

www.medicalcoupons.in


- **C2** - first palpable SP below the occipital bone
- **C7 or T1** - most prominent SP at base of neck (C7 will usually slide anterior from a palpating finger with cervical extension)
- **T4** - level with the root of the spine of scapula or apex of axillary fold
- **T7-T8** - level with the inferior angle of scapula

Thoracic TP palpation Rule of 3s


- **T1-T3 TPs:** at level of corresponding SP
- **T4-T6 TPs:** ~½ segment *above* SP
- **T7-T9 TPs:** at ~level of SP of vertebrae *above*

T10-T12 have SP's that project from a position similar to T9 & rapidly regress until T12 is like T1

- **T12** - level with the head of the 12th rib
- **L4** - level with the superior border of the iliac crest
- **PSIS & S2** - level with the most inferior portion of the PSIS
- **Sacral Apex** - level with upper greater trochanter (have patient rotate hip to locate trochanter)



5. What is the structure highlighted ?



a) Supraspinatus
b) Infraspinatus
c) Teres major
d) Teres minor

www.medicalcoupons.in

Muscle highlighted in this picture is TERES MINOR

Origin	Lateral border of scapula
Insertion	Greater tubercle of humerus
Action	Shoulder joint: Arm external rotation, arm adduction; Stabilizes humeral head in glenoid cavity
Innervation	Axillary nerve (C5, C6)
Blood supply	Suprascapular artery, dorsal scapular artery

6. Which single muscle is involved in the action of arm's flexion, adduction and medial rotation?

a) Serratus Anterior
b) Pectoralis major
c) Pectoralis minor
d) Subclavius

Muscle	Action
Pectoralis major	Adducts and medial rotation of arm
Pectoralis minor	Stabilizes the scapula
Subclavius	anchors and depresses the scapula
Serratus Anterior	• protracts the scapula • laterally rotates the scapula • overhead abduction

7. One of the nerves supplying the upper limb, the upper lateral cutaneous nerve of arm, supplying the skin overlying the posterior border of the deltoid muscle as well as covering the long head of the triceps, is a branch of:-

a) Musculocutaneous nerve
b) Radial nerve
c) Ulnar nerve
d) Axillary nerve

Axillary Nerve

This nerve originates from the posterior cord of the brachial plexus with contributions from the C5 and C6 ventral rami. The axillary nerve travels through the inferior-

www.medicalcoupons.in

lateral axilla, and around the surgical neck of the humerus with the posterior circumflex humeral artery. After providing motor innervation to the deltoid, **the axillary nerve gives off the lateral superior cutaneous nerve branch which innervates the lateral skin of the shoulder**

8. Which muscle originates from medial epicondyle of humerus (FMGE June 2019)

a) Supinator
b) Pronator quadratus
c) Pronator teres
d) Brachioradialis

PRONATOR TERES-Humeral head: medial supracondylar ridge of humerus Ulnar head: Coronoid process of ulna

SUPINATOR- Lateral epicondyle of humerus, radial collateral ligament, annular ligament, supinator crest of ulna

PRONATOR QUADRATUS- anterior surface of ulna.

BRACHIORADIALIS- muscle originates from the upper two-thirds of the lateral supracondylar ridge of humerus and the anterior surface of the lateral inter-muscular septum of the arm

9. An injury to the shown area can lead to fracture of which bone ?



a) Lunate
b) Scaphoid
c) Hamate
d) Trapezium

www.medicalcoupons.in

Anatomical snuff box: The given picture shows anatomical snuff box which is a depression formed at the
posterolateral side of the wrist and metacarpal by the extensor tendons passing into the thumb. Base of the triangle is at the wrist and apex is directed into thumb. It is most apparent when thumb is extended.

Boundaries:
Lateral: Abductor pollicis longus and extensor pollicis brevis tendons.
Medial :Extensor pollicis longus tendon
Floor: Scaphoid and trapezium the distal ends of the tendons of the extensor carpi radialis longus and extensor carpi radialis brevis

Scaphoid Fracture:
The commonest carpal injury is a fracture across the waist of the scaphoid bone It is uncommon to see other injuries. In approximately 10% of individuals, the scaphoid bone has a sole blood supply from the radial artery which enters through the distal portion of the bone to supply the proximal portion. When fracture occurs across the waist of the scaphoid, the proximal portion therefore undergoes avascular necrosis.

10. A patient presents with hypothenar muscle wasting and loss of sensation of medial one and a half digits. Which nerve is involved ?

a) Musculocutaneous
b) Radial
c) Ulnar
d) Median



www.medicalcoupons.in

11. Testicular artery is a branch of -

a) Abdominal aorta
b) Common iliac artery
c) External iliac artery
d) Internal iliac artery

Branches of Abdominal aorta:

Segment Branches

- **Ventral branches**:

→ Celiac trunk -T12
→ Superior mesenteric artery - L1
→ Inferior mesenteric artery - L3

- **Dorsal branches**

→ 4 pairs of lumbar artery
→ Median sacral artery

- **Lateral branches**

→ Inferior phrenic artery (1t branch)
→ Middle suprarenal artery
→ Renal artery
→ Gonadal artery
→ Terminal branches Common iliac arteries

12. Deep inguinal ring is formed in

a) External oblique aponeurosis
b) Internal oblique aponeurosis
c) Transversus abdominis
d) Transversalis fascia

The deep (internal) ring is found above the midpoint of the inguinal ligament. which is lateral to the epigastric vessels, The ring is created by the transversalis fascia, which invaginates to form covering of the contents of the inguinal canal.

13. Inguinal ligament is formed due to

a) Cremaster muscle
b) External oblique aponeurosis
c) Internal oblique aponeurosis
d) Transversus abdominis

www.medicalcoupons.in


The inguinal ligament(also known as Poupart's ligament or the groin ligament) is a fibrous band extending from the anterior superior iliac spine to the pubic tubercle. It is formed by the external oblique aponeurosis and is continuous with the fascia lata of the thigh.

14. Abduction at the hip joint is done by

a) Gluteus maximus
b) Gluteus medius
c) Obturator internus
d) Quadratus femoris

Gluteus medius is the prime mover of abduction at hip joint. Anterior portion of Gluteus medius abduct, assist in flexion and medial rotation of hip. Posterior portion of Gluteus medius abduct, assist in external and lateral rotation of hip.

15. Identify the marked tendon in the following image ?



a) Extensor hallucis longus
b) Flexor hallucis longus
c) Extensor digitorum longus
d) Tibialis anterior

Extensor Hallucis Longus(EHL):
Origin : Middle one third of anterior surface of Fibula and adjacent interosseous membrane.

www.medicalcoupons.in

Innervation: Deep fibular nerve
Action: Extension of greater toe and dorsiflexion of foot.

ANASTHESIA JUN-2019

1. In CPR, number of chest compression per minute in an adult:

a) 30-50 per minute
b) 50-72 per minute
c) 100-120 per minute
d) 120-200 per minute

Rate : 100-120/ minute

Depth:
Adult - 5-6 cm
Pediatric - 4-5 cm
Infants - At least 4 cm

Allow complete recoil for proper filling of heart
Adequate ventilation avoid excessive ventilation

2. Spinal anaesthesia in an adult is given at this level:

a) T12-L1
b) L1-L2
c) L3-L4
d) L5-S1

Spinal cord ends at lower border of L, in adults .Hence spinal anesthesia can be given at levels below L1. L2-L3 or L4-L5 are considered best in adults.

3. Colour of nitrous oxide cylinder is?

a) Blue
b) Blue body with white shoulder
c) White
d) Black

www.medicalcoupons.in

Gas	State in cylinder	International	USA
Air	Gas	Black body and shoulders with black and white.	Yellow
Oxygen	Gas	Black body and white shoulders	Green
Nitrous oxide	Gas + liquid(below 98°F)	Blue	Blue
Carbon oxide	Gas + liquid(below 88°F)	Grey	Grey
Helium	Gas	Brown	Brown
Entonox		Blue body and shoulders with blue and white quarters	
Cyclopropane		Orange	Orange
Nitrogen	Gas	Black	Black

4. The duration of spinal anaesthesia is based directly on:

a) Dose
b) Height
c) Age
d) Total body fat

The duration of spinal anesthesia primarily depends upon the dose, intrinsic properties of local anesthetic
used, use of additives
The level and duration is directly based on dose of the drug

5. On repeated use, which of the following inhalational anaesthetic agent can cause hepatitis

a) Isoflurane
b) Halothane
c) Sevoflurane
d) Ether

Halothane causes auto immune hepatitis
Halothane produces byproduct during metabolism which causes autoimmune hepatitis
Auto immune hepatitis d/t repeated use of halothane, preexisting liver dysfunction
More seen in female patients, obese patients & middle age patients

www.medicalcoupons.in

6. Which of the following is used for day care surgery?

a) Ketamine
b) Thiopentone
c) Propofol
d) Etomidate

Drugs Used in Day Care Surgery	• Propofol (Single Best Drugs for Day Care Surgery) • Desflurane / Sevoflurane • Fentanyl / Alfentanil / Sufentanil / Remifentanil • Midazolam • Intermediate or short acting neuromuscular blocker (vecuronium , rocuronium, atracurium , cisatracurium) • Beta blocker – Esmolol • Vasodilator - Nitroglycerine, sodium nitroprusside **Anesthetic agents to be avoided in Day Care Surgery :** 1. Thiopentone 2. Halothane 3. Morphine 4. Long acting neuromuscular blockers 5. Diazepam / Lorazepam

MICROBIOLOGY JUNE- 2019

1. Which is correct about Diphtheria membrane?

a) Grey membrane that bleeds on removal
b) Grey membrane that does not bleed on removal
c) Grey pseudo-membrane that bleeds on removal
d) Grey pseudo-membrane that does not bleed on removal

Diphtheria presents with grey pseudo-membrane that bleeds on removal.
Pseudo membrane on posterior pharynx is a tough leathery coat composed of inner band of fibrin surrounded by neutrophils, RBCs and bacteria.

www.medicalcoupons.in

Pseudo membrane:



- It is adherent to basal mucosa and bleeds on removal.
- It can be localized (Tonsillar, pharyngeal etc) or widely distributed i.e.in the entire tracheobronchial tree
- Most common site - faucial pillars (Tonsillar).
- Most dangerous site laryngeal as it causes respiratory obstruction.

Bull's neck appearance seen in diphtheria is due to cervical edema and cervical lymphadenopathy.

Bull's neck appearance:



www.medicalcoupons.in

2. DNA transfer in bacteria via phage is:

a) Conjugation
b) Transduction
c) Transformation
d) Translation


Transformation

- There is direct uptake of donor DNA by the recipient cell, which may be natural or forced.
- **Transduction - donor DNA is carried in a bacteriophage and is transferred to the recipient by the mechanism used for phage infection , two type lytic and lysogenic**
- Lysogenic conversion, bacteriophages exhibit two types of cycles, virulent or lytic cycle where the host bacterium is lysed and the temperate or non-lytic cycle where host bacterium is unharmed. Eg diphtheria
- Conjugation- The donor cell contributes energy and building blocks to the synthesis of a new DNA strand, which is physically transferred into the recipient cell, usually by tube called sex pilus.
- Transferable drug resistance mediated by the factor is the most important method of drug resistance
- Mutational resistance is of great importance in tuberculosis. If a patient is treated with streptomycin alone. Initially the appear and multiply unchecked. If two or more anti-re-population by resistant mutants does not occur, as a mutant resistant to one drug will be destroyed by the other drug.

3. Sporulation occurs in which phase of bacterial growth curve:

a) Stationary phase
b) Lag phase
c) Log phase
d) Decline phase

www.medicalcoupons.in



1.**Lag phase**- Adaptation phase to culture media decreased Accumulate nutrient size
2. **Log Phase** - Exponential increases in number
3. **Stationary phase** - Start of Accumulation of toxic metabolites = Dead Live -Sporulation occur in stationary phase (S-S)
4. **Decline -** Complete accumulation of Toxin metabolites

4. Bacillus Anthrax is:

a) Gram positive cocci in cluster
b) Gram positive rods with square ends
c) Gram positive bacili with spherical ends
d) Gram negative cocci in cluster

Gram Positive bacilli

Bacillus Anthrax-

- Gram positive rods with square ends (non bulging spore) (bulging spore seen in cl. tetani)

www.medicalcoupons.in

- GPB, Non Motile, capsulated.
- Mc fadyeans reaction

M - Mc'fadyean reaction
Medusa Head appearance
I- Inverted fir tree appearance
S - String of pearl reaction
Bamboo - Bamboo stick appearance (on Gram stain)

5. Which of the following is the most likely cause in a case of granuloma with positive AFB?

a) Cat scratch disease
b) Trench fever
c) Leprosy
d) Syphilis

- Granuloma with positive AFB Is Leprosy caused by M. leprae. M leprae is acid fast bacilli.
- Syphilis caused by Treponema pallidum (spirochaete), which is not stained by acid fast stain. Diagnosed by dark ground microscopy and other serological tests.
- Trench fever caused by Bartonella quintana. Bartonella species are Gram-negative pleomorphic rods that stain poorly with the Gram stain but better with the Gimenez stain.
- Cat scratch disease caused by Bartonella henselae leads to bacillary angiomatosis.


6. Legionnaire disease is implicated in causation of which among the following?

a) Respiratory disease
b) U.T.I
c) Retroperitoneal fibrosis
d) Acute gastroenteritis

Legionellosis (Atypical pneumonia)
- Fever, muscle pain, headache
- Bacterium found in AC fresh water
- Spread by breathing


Diagnosis By urinary Antigen test & Sputum culture
Treatment - Fluoroquinolones, Azithromycin or doxycycline

www.medicalcoupons.in



Culture for legionella → BCYE (Buffered charcoal yeast Extract with Fe & Cysteine)





7. When should you perform Widal test in a case of Typhoid?

a) 1st week
b) 2nd week
c) 3rd week
d) 4th week

Widal test should be performed in 2nd week.

Lab diagnosis for typhoid:
B - Blood culture - 1st week
A - Agglutination Test 2nd week
S - Stool culture - 3rd week
U - Urine culture - 4th week


Widal test
Highly sensitive but poorly specific.
Also biological false positive in -
Infectious mononucleosis

www.medicalcoupons.in

Malaria
SLE [auto immune Disease]

Antibody titre against O → 1:100 and H → 1:200 is positive.

Flagellar antigen is more immunogenic.
Paired testing should be done, after 2 weeks of initial testing.



8. Antemortem diagnosis of rabies is made with:

a) Rabies virus specific antibodies
b) Inoculation in culture media
c) Negri bodies in hippocampus
d) Corneal impression smear

Ante mortem diagnosis
- Cornual impression smear
- Hair follicle from Nape of neck and of DFAT (direct fluorescent antigen test)

9. Hemolytic uremic syndrome associate with

a) E. coli
b) Malaria
c) Parvovirus B19
d) Bartonella henselae

EHEC
-Entero hemorrhagic E.coli
-Cause HUS (Hemolytic Uremic Syndrome)
-Shiga like toxin or Verotoxin (VT-1& VT-2)
Aka verotoxigenic E.coli
Culture media is Sorbitol Mac-Conkey agar (SMAC) and Rainbow agar used

www.medicalcoupons.in

- Diarrhoea causing E coli

1. EPEC -Entero Pathogenic Attaching/Effacing lesions to villi
2. EIEC - Entero Invasive resemble shigellosis & sereny test +ve 3.
3. ETEC-> Traveler's diarrhea CFA (colonization factor antigen 4.
4. EAEC -Aggregative persistent type of diarrhea (EAST-1 Toxin)

Parvo Virus B19: Single standard DNA Virus cause Aplastic Crisi

10. Which of the following virus is from Herpes virus family?

a) Rubella
b) Measles
c) Rabies
d) EBV

HSV - 1 - Herpes simplex virus 1
HSV - 2 - Herpes simplex virus 2
HHV - 3 -varicella zoster [chicken pox virus]
HHV - 4 - EBV
HHV - 5 - Cytomegalovirus
HHV - 8 - Kaposi saroma causing virus

11. Molluscum contagiosum is caused by ?

a) Adenovirus
b) Flavi virus
c) Rubi virus
d) Pox virus

Molluscum contagiosum is an infection caused by a poxvirus
Pearly white umbilicated nodule over the skin / genitals
Caused by MCV 1,2, 4
On biopsy, Henderson Peterson [HP] bodies seen [intra cytoplasmic]
Rx - podophyllin resin + cryosurgery

12. A patient presented with some unknown fungal infection. Microscopic examination revealed brown colored spherical fungi with septate hyphae. Possible condition:

a) Histoplasmosis
b) Chromoblastomycosis
c) Coccidioidomycosis
d) Candida albicans

Dematiaceous Fungi
- Involves skin & sub cutaneous tissues

www.medicalcoupons.in

- Sclerotic bodies seen in tissue biopsy Brown colored globose bodies Also called as bodies Show copper penny appearance

Examples -
1) Phialophora
2) Cladosporium

13. Which is correct about larval stage of Taenia solium?

a) Larva currens
b) Cysticercosis cellulose
c) Cutaneous larva migrans
d) Visceral larva migrans

- **Larva stage of taenia solium**
→ Cysticercosis cellulose
→ Taenia saginata
→ Cyst cercus bovis
- **Visceral larva migrans**
→ Toxocara canis
→ Toxocara catis
→ Angio stronglus cantonesis
- **Cutaneous larva migrans**
→ Ancylostoma
→ Brasilencs

14. N. meningitidis can be due to deficiency of this complement system:

a) C1-C4 deficiency
b) C5-C9 deficiency
c) C3 deficiency
d) C2 deficiency

- C5 - C9-MAC (membrane attack complex) deficiency can lead to increased risk of meningococcal disease.
- C1. C2, and C4 deficiencies are associated with pyogenic infections and rheumatologic disorders, particularly systemic lupus erythematosus.
- C3 deficiency is also associated with pyogenic infections and rheumatologic disorders

15. Classical complement activated by:

a) C1
b) C3 convertase
c) IgA
d) Ag-Ab complex

www.medicalcoupons.in

The complement pathway. Complement can be activated through three pathways classical, lectin, and alternative. **The classical pathway is activated when C1q binds to antibody attached to antigen**, activating C1r and C1s, which cleave C4 and C2.

PSYCHIATRY JUNE 2019

1. **A 25-year-old lady presented with sudden onset chest pain , palpitations lasting for about 20 minutes. She says there were similar episodes in the past. All the investigations were normal. What is the likely diagnosis?**

a) Acute psychosis
b) Panic attack
c) Depression
d) Mania

Panic Attack: It is an acute attack of intense anxiety with a "feeling of impending Doom" and Patient feels he is about die or he feels he is having a heart attack or he feels he is about to go crazy or he feels he will lose control.

Panic Disorders: -
Recurrent & unexpected panic attacks
Anticipatory anxiety in between the attacks

2. **All are stages of death, except:**

a) Denial
b) Anger
c) Bargaining
d) Agitation

Five Stages of Dying. **Denial, Anger, Bargaining, Depression, and Acceptance** (DABDA).

PHARMACOLOGY JUNE-2019

1. **Shortest acting corticosteroid he following is?**

a) Dexamethasone
b) Hydrocortisone
c) Triamcinolone
d) Deflazacort

www.medicalcoupons.in

- Short acting steroids → Cortisone

Hydrocortisone

- Intermediate acting → Prednisone

Prednisolone
Triamcinolone

- Long acting → Dexamethasone

Betamethasone
Paramethasone

2. Which of the following anti-epileptic drug has the highest teratogenic potential and is not prescribed to pregnant women?

a) Valproate
b) Carbamazepine
c) Phenytoin
d) Lamotrigine

- Anti-epileptic with highest risk of teratogenicity → Valproate - causes neural tube defects
- Anti-risk epileptic with lowest of teratogenicity → Lamotrigine Levetiracetam

3. You, being an ardent researcher in the field of vaccine production, are the in-charge of preparing samples to make a vaccine against the new variant of SARS-CoV 2. After getting satisfactory results in pre-clinical studies, you seek approval to do clinical studies, which requires healthy volunteers. Which of the following phase does this refer to?

a) Phase 1
b) Phase 2
c) Phase 3
d) Phase 4

- Phase 1 → Done on healthy people. Maximum tolerable dose of drug is tested.
- Phase 2 → Done on fewer patients (20-200). Efficacy and safety of drug is tested here.
- Phase 3 → Done on more patients (up-to 5000). Efficacy is confirmed in this phase.
- Phase 4 → Post marketing surveillance. Long term and rare side effects are monitored in this phase.

www.medicalcoupons.in

4. Long term use of which of the following drug is most likely associated with development of tremours

a) Salbutamol
b) Propofol
c) Betaxolol
d) Timolol

Tremors occur with β_2-agonists like salbutamol

Side effects of salbutamol : Tremor- Most common side effect Tachycardia (in overdose) Tolerance (on long term use) Hypokalemia

5. Which of the following is a long acting bronchodilator?

a) Salbutamol
b) Terbutaline
c) Adrenaline
d) Formoterol

Bronchodilators are β_2-agonists

SABA(short acting β_2 agonist) Used in acute attacks -Salbutamol ,Terbutaline
LABA (Long acting β_2 agonist) Used for prophylaxis -Salmeterol, Formoterol- fast acting , hence can be used in acute attacks.

6. Drug of choice for neurogenic diabetes insipidus is:

a) Desmopressin
b) Vasopressin
c) Terlipressin
d) Conivaptan

- Diabetes insipidus is due to deficient ADH, ADH function is to reabsorb water from kidneys
- Deficient ADH (ADH not released) →Neurogenic DI
- Resistance to ADH in kidneys → Nephrogenic DI
- **Desmopressin is DOC for Neurogenic DI**
- Thiazides are DOC for nephrogenic DI

7. Which of the following inhalational anaesthetic agent can cause hepatitis on repeated use?

www.medicalcoupons.in

a) Halothane
b) Isoflurane
c) Sevoflurane
d) Ether

Halothane causes hepatitis .It can also cause arrhythmia by making adrenaline more adrenogenic, i.e, in conditions Adrenaline Shows more adrenogenic action when halothane is used. Hence it should not be used where adrenaline is high ,i.e, pheochromocytoma

8. Preferred drug for the treatment of ventricular tachycardia is

a) Digoxin
b) Propranolol
c) Diltiazem
d) Lignocaine

Doc for ventricular tachycardia – Lignocaine

9. Which of the following drugs mentioned below is the first choice for chronic gout?

a) Allopurinol
b) Febuxostat
c) Probenecid
d) Sulfinpyrazone

Doc for acute gout : NSAID except paracetamol, aspirin
Most effective drug for acute gout : Colchicine
DOC for chronic gout: **Allopurinol** --- Inhibits Xanthine Oxidase and thus reduces production of uric acid
Febuxostat is also Xanthine Oxidase inhibitor - used in patients allergic to uric acid

10. Which of the following is used for day care surgery?

a) Ketamine
b) Thiopentone
c) Propofol
d) Etomidate

Drugs used in day care surgery:

D-Desflurane
Manmohan- Midazolam
Singh -Sevoflurane
Is -Isoflurane
A –Alfentanyl

www.medicalcoupons.in

Prime -Propofol- mc drug used in day care surgery
Minister- Mivacurium

Ketamine causes dissociative anesthesia
Thiopentone is shortest acting due to redistribution
Etomidate causes adrenal suppression

11. A female with 20 weeks of pregnancy presents with fever and dysuria. A preliminary diagnosis of cystitis was made. Which of the following drugs will be safe to use for this patient?

a) Ciprofloxacin
b) Gentamicin
c) Cotrimoxazole
d) Amoxicillin

***Amoxicillin is safe in pregnancy.**

Antibiotics safe in pregnancy
P - Penicillin
C - Cephalosporin
M - Macrolides

Ciprofloxacin causes cartilage and bone destruction
Gentamycin causes hearing loss in the fetus
Cotrimoxazole contains sulfonamides which can cause kernicterus in the child

12. A 66 year old known diabetic, on poor compliance, was referred to your institution as he had symptoms of DKA. Which of the following insulin would you prefer for the treatment of this condition?

a) Regular Insulin
b) NPH insulin
c) Insulin glargine
d) Insulin lispro

- Diabetic ketoacidosis is a medical emergency that requires immediate treatment with insulin by I/V route.
- Insulin available in W/V route is regular insulin (short-acting, soluble, crystalline zinc insulin).
- Regular insulin is DOC for diabetic ketoacidosis.

www.medicalcoupons.in

13. A 45 year old walks in to your clinic with complaints of increased hunger and thirst. On testing, was noted to levels and HbA1c of 11. If you decide to start insulin therapy, which of the have elevated blood sugar attain following long-acting insulin would you pick, knowing it does not peak concentration in plasma?

a) Insulin lispro
b) Insulin aspart
c) Insulin glulisine
d) Insulin glargine

Insulin glargine:

- Long acting insulin.
- Subcutaneous glargine given, most of it stay in subcutaneous area, very small amounts comes into the blood, gradually it keeps coming into the blood (maintains basal level)
- Never become high in plasma
- Single injection work for 24 hrs
- Glargine is called as peakless insulin
- Rapid acting insulins (insulin Lispro, insulin aspart, insulin glulsine)

14. Praziquantel is used for the treatment of

a) Strongyloidiasis
b) Trichomoniasis
c) Schistosomiasis
d) Rhinosporidiosis

Platyhelminthes (flukes & tapeworms)

- DOC for all the platyhelminths is praziquantel
- DOC for Fasciola hepatica (liver fluke) is triclabendazole
- DOC for dog tapeworm is albendazole
- DOC for trichomoniasis is metronidazole
- Surgery is done for rhinosporidiosis
- Nematodes
- DOC for nematodes is albendazole
- DOC for filaria is diethyl carbamazine
- DOC for strongyloidiasis & onchocerca volvulus is Ivermectin

15. A young male presents with reduced sleep, hyperactivity and elevated mood. He has a family history of mania. Which of the following drug should be used for long term treatment of this patient?

www.medicalcoupons.in

a) Sodium valproate
b) Lithium carbonate
c) Carbamazepine
d) Barbiturates

DOC for long term treatment of mania is lithium
L-leucocytes
I-Increase
T-Tremors (M/C side effect)
H-Hypothyroidism
I-Increase
U-Urine
M-Mothers (teratogenic), C/I in pregnancy, baby may develop Epstein anomaly

16. Drug of choice for nasal carriers of MRSA is:

a) Vancomycin
b) Teicoplanin
c) Mupirocin
d) Linezolid

Vancomycin is the DOC for the treatment of MRSA
DOC for nasal carriers of MRSA is mupirocin
DOC for VRSA is daptomycin

17. Anti-depressant drug which is used for smoking cessation is?

a) Venlafaxine
b) Topiramate
c) Bupropion
d) Amitriptyline

- Bupropion is an antismoking drug
- Buspirone is a drug which is used for anxiety

18. Which of the following drug is a selective COX-2 inhibitor?

a) Ketorolac
b) Etoricoxib
c) Piroxicam
d) Nimesulide

- Selective COX 2 inhibitor, COX1B (COX 2 inhibitor)
- Etoricoxib > longest acting COX1B
  Major side effects of COX1B (long term use causes myocardial infarction)
- Other drugs like :

→ Ketorolac

→ Piroxicam inhibit both COX & COX 2 (nonselective COX inhibitors)
→ Nimesulide

19. Aerosol drug used for treatment of RSV infection in a child is:

a) Indinavir
b) Amantadine
c) Ribavirin
d) Tenofovir

DOC for treatment of RSV infection is ribavirin
Indinavir → protease inhibitor used in HIV
Tenofovir → NRTI used in HIV
Amantadine → used for influenza virus

20. Drug of choice for treatment of Alzhiemer's disease is:

a) Donepezil
b) Atropine
c) Physostigmine
d) Fluoxetine

Drug of choice for treatment of Alzheimer's disease is: Donepezil

- Atropine is DOC for organophosphate poisoning
- Physostigmine is DOC for atropine poisoning
- Fluoxetine is DOC for depression
- AcHE inhibiting drugs

→ Donepezil
→ Rivastigmine
→ Galantamine

21. Mechanism of action of atropine in treatment of organophosphate poisoning is?

a) It inhibits secretion of acetylcholine
b) It is reactivator of acetylcholine esterase enzyme
c) It has antimuscarinic activity
d) It is agonist of acetylcholine receptors

→ Organophosphates
→ ACHE inhibitors (acetylcholine cannot be broken)
→ Level of acetyl choline increases
→ Stimulate nicotinic & muscarinic receptors
→ Over activity of M1- acidity

M2-- bradycardia
M3-- bronchoconstriction, secretion, diarrhea & urination

- Atropine is DOC for organophosphate poisoning
- But atropine doesn't block the nicotinic receptors (muscle weakness)

22. Secretion of prolactin is inhibited by?

a) Dopamine
b) Nor - adrenaline
c) Adrenaline
d) Serotonin

- PIH is secreted from hypothalamus
- inhibiting hormone PIH (dopamine)
- Dopamine in the pituitary it stimulates D2 receptors
- Stimulated D2 receptors decrease prolactin
- D2 receptors are stimulated by drugs called as cabergoline
- DOC for hyperprolactinemia is cabergoline
- DA stimulates D2 receptors — decrease prolactin

OBG JUNE 2019

1. 15-year-old girl presented with primary amenorrhea with orderly appearance of secondary sexual characteristics like breast and pubic hair. What is the next best step for this patient:

a) Reassure
b) USG
c) HSG
d) Hormonal study

In the question the girl is 15 yrs with normal secondary sexual characteristics, she is a case of primary amenorrhea and needs evaluation (reassurance could have been given had the patient been younger than 15 years)
The next important step would be an USG to check whether she has normal mullerian structures (uterus and tubes) and ovaries or not
Hormonal studies and karyotyping are done subsequently to find but the cause.

2. A 19-year-old primigravida, presents with 8 weeks amenorrhea, light bleeding and pain. O/E Uterine size corresponds to POG and Os is closed. USG reveals intra uterine pregnancy. What is preferred management in this case?

www.medicalcoupons.in

a) Estrogen plus Progesterone therapy
b) Dilatation and Curettage
c) Bed rest and Progesterone
d) Beta hCG

The above case history is suggestive of Threatened abortion.


Management -
1. Bed Rest
2. Avoid Coitus
3. Avoid Lifting weights
4. Progesterone is given to compliment the placental functio

3. A 35 year female presented with complaints of infertility. She has previous history of PID. Preliminary investigations like USG showed normal organs and hormone levels were also normal. What is the next best investigation?

a) Repeat USG
b) Hysterosalpingography
c) Endometrial biopsy
d) Urine culture and sensitivity

A past history of PID and normal pelvic USG and hormonal assays is most likely suggestive of tubal block (secondary to adhesions due to PID) as a cause of the patient's infertility.

HSG




It is the procedure to see the uterus and the tubes
Done using leech Wilkinson cannula
Done in 1st half of the menstrual cycle preferably
Drawbacks of hsg : 1) exterior of the tubes is not visible

NOTE : the best test and a better answer would have been laparoscopic chromopertubation, which would be able to visualize inside and outside of uterus and tubes and also look for extrauterine adhesions and endometriotis implants which is another important differential in this case.

4. Couvelaire uterus is seen in:

a) Placenta previa
b) Vasa previa
c) Abruptio placenta
d) Placenta percreta

Couvelaire uterus: widespread extravasation of blood into the uterine musculature and beneath the serosa. Effusions of blood are also seen beneath the tubal serosa, between the leaves of the broad ligaments, in the substance of the ovaries, and free in the peritoneal cavity.



Type III Abruption
- Most severe type of Abruptio placenta
- FHR problem, Fetal distress
- II A - without DIC
- III B - with DIC

Presentation
- Painful bleeding
- Protraction of labour
- PPH

www.medicalcoupons.in

5. Risk of endometrial cancer is least in:

a) A positive family history
b) Obesity
c) Late menopause
d) Multigravida

Risk factors for endometrial cancer-

1) Late menopause
2) Nulliparity
3) Unopposed estrogen therapy
4) Obesity, diabetes, hypertension
5) History of persistent anovulation (PCOS)
6) History of irregular & excessive premenopausal bleeding
7) Atypical endometrial hyperplasia
8) Tamoxifen therapy
9) Radiation menopause.

6. A grand multipara is a women who has given birth _ times

a) >2
b) >3
c) >4
d) ≥5

Grand multigravida - ≥ 5 times pregnant
Problems related:
- Malpresentation
- Meconium stained liquor
- Placenta previa
- Abruption'
- PPH
- Umbilical cord prolapse
- Low apgar score

7. A 26 weeks pregnant female presented with HTN for the first time. There is no proteinuria. Diagnosis of such condition

a) Chronic hypertension
b) Eclampsia
c) Gestational Hypertension
d) Preeclampsia

www.medicalcoupons.in

Gestational hypertension - bp > 140/90, > 20weeks of gestation
Pre eclamptic toxemia - gestational hypertension + proteinuria >300mg
Eclampsia - above 2 + convulsions
Chronic hypertension - hypertension before she is pregnant

8. Minimum sperm count for normal semen analysis according to WHO:

a) 5 million/mL
b) 10 million/mL
c) 15 million/mL
d) 20 million/mL

Parameters (WHO Criteria 2010)	
Parameter	**Lower Reference Limit**
Semen volume (ml)	1.5
Sperm concentration (10^6/ml)	15
Total sperm number (10^6/ejaculate)	39
Progressive motility (PR %)	32
Total motility (PR + NP. %)	40
Vitality (live sperms, %)	58
Sperm morphology (N %)	4
pH*	>/=7.2
Leucocyte* (10^6/ml)	<14
MAR/Immunobead test* (%)	<50

9. A 26 year old woman presents with infertility. She has regular cycles of 28 days with no dysmenorrhea. Her LMP was on 8th of May. When should her endometrial biopsy be planned ?

a) 22nd May
b) 8th-12th May
c) Anytime during cycle
d) 26th May


Endometrial biopsy

Best done preferably around 7 days before the onset of menstruation (that is around 21st day of a 28 day cycle)
Secretory changes prove that the cycle has been ovulatory

www.medicalcoupons.in

Can predict the functional integrity of the corpus luteum
Can detect luteal phase defect

10. In this normal menstrual cycle graph, the mark X represents levels of which hormone?



a) LH
b) FSH
c) Estrogen
d) Progesterone

Green line: LH surges and peaks before ovulation
Blue line: FSH responsible for follicular development in the early phase of the cycle, increases gradually up-to ovulation
Red line: Estrogen : increases steadily in follicular phase secondary to follicular development as estrogen is produced by the follicle and is responsible for growth of the endometrium in follicular phase
Dotted line (marked X): Progesterone: starts to rise after ovulation (after formation of corpus luteum), responsible for stabilization of endometrium. Withdrawal of progesterone leads to endometrial shedding (menstrual bleed)

11. A patient who was using CuT for contraception, presented with a 20 weeks pregnancy. The IUCD is placed at fundus, tail visible at os and she wants to continue the pregnancy. What is your next step this patient:

www.medicalcoupons.in

a) Leave IUD in-situ, continue pregnancy
b) Do medical termination of Pregnancy
c) Remove IUCD and continue pregnancy
d) Remove IUCD and do MTP

→ With intrauterine pregnancy, if the tail is seen, it should be grasped and the IUD removed by gentle outward traction This action reduces complications such as abortion, chorioamnionitis and preterm birth.
→ Specifically, in one cohort, a 54% abortion rate and 17% preterm delivery rate was noted if the device remain insitu

12. Which of the following can be a cause of Oligohydramnios?

a) Multiparity
b) Twins
c) Renal agenesis
d) Macrosomia

Oligohydramnios causes:
1) Fetal chromosomal or structural anomalies
2)Renal agenesis
3) Obstructed uropathy
4) Spontaneous rupture of membranes
5) Intrauterine infections
6) Postmaturity
7) Iugr
8) Amnion nodosum

PEDIA JUN 2019

1. A 3-year-old child with weight of 12 kg is having loose motions. He is thirsty, drinks eagerly, tears are absent and skin pinch goes back slowly. What is the best treatment plan for this child?

a) 1200 ml RL over 12 hours
b) 600 ml RL over 6 hours
c) 720ml ORS over 4 hours
d) 300 ml ORS per episode of loose stool

www.medicalcoupons.in

Assessment of severity of dehydration in children

Parameters	No Dehydration	Some dehydration	Severe dehydration
Condition	Well, alert	"Restless, irritable	"Lethargic"
Eyes	Normal	Sunken	Very sunken
Tears	Present	Absent	Absent
Mucosa	Moist	Dry	Very dry
Thirst	Normal	"Thirsty drinks eagerly"	"Drinks poorly"
Skin pinch	Goes back quickly	Goes back slowly	Goes back slowly

Management of dehydration in children

Plan A	Plan B	Plan C
No dehydration	Some dehydration	Severe dehydration
WHO ORS 5-10 ml/kg/ loose stool (Replacement of ongoing losses)	WHO ORS 75 ml/kg over 4 hours + replacement	IV RL 100ml/kg Over 6 hours(< 1year age) 3 hours (>1year age)

2. Wide open posterior fontanelle large tongue ,rough dry skin with constipation is seen in?

a) Pellagra
b) Down syndrome
c) Hypothyroidism
d) Nutritional rickets

Clinical features Hypothyroidism

- Wide open Anterior & Posterior fontanelle
- Prolonged physiological jaundice (earliest sign sometimes)
- Myxedematous facies
- Large, protruded tongue
- Skin is dry & scaly
- Hypothermia, hypothermia, hoarse cry
- Constipation
- Abdominal distension
- Umbilical hernia

3. Chronic malnutrition in child is best evaluated by?

www.medicalcoupons.in

a) Weight for height
b) Weight for age
c) Height for age
d) Ponderal index

Acute malnutrition → wasting (↓ in Weight for Height)
Chronic malnutrition → stunting (↓ in Height for age)

Neonatal hyperbilirubinemia, defined as a total serum bilirubin level **above 5 mg per dL (86 □mol per L)**, is a frequently encountered problem. Although up to 60 percent of term newborns have clinical jaundice in the first week of life, few have significant underlying disease.

4. Neonatal jaundice becomes detectable at serum bilirubin exceeding?

a) 2 mg%
b) 3 mg%
c) 4 mg%
d) 5 mg%

Neonatal hyperbilirubinemia, defined as a total serum bilirubin level **above 5 mg per dL (86 □mol per L)**, is a frequently encountered problem. Although up to 60 percent of term newborns have clinical jaundice in the first week of life, few have significant underlying disease.

5. The pathophysiology of a tet spell is a decrease in systemic vascular resistance, increasing right to left shunt which leads to a bunch more things that all just keep increasing the right to left shunt. The severity of cyanosis best described by?

a) Size of V.S.D
b) Overriding of aorta
c) Degree of RVH
d) Degree of pulmonary stenosis

Severity of TOF is:
- Directly proportional to the severity of cyanosis &
- Inversely proportional to the intensity & duration of the murmur

→ **which depends on the degree of pulmonary stenosis**

The more severe the pulmonic stenosis, the less flow into the pulmonary artery and bigger the right to left shunt. Thus, the severity of cyanosis is directly proportional to the severity of pulmonic stenosis, but the intensity of the systolic murmur is inversely related to the severity of pulmonic stenosis.

6. Infant is admitted with respiratory distress and prolonged expiration with rhonchi in chest. CXR shows hyperinflation. What is the diagnosis?

a) Pneumonia
b) Croup
c) Asthma
d) Bronchiolitis

Given scenario suggests diagnosis of Bronchiolitis: MC LRTI in infants.

Acute Bronchiolitis:
- Age group 6 months to 2 years.
- MC agent is RSV
- Clinical features: Viral prodrome followed by tachypnea, retractions and hypoxemia, wheeze and crepitation heard on examination
- Chest x-ray: Hyperinflated chest

Treatment:
- Moist Oxygen
- In immunocompromised children and those on ventilator: Nebulized Ribavirin is indicated

Acute Laryngotracheobronchitis (Croup):
- MC organism: Parainfluenza virus
- C/F's: Starts as viral prodrome and progresses to stridor, barking cough and respiratory distress
- x ray shows 'Steeple sign' due to subglottic narrowing

Treatment:
- Oxygen support
- Mild cases: Single dose dexamethasone
- Moderate to severe cases: Supportive care Single dose dexamethasone + Nebulized epinephrine

Pneumonia: h/o Fever, cough, tachypnea, retractions, sick looking child, CXR shows consolidation
Asthma: recurrent h/o wheezing present, cough which worsens at night or early morning, family h/o atopy present.

7. Protein in Cow's milk in comparison to human milk is?

a) Double
b) Triple
c) Quadruple
d) Same

www.medicalcoupons.in

	Breast Milk	Cow Mil
Proteins	1 g/dl	3.5 g/dl
Carbohydrates	7 g/dl	4.5 g/dl
Lipids	Richer in PUFA	Poorer

Composition of Breast milk (BM):

→ **Carbohydrates**:
BM contains more lactose (7g/dl) compared to cow's milk (CM), which contains 4.5 g/dl

→ **Proteins:**
BM contains lesser protein (1g/dl vs 3.5 g/dl) as compared to CM → lesser solute load on
BM contains adequate cysteine, taurine → help in brain development
BM is richer in whey proteins (lactalbumin), which are easily digestible as compared CM, which is richer in casein.

→ **Lipids:**
Breast milk is richer in PUFA (poly unsaturated fatty acids)
It contains adequate amounts of DHA (Docosa Hexaenoic Acid) = helps in brain & eye development
It is more easily digestible due to presence of lipase in breast milk.

→ **Minerals:**
Calcium : Phosphate ratio in breast milk favours calcium absorption
Iron present in breast milk is more easily absorbable than in cow's milk

→ **Vitamins:**
Breast milk contains adequate amounts of all vitamins except vitamin D. K(& B12 in strictly vegan mothers).

8. 6th day disease is?

a) Erythema Infectiosum
b) Exanthema subitum
c) Erythema marginatum
d) Erythema nodosum

www.medicalcoupons.in

Terminology	Other names for the disease	Etiology
First disease	Rubeola/ measles	Measles virus
Second disease	Scarlet fever	Streptococcus pyogenes
Third disease	Rubella/German measles/3- day measles	Rubella virus
Fourth disease	Staphylococcal scalded skin syndrome/Ritter's disease	Staphylococcus aureus
Fifth disease	Erythema infectiosum	Parvovirus B19
Sixth disease	Erythema infectiosum	Human Herpes Virus 6 or 7

MEDICINE—JUNE 2019

1. Which of the following statements are true about tremors?

1)PD is characterized by resting tremor
2)Tremor consist of alternate contraction of agonist and antagonist muscle in an oscillating rhythmic manner
3)Essential tremor is an uncommon movement disorder affecting 5%of the population
4)Normal individuals can have physiologic tremors that manifest as mild high frequency, postural or action tremors.

a) 1,2,3 only
b) 2,3,4 only
c) 1,2,4 only
d) All of the above

Options 1, 2, and 4 are true

Essential tremor is most common movement disorder.
Tremors may be most prominent:

- At rest-Resting tremor
- On assuming posture- postural tremor
- On actively reaching for target- Kinetic tremor
- On carrying out movement- Action tremor

Physiologic tremor- Normal individual and they affect upper extremities.

- Enhanced physiologic tremor - anxiety, fatigue, metabolic disturbance, drugs, toxins(alcohol, smoking and caffiene).

www.medicalcoupons.in

2. MC joint involved in diabetes is?

a) Ankle
b) Knee
c) Shoulder
d) Foot

Most common joint involved in diabetes is Foot Joint

Charcot's joint –
Painless, Paralyzed joint
Neuropathy
Delayed wound repair
Angiopathy
Non-heading ulcer on soul of test

Ankle Joint - in tables dorsalis
Knee - OA
Shoulder- Syringomyelia - Dilation of central canal of Spinal cord. MC Cervical spine involved

3. All are true about steps followed in management of ventricular fibrillation except?

a) CPR cycle duration is 2 minutes
b) Chest compressions at 100-120/min
c) Start CPR followed by immediate defibrillation
d) Intravenous access followed by Immediate defibrillation

CPR DC shock 200 J Biphasic *3 *2 min
Asystole - IV adrenaline

4. A 70 yrs old male patient was diagnosed of having heart failure.His ejection fraction was found to be <40%. Which of the following drug does not reduce the mortality for this patient?

a) Metoprolol
b) Digoxin
c) Captopril
d) K*sparing diuretics

Digoxin is not mortality reducing drug in HFrER
Mortality reducing drug in patients with HFrEF:

- ACE inhibitors
- Angiotensin receptor blockers (ARBS)

www.medicalcoupons.in

- Angiotensin receptor blocker with Neprilysin Inhibitor (ARNI)
- Beta blockers
- Mineralocorticoid receptor antagonists
  —Spironolactone
  —Eplerenone
- Isosorbide dinitrate + Hydralazine

5. Which of following leads to a continuous murmur ?

a) Peripheral Pulmonic stenosis
b) Severe Pulmonary artery hypertension
c) Type A aortic dissection
d) Rupture of cardiac chamber

Continuous murmurs begin in systole, peak near the second heart sound, and continue into all or part of diastole. Their presence throughout the cardiac cycle implies a pressure gradient between two chambers or vessels during both systole and diastole

Continues Murmur seen in
1. COA
2, PDA a/s with congenital rubella syndrome
3. Mammary souffle of pregnancy is created by enhanced arterial flow through engorged breasts and usually appears during the late third trimester or early puerperium.
4. Venous hump =dilated vein internal mammary artery
5. Rupture of sinus of Valsalva
6. Peripheral pulmonic stenosis

6. Which is true about Postural Hypotension?

a) Decrease in systolic blood pressure 20 mm Hg within 6 mins of postural change
b) Decrease in systolic blood pressure 20 mm Hg within 3 mins of postural change
c) Decrease in diastolic blood pressure 20 mm Hg within mins of postural change
d) Decrease in diastolic blood pressure 20 mm Hg within 3mins of postural change

Postural Hypotension – Supine 3 min stand
SBP↓ > 20 mmHg
DBP↓ > 10 mmHg

7. A 45-year patient working in a factory for past 20 years presents with breathlessness. HRCT chest shows pleural thickening and fibrosis. Diagnosis is?

www.medicalcoupons.in

a) Asbestosis
b) Coal worker pneumoconiosis
c) Silicosis
d) Berylliosis

ASBESTOSIS Pleural thickening + Fibrosis - Mesothelioma
Silicosis B/L Hilar LN + / miliary picture

8. Most commonly observed autosomal aneuploidy leading to spontaneous abortions is?

a) Trisomy 16
b) Trisomy 21
c) Monosomy
d) Trisomy 18

Trisomy 16: Never seen postnatally 100% abortion Rate

9. Which of these is a Nephritic syndrome?

a) Minimal change disease
b) Membranous Glomerulopathy
c) Post infectious Glomerulonephritis
d) Focal segmental glomerulosclerosis

Nephrotic Syndrome	Nephritic Syndrome
Minimal changes	PSGN/Post infectious Nephritic syndrome
Membranous	IgA Nephropathy
FSGS	

10. The following serological status is noted in a patient: HbsAg positive and HbeAg positive. Diagnosis is?

a) Carrier
b) Chronic viral hepatitis
c) Acute viral hepatitis with infectivity
d) Remote infection

www.medicalcoupons.in

Hbs Ag positive IgM anti Hbc Ag Hbe Ag positive	Acute Hepatitis + Infective

Carrier - Hbs Ag positive > 6 month
Chronic Hepatitis Hbs Ag positive

IgG positive
Remote infection . Hb Ag negative
IgG anti Hbc Ag positive

11. Chronic viral hepatitis is seen with?

a) HBV
b) HCV
c) HDV
d) HEV

Chronic Viral Hepatitis-
HCV - Cirrhosis/ HCC
DOC – Lepsidavir

12. Acute auto-graft rejection occurs within?

a) Few hours
b) <1 month
c) < 6 months
d) 6-12 months

	Hypersensitivity reaction	Time
1.Hyperacute	II	Minutes
2.Acute	II	<6 months
3.Chronic	IV	>6 months

13. Mark the true statements .

1. Pernicious anemia is an example of Type II Hypersensitivity
2. Serum Sickness is example of Type III Hypersensitivity
3. Pathergy test is an example of Type IV Hypersensitivity
4. Pathergy test is done for Reiter's disease

www.medicalcoupons.in

a) 1,2 and 3 are true
b) 2, 3 and 4 are true
c) 1, 2, 3 and 4 are true
d) 3 and 4 are true

Pernicious anemia is an example of Type II Hypersensitivity

Pernicious anemia
- Characterized by antibodies against parietal cells of stomach
- Results in vitamin B12 deficiency ---- Anemia

Serum Sickness/ Arthus reaction are example of Type III HSN
Pathergy phenomenon
- For Behcet disease
- Type IV Hypersensitivity


14. Which of the following is a causative agent of Farmer's Lung?

a) Thermophilic Actinomycetes
b) Aspergillus Fumigatus
c) Actinobacter
d) Aspergillus Flavus

Farmer's lung-T. Actinomycetes
ABPA
- Black sputum plugs in Asthmatic Patients
- Aspergillus Fumigatus

HCC Aspergillus flavus

15. What is the diagnosis ?



a) Malaria
b) Hereditary spherocytosis
c) G6PD Deficiency
d) Thalassemia

www.medicalcoupons.in

The above image shows Banana shaped gametocytes which are characteristic of Plasmodium falciparum malaria.
Hereditary spherocytosis --- Spherocytes (absence of central pallor)
G6PD --- Bite cells
Thalassemia Target cells


16. True statement regarding Whipple's disease is/ are:

1. Caused by a Protozoa Tropheryma Whipplei
2. PAS positive macrophages in the lamina propria of the small intestine
3. Culture-negative endocarditis is a common cardiac presentation
4. Treated with IV Ceftriaxone

a) 1,2 and 4 are true
b) 2, 3 and 4 are true
c) 1, 2 and 3 are true
d) 1 and 2 are true

Whipple's disease

By Bacteria Tropheryma Whipplei
PAS positive (Schiff reagent- Intracellular bacteria found in macrophage in gut)
PAS positive macrophages in the lamina propria of the small intestine in Whipple's disease:



Rx

IV - Ceftriaxone
Oral - Cotrimoxazole
Whipple's triad -seen in Insulinoma
Whipple's Sx- is performed in Ca of head pancreas.
Extra Edge: The most common type of cardiovascular presentation in Whipple's disease is culture-negative endocarditis.

www.medicalcoupons.in

17. Most common site of Brain metastasis is?

a) Lung cancer
b) Head and neck cancer
c) Prostate cancer
d) Breast cancer

Oat Cells Ca > Breast Ca > Malignant Melanoma

18. Which is not seen in Tumour lysis Syndrome?

a) Hypophosphatemia
b) Hypocalcemia
c) Hyperuricemia
d) Hyperkalemia

Tumor lysis syndrome is characterized by **high blood potassium (hyperkalemia)**, high blood phosphate (hyperphosphatemia), low blood calcium (hypocalcemia), high blood uric acid (hyperuricemia), and higher than normal levels of blood urea nitrogen (BUN) and other nitrogen-containing compounds (azotemia).

19. Comment on the diagnosis for SAP normal, PTH normal, Vitamin D3 normal with elevated serum Calcium values?

a) Vitamin D intoxication
b) Hyperparathyroidism
c) Multiple myeloma
d) Nutritional rickets

Multiple myeloma:
→ SAP Negative
→ PTH Negative
→ D3 Negative
→ ↑ Sr.Cal

20. Which of the following interfere with iron absorption?

a) Vitamin
b) Phytates
c) Oxalate
d) Myoglobin

Phytates - compete with iron absorption

21. Most sensitive and specific marker for MI is?

a) Troponin
b) Cytokeratin
c) Myoglobin
d) CPK- MM

- Most sensitive & specific marker for MI = Troponin
- 1st Investigation - ECG
- Best Investigation - Troponin I * 2 times
- Myoglobin 1 - Skeletal Muscle damages also
- Cytokeratin - Cancer
- CPK MM - Only right at Skeletal Muscle


PATHO- JUNE 20[illegible]

1. Berry aneurysm most commonly occurs due to?

a) Endothelial injury of vessel due to HTN
b) Muscle intimal elastic lamina layer defect
c) Muscle and adventitial layer defect
d) Adventitia defect

Berry aneurysm due to defect in tunica media and intimal elastic lamina layer leading to weakening of vessel wall and out-pouching in anterior portion of circle of willis. Saccular Aneurysms: is a/the most common cause of clinically significant subarachnoid hemorrhage is rupture of a saccular ("berry") aneurysm in cerebral artery following trauma. Patient will describe the pain as worst headache of his life. Bloody tap on lumbar puncture is seen. Nimodipine is a cerebro-selective vasodilatory drug helpful for treatment of symptoms associated with subarachnoid hemorrhage. Berry aneurysm is associated with adult polycystic kidney disease (ADPKD).

www.medicalcoupons.in



2. A 45-year patient working in a factory for past 20 years presents with breathlessness. HRCT chest shows pleural thickening and fibrosis, What is the person suffering from

a) Asbestosis
b) Coal worker pneumoconiosis
c) Silicosis
d) Berylliosis

Occupational lung disease/pneumoconiosis:

Caused by dust particle with range of 1-5 micron and it mainly involves the upper lobe of lung.
HRCT will show pleural thickening and fibrosis.

Asbestosis: In contrast to coal workers' pneumoconiosis and silicosis, asbestosis begins in the lower lobes and sub pleurally. The middle and upper lobes of the lungs become affected as fibrosis progresses. The scarring may trap and narrow

www.medicalcoupons.in

pulmonary arteries and arterioles, causing pulmonary hypertension and cor pulmonale. Pleural plaques, the most common manifestation of asbestos exposure, are well-circumscribed plaques of dense collagen that are often calcified. They develop most frequently on the anterior and posterolateral aspects of the parietal pleura and over the domes of the diaphragm.
Asbestos is responsible for causing mesothelioma and bronchogenic carcinoma. Most common lung cancer associated with asbestosis bronchogenic carcinoma. Most specific lung cancer associated with asbestosis is mesothelioma. Most common pneumoconiosis: silicosis/sand blaster disease.
Coal worker pneumoconiosis: anthracosis (black lung) Berylliosis: associated with aerospace industry.

3. Correct about of Anaemia of chronic disease

a) High ferritin
b) Increased percentage of saturation of transferrin
c) High transferrin
d) High serum iron

In patient with Anemia of Chronic disease, there is increase in blood level of hepcidin protein which interferes with metabolism of iron. Hepcidin is a negative iron regulator. It inhibit the usage of iron from storage form of iron and GI tract. Therefore these patient develops microcytic anemia. Lab findings:
→ Serum iron decrease
→ Transferrin saturation is decrease
→ Increase in the concentration of serum ferritin
→ Decrease in total iron binding capacity

Microcytic hypochromic anemia:

- S: Sideroblastic
- I: Iron deficiency
- T: Thalassemia
- A: Anemia of chronic disease

4. Which of these is a Nephritic syndrome?

a) Minimal change disease
b) Membranous Glomerulopathy
c) Post infectious Glomerulonephritis
d) Focal segmental glomerulosclerosis

The nephritic syndrome is characterized by hematuria, oliguria with azotemia, proteinuria, and hypertension.

www.medicalcoupons.in

The most common causes are immunologically mediated glomerular injury; lesions are characterized by proliferative changes and leukocyte infiltration.

- Acute postinfectious glomerulonephritis typically occurs after streptococcal infection in children and young adults but may occur following infections with many other organisms; it is caused by deposition of immune complexes, mainly in the subepithelial spaces, with abundant neutrophils and proliferation of glomerular cells. Most affected children recover; the prognosis is worse in adults.
- Rapidly progressive glomerulonephritis (RPGN) is a clinical entity with features of the nephritic syndrome and rapid loss of renal function. Nephrotic syndrome is caused by derangement in glomerular capillary walls resulting in increased permeability to plasma proteins.

The manifestations of the syndrome include:

→ Massive proteinuria, with the daily loss of 3.5 g/m² or more of protein less in children)
→ Hypoalbuminemia, with plasma albumin levels less than 3 gm/dl
→ Generalized edema
→ Hyperlipidemia and lipiduria

5. The following serological status is noted in a patient: HbsAg positive and HbeAg positive. Diagnosis is?

a) Acute viral hepatitis
b) Chronic viral hepatitis
c) Acute viral hepatitis with infectivity
d) Remote infection

Acute viral hepatitis: HBsAg +/IgM antiHBc
Chronic viral hepatitis: HBsAg +/IgG anti HBc
Acute viral hepatitis with infectivity: HbeAg+
Remote infection: anti HBs

6. Chronic viral hepatitis is seen with all of the following viruses, except?

a) HBV
b) HCV
c) HDV
d) HEV

- Chronic hepatitis is defined as symptomatic, biochemical, or serologic evidence of continuing or relapsing hepatic disease for more than 6 months.
- Etiology rather than the histologic pattern is the most important determinant of the probability of developing progressive chronic hepatitis.
- Chronic viral hepatitis is seen in HBV, HCV, HDV.
- Hepatitis A virus (HAV) infection is a self-limited disease and does not lead to chronic hepatitis or carrier state.

www.medicalcoupons.in

- HAV and HEV (in immunocompetent hosts) do not cause chronic hepatitis

7. Acute graft rejection occurs within?

a) Few minutes
b) Few hours
c) < 6 months
d) 6-12 months

Hyper acute rejection: type 2 HSR, happens over span of few minutes to hours. Eg: kidney transplant
Acute rejection: Few weeks to months(< 6 months)
Chronic rejection: After 6 months.
Graft vs host disease (Runt Disease):
Acute GVH: 100 days
Chronic GVH: > 100 days

8. Type II Hypersensitivity is seen in?

a) Pernicious anaemia
b) Serum sickness
c) Arthus phenomenon
d) Pathergy phenomenon

Type 2 HSR is seen in following condition:

My : Myasthenia gravis,
Blood : Blood transfusion (mismatch)
Group : Graves, goodpasture
Is: Immune hemolytic anemia
Rh: Rheumatic heart disease
Positive: Pernicious anemia/ pemphigous vulgaris

Type 3 HSR: SHARP

S: Serum sickness/SLE
H: Henoch schonlein purpura
A: Arthus reaction
R: Reactive arthritis
P: Post strep glomerulonephritis
Pathergy phenomenon (Type 4 HSR) is useful in diagnosis of Behcet syndrome(oculo-oro-genital syndrome): patient complains of recurrent oral ulcer eye lesion, skin lesion and genital lesion.

www.medicalcoupons.in

9. A stable boy admitted with acute onset of fever. cough and dyspnea. Chest radiograph showed findings of pneumonia for which he was treated with antibiotics; showed good response. After month he again presented in hypoxemia state with progressive dyspnea. CT chest showed diffuse ground glass changes. Transbronchial biopsies showed numerous small granulomas and marked lymphocytosis. Which of the following is a causative agent for given disease?

a) Thermophilic Actinomycetes
b) Aspergillus Fumigatus
c) Actinobacter
d) Aspergillus Flavus

Farmer's lung is. subtype of Hypersensitivity pneumonitis resulting from exposure to dusts generated from humid, warm, newly harvested hay that permits the rapid proliferation of the spores of thermophilic actinomycetes.

	Pathogenesis	C/F
Acute Phase	1st exposure Ab formation Repeat exposure - Ag-Ab immune complex formation ~ type III hypersensitivity	Cough / Dyspnea /Fever
Chronic Phase	Non caseating granuloma Type IV HR > Type III HR	Interstial fibrosis – Progressive dyspnea Respiratory Failure Cyanosis

10. Most sensitive indicator for Iron deficiency anaemia?

a) Serum Ferritin
b) TIBC
c) Percentage saturation of transferrin
d) Bone marrow iron

→ Most specific test for ron deficiency anemia is bone marrow iron. But bone marrow iron test is invasive procedure
→ Most sensitive indicator for iron deficiency anemia: serum ferritin.

Lab finding in iron deficiency anemia:
- Serum ferritin : decrease (early indicator)
- Serum iron: decrease
  % Tf saturation: decrease
- TIBC: increase

www.medicalcoupons.in

11. Whipple's disease is caused by ?

a) Bacteria
b) Virus
c) Protozoa
d) Helminths

Whipple disease is cause by bacteria called tropheryma whipplii is responsible fot causing infux of lot of macrophages leading to compression of lacteals and impairment in absorption of nutrients.

Rod shaped bacillis in macrophage is seen. Patient complains of malabsorption. Pas +ve diastase resistant is seen macrophage.
Treatment with antibiotic will show dramatic improvement.

12. Most common primary for brain metastasis is which of the following?

a) Lung cancer
b) Head and neck cancer
c) Prostate cancer
d) Breast cancer

Most common Primary for brain metastasis is - LUNG CANCER
Particular Lung cancer which is notorious for causing CNS metastasis - SMALL CELL LUNG CARCINOMA
Most common Primary malignant CNS tumor seen in children - MEDULLOBLASTOMA
Most common Primary malignant CNS tumor seen in adults GLIOBLASTOMA
Most common Primary benign CNS tumor seen ir adults. MENINGIOMA

13. 12-year-old Boy came with complaints of Severe Abdominal pain and headache for the past 2 days, H/0 On & off for we the past 1 week, associated with myalgia, and several episodes vomiting self-medicated with Tab Paracetamol with no relief. Presents with Dengue NS -1 antigen (+). What is the first line of defense of the body in the given scenario?

a) NK cell
b) T cell
c) Histiocyte
d) Macrophage

The major components of innate immunity are epithelial barriers that block entry of microbes, phagocytic cells (mainly neutrophils and macrophages), dendritic cells,

natural killer cells and other innate lymphoid cells and several plasma proteins, including the proteins of the complement system.

NK cells

- Ability to kill virus-infected cells and tumor cells
- NK cells act as an early line of defense against viral infections and some tumors.
- Two cell surface molecules, CD16 and CD56, are commonly used to identify NK cells.
- CD16 is an Fc receptor for IgG, and it confers on NK cells the ability to lyse IgG-coated target cells. This phenomenon is known as antibody-dependent cell-mediated cytotoxicity (ADCC).

The functional activity of NK cells is regulated by a balance between signals from activating and inhibitory receptors

- NK cell inhibitory receptors recognize self-class I MHC molecules, which are expressed on all healthy cells. The inhibitory receptors prevent NK cells from killing normal cells.
- Virus infection or neoplastic transformation often enhances expression of ligands for activating receptors and at the same time reduces the expression of class I MHC molecules.

14. Which of the following interfere with iron absorption?

a) Vitamin
b) Phytates
c) Oxalate
d) Myoglobin

Maximum site of Iron absorption - DUODENUM
The form in which Iron is absorbed - Fez+
Absorption of inorganic iron is enhanced by ascorbic acid, citric acid, HCL, amino acids, and sugars the diet and inhibited by Phytates and tannates (found in tea), carbonates, oxalates, and phosphates.

15. Most sensitive and specific marker for myocardial infarction is?

a) Troponin
b) Cytokeratin
c) Myoglobin
d) CPK-MM

The laboratory evaluation of MI is based on measuring the blood levels of proteins that leak out of irreversibly damaged myocytes, the most useful of these molecules are cardiac specific troponin T and I (cTnT and cTnI).

→ Most sensitive and specific marker for MI-Troponin
→ Most important enzyme or making the diagnosis of Reinfarction - Troponin.

www.medicalcoupons.in

→ 1st enzyme that increases after the attack of MI- Myoglobin.

16. Antecedent diagnosis of Group A streptococcal infection in Acute rheumatic fever can be made by?

a) ASO
b) CRP
c) ESR elevation
d) Low C3 levels

Acute RF typically appears 10 days to 6 weeks after group A streptococcal infection in about 3% of patients.
It occurs most often in children between ages 5 and 15, but first attacks can occur in middle to later life. Although pharyngeal cultures for streptococci are negative by the time the illness begins, antibodies to one or more streptococcal enzymes, such as streptolysin O and DNase B can be detected in the sera of most patients with RF.

17. Most common anterior mediastinal tumor?

a) Thymoma
b) Neurofibroma
c) Pericardial cyst
d) Bronchogenic cyst

The designation thymoma is restricted to tumors of thymic epithelial cells. Such tumors typically also contain benign immature T cells.
The tumors usually occur in adults older than 40 years of age; thymomas are rare in children. Males and females are affected equally. Most arise in the anterior superior mediastinum, but sometimes they occur in the neck, thyroid, pulmonary hilum or elsewhere

18. Most characteristic finding of diabetic nephropathy?

a) Kimmelstein Wilson disease
b) Diffuse glomerulosclerosis
c) Focal segmental glomerulosclerosis
d) Armani Ebstein change

Diabetic Nephropathy Three lesions are encountered: (1) glomerular lesions; (2) renal vascular lesions, principally arteriolosclerosis; and (3) pyelonephritis, including necrotizing papillits.

The most important glomerular lesions are capillary basement membrane thickening, diffuse mesangial sclerosis, and Nodular glomerulosclerosis.

www.medicalcoupons.in

Capillary Basement Membrane Thickening. Widespread thickening of the glomerular capillary basement membrane (GBM) occurs in virtually all cases of diabetic nephropathy and is part and parcel of the diabetic microangiopathy. Diffuse Mesangial Sclerosis. This lesion consists of diffuse increase in mesangial matrix. Nodular Glomerulosclerosis. This is also known inter capillary glomerulosclerosis or Kimmelstiel-Wilson disease. It is the Most characteristic finding of Diabetic Nephropathy.

19. A 23 year old individual with serum alkaline phosphatase normal, PTH normal, Vitamin D3 normal with elevated serum Calcium values what will be the diagnosis ?

a) Vitamin D intoxication
b) Hyperparathyroidism
c) Multiple myeloma
d) Nutritional rickets

Multiple myeloma is a plasma cell neoplasm commonly associated with lytic bone lesions, hypercalcemia, renal failure, and acquired immune abnormalities.

Clinical Features.
The clinical features of multiple myeloma stem from
(1) the effects of plasma cell growth in tissues, particularly the bones
(2) the production of excessive Igs, which often have abnormal physicochemical properties; and
(3)The suppression of normal humoral immunity.

Bone resorption often leads to pathologic fractures and chronic pain. The attendant hypercalcemia can give rise to neurologic manifestations, such as confusion, weakness, lethargy, constipation, and polyuria, and contributes to renal dysfunction. Decreased production of normal Igs sets the stage for recurrent bacterial infections. Cellular immunity is relatively unaffected. Of great significance is renal insufficiency, which trails only infections as a cause of death. The pathogenesis of renal failure ,which occurs in up to 50% of patients, is multifactorial. However, the single most important factor seems to be Bence Jones proteinuria, as the excreted light chains are toxic to renal tubular epithelial cells. Certain light chains (particularly those of the λ6 and λ3 families)are prone to cause amyloidosis of the AL type, which can exacerbate renal dysfunction and deposit in other tissues as well.

20. MC cause of atypical pneumonia?

a) Mycoplasma pneumoniae
b) Klebsiella pneumoniae
c) Hemophilus influenzae
d) Chylamydia

www.medicalcoupons.in

MC cause of Atypical pneumonia- Mycoplasma Pneumonia aka "Eaton's agent".
This organism does not have cell wall which makes them resistant to the treatment by Penicillin's.
The infection will be treated by the use of Macrolide antibiotic (Azithromycin).

BIOCHEM JUNE 2019

1. Enolase is inhibited by which of the following substances ?

a) NaF
b) Fluoroacetate
c) iodoacetate
d) Potassium oxalate

- NaF is added for in vitro test of blood glucose estimation.
- NaF Inhibits glycolysis
- If not added in the blood testing vial, glucose used by RBC and WBCs is not measured, then the reported glucose value will be falsely decreased or underestimation of real value.

Fluoroacetate--non-competitive inhibitor of Aconitase (TCA cycle)

Iodoacetate--Inhibits glyceraldehyde 3-phosphate Dehydrogenase (glycolysis)

Potassium oxalate--anti-coagulant

2. Maple syrup urine disease is due to deficiency of:

a) □ keto acid decarboxylase
b) α keto acid carboxylase
c) α keto acid desulphate
d) α keto acid chain oxidase

BCAA —b6→ BCKAL →
transamination α-keto acid decarboxylase

accumulation of BCKA (branched chain keto acid)

Branched chain amino acid - valine, leucine, isoleucine this amino acid are essential amino acid
Maple syrup urine disease - defect in branched chain amino acid metabolism
Enzyme - alpha keto acid dehydrogenase alpha- keto acid decarboxylase
C/F- burnt sugar like odour in urine

www.medicalcoupons.in

- Ketosis
- Mental retardation
- Abnormal muscle tone
- Coma & death

3. Collagen is rich in which amino acid:

a) Glycine
b) Arginine
c) Phenyl - alanine
d) Tyrosine

The Main Amino Acid in Collagen

Collagen is a structural protein that contains high amounts of **glycine**. In fact, every third to fourth amino acid in collagen is glycine. Collagen is the most abundant protein in your body. It provides strength for your muscles, skin, cartilage, blood, bones and ligaments.
The triple-helical structure of collagen arises from an unusual abundance of three amino acids: **glycine, proline, and hydroxyproline**.

4. Nitric oxide is derived from

a) Arginine
b) Histidine
c) Glycine
d) Aspartate

Nitric oxide (NO) is a gaseous molecule with autocrine and paracrine effects on many cell types. NO is synthesized from **the amino acid l-arginine by NO synthase (NOS)** and is involved in a myriad of cellular functions, including muscle relaxation, neuronal signaling, and immune function.

5. De--Novo synthesis of Fatty Acid requires which coenzyme?

a) NADPH
b) TPP
c) FAD
d) NAD

NADPH Reductive Biosynthesis Pathway

Fatty acid Synthesis

www.medicalcoupons.in

1st enzyme Acetyl CO A carboxylase

- Require biotin, ATP, CO2 and Bicarbonate

2nd enzyme -Fatty Acid Synthesis complex requires NADPH

- TPP is required for carbohydrate metabolism & Trans ketolase (HMP Pathway)
- FAD and NAD are required in beta oxidation of fatty acids and carbohydrate metabolism (Glycolysis and TCA)

6. All are cofactors for Dehydrogenase except:

a) NADP
b) FAD
c) NAD
d) SAM

Dehydrogenases --remove H2 and which is usually taken up by their cofactors;


1. FAD → FADH2 →(ETC)
2. NAD → NADH →(ETC)
3. NADP → NADPH
   - NADPH → used in anabolic pathway

→ SAM (S - Adenosyl Methionine) (Donor of methyl group) is a nucleotide but not a cofactor.

7. After strenuous exercise alkaline pH of skeletal muscle is in which glycogen storage disease?

a) Mc Ardle's disease
b) Von- Gierke's disease
c) Her's disease
d) Pompe's disease

Glycogen Phosphorylase absent an Mc Ardle's disease lactate will not formed alkaline PH
Normal Condition



www.medicalcoupons.in

8. Most potent lipid phase antioxidant:

a) Vitamin A
b) Vitamin E
c) Vitamin C
d) Vitamin K

Vit E delays the progression of atherosclerotic plague because its convert oxidased LDH to Normal LDH
Vit E act simultaneously with selenium & selenium is part of glutathione peroxidase which is involved in the oxidant Pathway in cell

Other anti-oxidant Vitamin - Vit A, C & D

Vit K - Fat soluble vitamin
only fat soluble vitamin which has co-enzyme role in carboxylation of glutamate in clotting factors 2,7,9,10

9. Wernicke encephalopathy caused by:

a) Vit B1
b) Vit B2
c) Vit B7
d) Vit B12

Vit B1 deficiency seen in chronic alcoholics & lead to Wernicke's Korsakoff psychosis and patient has memory loss / confabulation ,cerebral hemorrhage, muscles paralysis

Vit B2 → riboflavinosis → 2C
- Corneal Vascularization
- Cheilosis

Vit B7 → alopecia, bowel inflammation, muscles pain

Vit B12 → megaloblastic anemia + Neurological manifestation

10. Pheochromocytoma is a rare, usually a benign tumor that develops in chromaffin cell (pheochromocytes) of adrenal gland. The most specific biomarker pheochromocytoma is which one of following?


a) VMA
b) Catecholamine
c) HIAA
d) Serotonin

www.medicalcoupons.in

Vanillylmandelic acid (VMA) and other catecholamine metabolites (homovanillic acid: HVA and dopamine) are typically elevated in patients with catecholamine-secreting tumors example: neuroblastoma, pheochromocytoma, and other neural crest tumors.

Pheochromocytoma - tumor or Adrenal gland which secretes high amount of epinephrine & Nor-epinephrine
catabolic products of catecholamine – VMA

End Products of Catecholamines



11. Which cell releases histamine?

a) Mast cells
b) Eosinophils
c) T-cells
d) NK cells

Mast cell release Histamine & Heparin
Histamine is vasodilators & increase permeability for WBC in Blood vessels.

FORENSIC JUNE 2019

1. Grounds of divorce:

a) Sterility
b) Frigidity
c) Impotence developing after the marriage
d) Poverty

Impotence at the time of marriage is a ground for nullity of marriage.
Impotence developing after the marriage is a ground for divorce.
Sterility is not a ground for divorce.

Impotence: is the inability of a person to perform sexual intercourse and achieve gratification.

www.medicalcoupons.in

Frigidity: It is the inability to initiate or maintain the sexual arousal pattern in female (absence of desire for sexual intercourse or incapacity to achieve orgasm).

Sterility: It is the absolute inability of either a male or a female to procreate. In male, it is inability to make a female conceive, and in females, it is inability to conceive children.

2. IPC for causing abortion without women's consent:

a) 312
b) 313
c) 314
d) 315

IPC - Abortion

Section	Defines/Punishes
312	Punishment for Criminal abortion with consent of woman
313	Punishment for Criminal abortion without consent of woman
314	Punishment for Criminal abortion leading to death of mother
315	Punishment for Causing death of quick unborn child

Criminal Abortion is dealt under Section 312-316 IPC:

Sec. 312 I.P.C. whoever voluntarily causes criminal abortion is liable for imprisonment up to three years, and/or fine; and if the woman is quick with child the imprisonment may extend up to seven years. It is necessary that the woman should be pregnant and that abortion should be carried with her consent. Both the person causing the abortion and the woman are liable for punishment.

Sec. 313 IPC: if the miscarriage is caused without the consent of the woman, the imprisonment may be up to ten years.

Sec. 314 PC: if a pregnant woman dies from an act intended to cause miscarriage, the offender is liable to be punished with rigorous imprisonment which shall not be less than ten years and also fine up to two lakh rupees.

www.medicalcoupons.in

Sec. 315 IPC: a person doing an act intended to prevent the child from being born alive or to cause to die after its birth, is liable to be punished with imprisonment up to ten years.

Sec. 316 IPC: causing death of quick unborn child by any act amounts to culpable homicide, and the punishment may extend up to ten years imprisonment.

3. **In corrosive acid case is opened along:**
   a) Lesser curvature
   b) Greater curvature
   c)Vertical
   d) Pylorus

In corrosive acid poisoning, it is open along the greater curvature.

Toxic substances may be held in high concentrations in the rugae and crypts of the mucosa in the actual stomach wall.
The pathway of acids and alkalis in food-filled stomach starts along the lesser curvature of the stomach and leads to the pylorus, which explains the location of greatest damage in stomach.

4. **Inhaling the cloth soaked in drugs**

   a) Bagging
   b) Spraying
   c) Huffing
   d) Sniffing

Huffing: Applying the chemical to a cloth/rag and then inhaling by covering nose and mouth with the cloth/ rag

Sniffing: Inhaling fumes from the liquid in an open container.

Bagging: Placing the chemical in a bag and then putting it over the face.

5. **One of the victims of the fire which had destroyed 6 huts in a nearby slum has decided to give a dying declaration. The doctor was called in as part of the team to record the statement. Which among the following is the role of the doctor ?**

   a) To assess compos mentis
   b) To record statement even in presence of magistrate
   c) Cross examine the person
   d) Put person under oath before declaration

www.medicalcoupons.in

Dying declaration : written or oral statement of a person, who dying as a result of some unlawful act, relating to the material facts of the cause of death or the circumstances surrounding it.

The law does not provide who can record dying declaration, nor is there any prescribed form, format or procedure for the same.

The doctor should certify that the person is conscious and his mental faculties are normal, i.e. he is in compos mentis.

Dying Declaration VS Dying Deposition

	Dying Declaration	Dying Deposition
Recorded by	Magistrate/doctor/lay person	Magistrate only
Oath	-	+
Cross examination of person	-	+
Superiority	Less	Superior
Accepted under	SEC(3) IEA	Not accepted in India

PSM JUNE - 2019

1. Inertization is:

a) Reducing organic and combustible waste to inorganic
b) Burning biomedical waste
c) Biomedical waste converted into non-harmful
d) To avoid water contamination

Inertization technique is used when large volume of toxic Bio medical waste is converted to non- toxic waste.

It is achieved by mixing,
→ 15% lime and
→ 15% cement with
→ 65% biomedical waste
→ 5% water

2. Amplifier host of Japanese encephalitis:

a) Pig
b) Dogs
c) Cats
d) Birds

www.medicalcoupons.in

Japanese Encephalitis

→ Actual host →Ardeid birds
→ Natural / Amplifier host → Pig
→ Vector Culex → Tritaeniorhynchus
→ Live vaccine → SA - 14-14-2
→ Two doses of vaccine SA-14-14-2 is given, 1st at 9 months & 2nd at 16 to 24 months of age.
→ Mainly affects children under 1-15 years of age.

3. Best index for calculation of nutrient value of protein

a) Serum albumin concentration
b) Arm-muscle circumference
c) Creatinine-height index
d) Serum transferrin level

- Assessment of protein nutritional status.
- 5 Major indicators

→ Arm muscle circumference
→ Creatinine height index
→ Serum Album
→ Serum Transferrin
→ Total Body Nitrogen (TBN)

- Serum albumin is the best indicator
- Serum albumin concentration

→ >3.5 g/ dl - Normal
→ 3.0-3.5 Mild to moderate malnutrition
→ < 3.0 - Severe malnutrition

4. Best index for measurement health indicator in society:

a) Under 5 mortality rate
b) Infant Mortality Rate
c) Child death rate
d) Maternal mortality rate

Infant mortality is the indicator / marker of
→ Health status population
→ Health care infants
→ Health services

Under 5 mortality is marker of → Social development of the Country.
Material mortality rate is the indicator of → Maternal care.

www.medicalcoupons.in

Perinatal mortality rate is the very good indicator of → Combined Maternal and newborn care.

5. A community medicine student was asked to explain about natural sources of Vit.A. Which of the following has the maximum natural source of vitamin A?

a) Carrot
b) Halibut liver oil
c) Cod fish oil
d) Cow milk

- Halibut Liver Oil has the maximum natural content of Vitamin A & also it has the richest source of the Vitamin D.
- Among fruits Ripe Mango & among vegetables carrot have high content of Vitamin A.
- Vitamin A deficiency causes Xerophthalmia.

6. According to WHO, marker for obesity index is:

a) Quetelet index
b) Chandler index
c) Pearl index
d) Sullivan's index

BMI is still referred to as the Quetelet index, for which the formula is: **BMI (kg/m2) = mass (kg) / height (m)2**.

Body Mass Index Values
- Normal → 18.5 to 25
- Overweight → 25 to 29.9
- Obesity → > 30
- Underweight → < 18.5

BMI in Percentile
- >95th Percentile → Obesity
- >85th Percentile → Overweight
- <5th Percentile → Underweight

7. According to WHO, which one is a notifiable disease?

a) Yellow fever
b) Polio
c)HIV
d) Malaria

www.medicalcoupons.in

The International Health Regulations (WHO IHR) does not cover HIVAIDS
Immediately Notifiable Diseases: Smallpox, Human Influenza, Wild Polio, SARS
Potentially Notifiable Diseases:

- Public health importance: Cholera, Plague, Yellow fever, Viral hemorrhagic fevers, West nile ever, Dengue, Rift valley fever, Meningococcal disease
- Biological/Chemical/Radiological events
- Serious illnesses of Unknown origin

8. Gujarat recently reported an massive earthquake disaster causing loss of life and leading to sufferings. Which of the following is the most common reported disease in post disaster period?

a) Acute gastroenteritis
b) Pneumonia
c) Leptospirosis
d) Malnutrition

- Most Common reported disease in post disaster phase is Acute Gastroenteritis, Typhoid & Cholera.
- Most important step during Post-disaster phase is, Chlorination of water > Vaccination
- All vaccines are CI in post-disaster phase; except Measles

Extra edge:
Chairman of National Disaster Management Authority (NDMA) of India is Prime Minister of India

9. Amount of sodium in ORS as per WHO:

a) 65 mmol/L
b) 75 mmol/L
c) 20 mmol/L
d) 100 mmol/L

www.medicalcoupons.in

New formula WHO-ORS

Content		Concentrations	
NaCl	: 2.6 g	Na^+	— 75 mM
KCl	: 1.5 g	K^+	— 20 mM
Trisod. citrate	: 2.9 g	Cl^-	— 65 mM
Glucose	: 13.5 g	Citrate	— 10 mM
Water	: 1 L	Glucose	— 75 mM

Total osmolarity 245 mOsm/L

(available as ORETRAL-A, ELECTROBION, ELECTRAL 21 g sachet for 1000 ml; WALYTE, RELYTE 4.2 g sachet for 200 ml).

10. As compared to cow milk, the protein content of human milk is:

a) 3-4 times more than cow milk
b) 3-4 times less than cow milk
c) Double than cow milk
d) Same as cow milk

	Human Milk	Cow's Milk
Total Proteins	11	33
Casein	4	28
Soluble Proteins	7	5

Human milk is higher in Lactose, Iron, Water and Ca: P ratio.
Cow's milk is higher in Proteins, Fats, Energy & absolute quantity of Calcium and Phosphorous.

11. ORS is diluted with:

a) Normal saline
b) Distilled water
c) Ringer lactate
d) Distilled water

www.medicalcoupons.in

BCG is available in Powder form, which is diluted with Normal saline and it should be used within 4 hours,
Otherwise Toxic shock syndrome can occur.

- Strain →Danish - 1331
- Dose → 0.1 ml intradermally in left deltoid in newborns within 28 days -
- Efficacy of dose →0 to 80%
- Protective duration → 20 years

12. Nowadays which is the most common pasteurization method?

a) Holder (VAT) method
b) Ultra-high temperature method
c) High temperature and short time method
d) Low temperature and longtime method

- High temperature and short time is known as Flash Method 72 °C X 15 seconds.
- In rural areas Holder VAT method is most popular 63-66 •C for 30 mins.
- Phosphatase test- used to test the sufficiency of pressurized milk.

13. True about Quarantine:

a) Restriction of movement of Infected patient
b) Restriction of movement of healthy contact of an infectious disease
c) Movement restricted for shortest incubation period
d) Isolation of diseased person

	QUARANTINE	ISOLATION
Is done for	Healthy contact of a case with maximum Insulation period	Cases until recovery
Level of prevention	Primary	Secondary

- Quarantine was first done for the disease plague.

14. A 26 year old pregnant female visited the sub centre for her check-up. Which of the following test is done at Subcentre during pregnancy?

a) USG
b) Haemoglobin
c) OGTT
d) Triple test

Tests done at Sub center:

- Pregnancy detection test
- Hemoglobin level estimation
- Urine test for albumin and sugar

www.medicalcoupons.in

- Rapid malaria test

Tests done at PHC/CHC/DH:

- Blood grouping with Rh
- HIV testing
- VDRL, RPR
- Rapid Malaria test
- Blood Sugar
- HBSAg.

15. 25 year old patient with diarrheal episodes presented to the hospital. The physician started his treatment with ORS. In preparation of ORS sugar is added, Which of the following is the correct reason to add sugar in ORS?

a) Enhance acceptability
b) Enhance salt absorption
c)Enhance shelf life
d) Enhance taste

Adding sugar to ORS, enhances absorption of salt & water.
Sugar in the ORS has found to have sym- transporter mechanism with sodium.

Composition	Weight
Sodium chloride	2.6 gm
Potassium chloride	1.5 gm
Sodium citrate	2.9 gm
Glucose anhydrous	13.5 gm
Total	20.5 gms

Osmolar concentration	(mmol/liter)
Sodium	75
Potassium	20
Chloride	65
Citrate	10

www.medicalcoupons.in

Glucose anhydrous	75
Total	245

ReSoMAL: Rehydration solution for MALnourished
Sodium • halved to 45 mmol/L : Potassium Doubled to 40 mmol / L
SUPER ORS: Rice /starch/ Alanine based [not mono-sugars]

16. Vitamin which helps in Iron absorption is:

a) Vitamin A
b) Vitamin B
c) Vitamin C
d) Vitamin E

Decrease Iron absorption
Phytate (cereals, Pulses)
Tannates (Tea, Coffee)
Oxalates (green vegetables)
Calcium
Drugs like Tetracycline

- Iron absorption in India is < 10%
- Vitamin- C is the only facilitators of Iron absorption

17. Wernicke encephalopathy is due to deficiency of:

a) Thiamine
b) Biotin
c)Niacin
d) Hydroxycobalamin

- Thiamin (Vitamin B1) deficiency leads to Beri- Beri seen in polished rice eaters.
- In polished rice outer layer of rice is removed which contains Thiamine, it is more common in Andhra Pradesh & Telangana States.
- Thiamine deficiency also causes Wernicke's encephalopathy & Wernicke's Korakoff psychosis (seen in alcoholics)

18. Healthy volunteers are recruited in which phase of Clinical trials?

a) Phase1
b) Phase 2
c) Phase 3
d) Phase 4

Phase 1: Healthy volunteers are re elevated to establish safety & Non- toxicity of drugs.
Phase 2: Patients are used to record the efficiency of the drug.
Phase 3: Patients are used mainly for compassion of existing drugs.

Phase 4:

- Done on patients where post marketing surveillance of rare and long term side effects is evaluated.
- This is the longest phase of trail with minimum 10 to 25 years preferably till lifelong.

Phase 1 → tell the Maximum Tolerated Dose of the drug.
Phase 2 → Maximum Drug Failure occurs.
Phase 3 → Is a Randomized control trail
Phase O → Used before phase with healthy volunteers with 1/100 of the dose (micro dosing) with very less volunteers

19. A 29 year old male patient with abdominal pain, fatigue, dark colored urine, clay colored stools and yellow sclera presented in the hospital. On serological examination the patient is showing HBsAg positive and HBeAg positive. Which of the following is the correct clinical diagnosis ?

a) Acute viral hepatitis
b) Chronic viral hepatitis
c) Acute viral hepatitis with infectivity
d) Remote infection

HBs Ag →Epidemiologic marker (1st antigen to appears in infection)
HBc Ag →Rarely appears in plasma.
HBe Ag →Marker of infectivity, indicates & active viral replication.

Condition	Serological Pattern
Acute Hepatitis B	HBsAg, HBeAg, IgM Anti-HBC
Chronic Hepatitis B with active viral replication	HbsAg, HBeAg, IgG Anti-HBc
Chronic Hepatitis B, with low viral replication	HbsAg, IgG Anti-HBc, Anti-HBe
Recovery from Hepatitis B infection	Anti-HBS, IgG Anti-HBC
Post Vaccination	Anti-HBS

20. Which of the following are agents of Farmer's Lung?

a) Thermophilic Actinomycetes
b) Aspergillus fumigatus
c) Aspergillus flavus
d) Aspergillus niger

www.medicalcoupons.in

Farmers lung-exposure to mouldy hay contaminated with Thermophilic Actinomycetes/ Micropolyspora Faeni .
Aspergillus fumnigatus and flavus are opportunistic pathogens and common causes of infections in humans.
Pneumoconiosis occurs with particles of size 0.5- 3μ diameter.

21. Which of the following interfere with Iron absorption?

a) Vitamin C
b) Phytates
c)Hemoglobin
d) Myoglobin

Decrease Iron absorption :

- Phytate (cereals, Pulses)
- Tannates (Tea, Coffee)
- Oxalates (green vegetables)
- Calcium
- Drugs like Tetracycline

Iron absorption in India is <10%
Vitamin- C is the only facilitators of Iron absorption

22. Chronic malnutrition in children is best evaluated by?

a) Weight for height
b) Weight for age
c) Height for age
d) Ponderal index

Low weight for age → underweight →acute or chronic malnutrition
Low Height for age→ stunting → chronic malnutrition
Low weight for height → wasting → Acute malnutrition

23. A married couple visited to the gynaecology department complaining of not able to conceive even after year of regular unprotected coitus. The physician sent the male partner for semen analysis. According to WHO which of the following is the correct minimum sperm count for Normal Semen analysis

a) 2 million / mL
b) 15 million /mL
c)10 million / mL
d) 20 million ml

Latest semen analysis guidelines of 2010 says:

www.medicalcoupons.in

- Minimum sperm count- 15 million / ml
- Minimum volume - 1.5 ml
- Motility- > 40%
- Progressive motility- > 32%
- Vitality- > 58%
- PH-> 7.2
- Normal morphology- > 4%

24. Most common cause of Blindness in India:

a) Cataract
b) Refractive error
c)Trachoma
d) Glaucoma

- According to latest 2015-2019 survey in India, prevalence of blindness is about 0.32% among that most common cause contributing to this cataract (66.2%), reflective error contributes 20%
- Cataract is also the most common cause of visual impairment in India.
- Blindness is defined as < 3/60 in better eye with best possible duration.

PHYSIO JUNE 2019

1. What is the cause of dicrotic notch?

a) Passive filling of blood in ventricles
b) Rapid ejection phase
c) Closure of semilunar valves
d) Isovolumic contraction

The dicrotic notch is a small and brief increase in arterial blood pressure that appears when the aortic valve closes. The dicrotic notch would therefore be the result of a short period of backward flow of blood immediately before the aortic valve closes.

2. Which of the following is correct regarding Isovolumic relaxation?

a) AV valves are closed
b) Corresponds to QT interval
c)C wave of JVP
d) Semilunar valves open

ISOVOLUMETRIC RELAXATION:

www.medicalcoupons.in

Heart:
At the beginning of this phase the AV valves are closed.

Pressures & Volume:
Throughout this and the previous two phases, the atrium in diastole has been filling with blood on top of the closed AV valve, causing atrial pressure to rise gradually.
The "v" wave is due to the back flow of blood after it hits the closed AV valve. It is the second discernible wave of the jugular venous pulse.
The pressure in the ventricles continues to drop.
Ventricular volume is at a minimum and is ready to be filled again with blood.

Heart sounds:
The second heart sound (S2, "dup") occurs when the semilunar (aortic and pulmonary) valves close. S2 Is normally split because the aortic valve closes slightly earlier than the pulmonary valve.

3. **Adequate 0_2 oxygen delivery at cellular level occurs in which type of hypoxia?**

   a) Hypoxic
   b) Anaemic
   c)Stagnant
   d) Histotoxic

Histotoxic hypoxia (also called histotoxic hypoxia) is **the inability of cells to take up or use oxygen from the bloodstream**, despite physiologically normal delivery of oxygen to such cells and tissues but the oxygen delivery to tissue is normal.

4. **If radius of a vessel is doubled, then the blood flow is increased by?**

   a) 8 times
   b) 16 times
   c)32 times
   d) 256 times

Vessel resistance (R) is directly proportional to the length (L) of the vessel and the viscosity (n) of the blood, and inversely proportional to the radius to the fourth power (r^4)
So when radius is increased by 2 times blood flow decreases by 16 times.

5. **Normal anion gap is__ mmol/L?**

   a) 4-6
   b) 6-12
   c) 12-24
   d) 24-50

www.medicalcoupons.in

The cation normally measured is Nat, and the anions are usually Cl® and HCO3 The "anion gap" is the difference between unmeasured anions and unmeasured cations and is estimated as

PLASMA ANION GAP = $[Na^+] – [HCO_3^-] – [Cl^-]$ = 144 – 24 – 108= **12 mEq/L**
Normal range = 8-16 mEq/L

6. **Breathing pattern disorders occur when ventilation exceeds metabolic demands, resulting in symptom-producing hemodynamic and chemical changes. As well as having a marked effect on the biochemistry of the body, BPDs can influence emotions, circulation, digestive function as well as musculoskeletal structures involved in the respiratory process. Which of the following have prolonged inspiratory spasms that resemble breath holding?**

   1. Biot breathing
   2. **Apneustic breathing**
   3. Cheyne stokes breathing
   4. Kussmaul breathing

APNEUSTIC BREATHING → long inspiratory spasms (resembles breath holding)
CHEYNE-STROKE BREATHING → apnea followed by hyperapnea
KUSSMAUL RESPIRATION → increased depth and prolonged inspiration
REGULARLY IRREGULAR→ seen in meningitis

SURGERY JUNE 2019

7. **A patient presented to emergency after RTA with multiple rib fractures. He is conscious, speaking single words. On examination, respiratory rate was 40/min and BP was 90/40 mm Hg, What is the immediate next step?**

   a) Urgent IV fluid administration
   b) Intubate the patient
   c)Chest x-ray
   d) Insert needle in 2nd intercostal space

- In this case, Patient having tachypnea, dyspnea, hypotension
- After RTA, there is also rib fracture
- Above all are suggestive of tension pneumothorax
- It is an emergency condition, so wide bore needle is inserted in 2nd intercostal space

According to ATLS (2018):
In adults, wide bore needle is inserted in 5th intercostal space slightly anterior to mid axillary line
TOC: ICD insertion in 5th intercostal space in triangle of safety

www.medicalcoupons.in

8. Correct procedure of inserting Nasogastric tube is?

a) Supine with neck flexed
b) Supine with neck extended
c) Sitting with neck flexed
d) Sitting with neck extended

- Ask the patient to sit and flex the neck
- Lubricate the tube and the nasal cavity
- And then insert the tube

9. A 7-day old infant presents with bilious vomiting and gross abdominal distention with absent bowel sounds. X ray abdomen shows multiple gas filled loops. Diagnosis is?

a) Hirschsprung disease
b) Congenital Hypertrophic pyloric stenosis
c)Duodenal atresia
d) Malrotation of gut

Malrotation of gut:

- Surgical emergency
- Bilious vomiting - first sign of volvulus
- Abdominal distention
- Multiple gas filled loops

The cardinal symptom of intestinal obstruction in the newborn is bilious emesis. Proximal intestinal obstructions typically present with bilious emesis and minimal abdominal distension. X-ray shows a paucity or absence of bowel gas, which should normally be present throughout GIT within 24 hours
Distal obstructions typically present with bilious vomiting and abdominal distension. Passage of black green meconium should have occurred within the first 24 to 38 hours

X-ray calcification - complicated meconium ileus
Pneumatosis or pneumoperitoneum - necrotising enterocolitis
Contrast enema - microcolon - jejunoileal atresia or meconium ileus
No microcolon - Hirschsprung's disease, small left colon syndrome or meconium plug syndrome.

10. Newborn infant presents with abdominal distension and bilious vomiting. There is failure to pass meconium beyond 24 hours. X-ray shows dilated small bowel loops with no air fluid levels and soap bubble appearance in right lower quadrant. This condition is initially treated by:

www.medicalcoupons.in

a) Paul mikulicz ileostomy
b) Bishop ileostomy
c) Contrast enema
d) Barium enema

Infant presents with signs and symptoms of small bowel obstruction (abdominal distension, bilious vomiting and inability to pass meconium). X ray shows typical features, suggestive of meconium ileus
Uncomplicated meconium ileus is initially treated by contrast enema failure of which surgical management is advocated.

11. A full-term, 2.0 Kg male baby, delivered at home presented with excessive frothing within a few hours of birth. The antenatal period was unsupervised and the delivery, uncomplicated. At presentation, the baby had mild respiratory distress. The first investigation to be done in this case IS:

a) Bronchoscopy with injection of methylene blue
b) NG Tube insertion and CXR to check position of tube
c) CT chest
d) Endoscopy

Two problems
→ Collection of saliva in the mouth
→ Air coming inside the mouth
Therefore, saliva is not going into the esophagus
Esophageal atresia with tracheoesophageal fistula

Diagnosis:
- The inability to pass a NG into the stomach is a cardinal feature for the diagnosis of EA
- If gas is present in the GIT below the diaphragm, an associated TEF is confirmed
- Inability to pass a NG tube with absent radiographic evidence for gastrointestinal gas is virtually diagnostic of an isolated EA without TEF
- For H-type, tracheobronchoscopy + endoscopy should be performed

12. Calculate the GCS of a patient exhibiting eye opening on pain, conscious but confused and cannot tell time and exhibits flexion on painful noxious stimuli to the arm.

a) 8
b) 9
c) 10
d) 11

BEHAVIOR	RESPONSE	SCORE
	Spontaneously	4
	To speech	3

www.medicalcoupons.in

Eye opening response	To pain	2
	No response	1
Best verbal response	Oriented to time, place, and person	5
	Confused	4
	Inappropriate words	3
	Incomprehensible sounds	2
	No response	1
Best motor response	Obeys commands	6
	Moves to localized pain	5
	Flexion withdrawal from pain	4
	Abnormal flexion (decorticate)	3
	Abnormal extension (decerebrate)	2
	No response	1
Total score	Best response	15
	Comatose patient	8 or less
	Totally unresponsive	3

13. A 27-year-old woman presents with 24 weeks of gestation with thyroid lesion which is found to be papillary carcinoma of thyroid (PCT). Which is the best treatment for this patient?

a) Thyroid ablation using radioactive Iodine
b) Total thyroidectomy
c)Observation
d) Hemi-thyroidectomy

In this case patient is pregnant therefore total thyroidectomy is the treatment of choice

Treatment of PCT:
- Total thyroidectomy (TOC)
- Radioactive iodine (RAI)

Contraindications of RAI:
- Pregnancy
- Lactation

14. 36 yr old African-american male presents to plastic surgeon with recurrent keloid lesion in the left earlobe with a history of previous injuries. The lesion was operated on, but always grows back postoperatively. Most appropriate management of Recurrent Keloid is?

a) Excisional surgery
b) Intramarginal excision followed by radiation
c)Cryosurgery
d) Silicone Gel Sheeting

www.medicalcoupons.in

- Keloids rarely regress with time, often refractory to medical and surgical intervention
- First line treatment: silicone in combination with pressure therapy and intralesional corticosteroid injection
- Refractory cases (after 12 months of therapy): excision + post operative therapy (external beam or
- brachytherapy)
- New treatment modalities: internal cryotherapy and 5% imiquimod

15. Seat Belt injury leads to?

a) Splenic laceration
b) Splenic contusion
c) Mesenteric tear
d) Mesenteric adenitis

→ In Seat belt injury there is sudden deceleration → Mesenteric tear
→ Classical seat belt sign is abrasion involving neck, chest and abdomen
→ Most commonly injured structure is mesentery
→ Two types of tears are seen
- Longitudinal tear - can easily repair
- Transverse tear - resection and anastomosis

Mesenteric adenitis is a condition that affects the lymph nodes in the membrane which connects bowel to the abdominal walll (mesentry). A viral infection is the usual cause.

16. Which complication is seen after Varicose vein stripping procedure?

a) Neuralgia
b) Deep vein thrombosis
c)Acrocyanosis
d) Telangiectasis

- During stripping of long saphenous vein most commonly injured nerve is saphenous nerve (saphenous neuralgia)
- During stripping of short saphenous vein most commonly injured nerve is sural nerve
- Neuralgia is prevented by limit the stripping of long saphenous vein till knee joint

17. Which of the following is not scanned by FAST USG?

www.medicalcoupons.in

a) Pericardium
b) Pleural cavity
c)Spleen
d) Liver

FAST- Focussed assessment with sonography for trauma
It is performed in 2-4 minutes-**4 views (4 P) -**

- Pericardium
- Perihepatic region
- Peri splenic region
- Pelvis

e- FAST- Extended FAST- 6 views

- Pericardium
- Perihepatic region
- Peri splenic region
- Pelvis
- Right thoracic view
- Left thoracic view

18. A 40-year-old man is suffering from heaviness in scrotum. A bag of worms feel is observed on scrotal examination and the swelling is seen to reduce in supine position. What is the best treatment?

a) Suction drainage
b) Varicocelectomy
c)Jaboulay procedure
d) Herniotomy

VARICOCELE

- Bag of worms like sensation
- Dilated tortuous pampiniform plexus of vein
- Treatment- Varicocelectomy/ high inguinal ligation of pampiniform plexus of veins
- Jabuloy procedure is eversion of sac performed for median sized hydrocele
- Herniotomy is preferred for Hernia

19. A 9-year-old boy presented in the urology outpatient department with dysuria, intermittently interrupted urinary flow, and pain in the lower abdomen for one year. USG of abdomen detected a single urinary bladder stone of 14.6mm size. All the blood tests and routine examination of urine were within normal limits. There was no significant growth in urine culture preoperatively. Open cystolithotomy was done under general anesthesia. He was catheterized and a drain was placed for two days for drainage of retro pubic space. Post-operative dressing done and amoxicillin-clavulanic acid (375mg) was prescribed thrice daily for ten days. He was discharged

www.medicalcoupons.in

with a urinary catheter on the seventh post-op day and was advised to apply mupirocin ointment topically at the surgical site daily. But the surgical wound never healed properly and after six weeks patient presented with the discharge of pus from the incision site with lower abdominal pain and a burning sensation in the urethra. In this patient, the occurrence of SSI could have been reduced if Prophylactic antibiotics were given before?

a) 60 minutes before skin incision
b) 1-3 hours before skin incision
c)At time of surgical incision
d) Night before surgery for peaking of effect

Prophylactic antibiotic to minimize SSI are given 30- 60 minutes before skin incision / at the time of induction

20. Which procedure is done in case of recurrent Ranula management?

a) Incision and drainage
b) Aspiration
c) Excision
d) Sclerosant injection

→ Incision over ranula and edges are sutured- Marsupialization
→ In cases of recurrence- Excision

21. Which of the following is best for diagnosis of pheochromocytoma?

a) 24-hour urinary Vanillyl mandelic acid
b) 24-hour urinary Fractionated Metanephrine
c)24-hour Urinary Hydroxy indole acetic acid
d) 24-hour Urinary Hydroxy tryptamine

Pheochromocytoma

- Most sensitive screening test for pheochromocytoma- Urinary catecholamines and Vanillyl mandelic acid (VMA)level.
- Best test for diagnosis of pheochromocytoma- 24 hours plasma fractionated Metanephrine >24 hours urinary plasma fractionated Metanephrine
- 24 hours urinary hydroxy indole acetic acid is performed for carcinoid tumors

www.medicalcoupons.in

22. Identify the instrument in the image below ?



a) Artery forceps
b) Kocher forceps
c)Allis forceps
d) Babcock forceps

Artery forceps are used **to compress an artery to stem bleeding**, because of this they belong to a group of surgical instruments called hemostats.

23. Identify the image ?



a) Tongue depressor
b) Doyen retractor
c)Self-retaining retractor
d) Langenbeck's retractor

www.medicalcoupons.in

Tongue depressor

- Used in surgery department for evaluation of oral cavity
- Used in ENT department generally

24. What is the Diagnosis ?



a) Sebaceous cyst
b) Implantation dermoid
c)Angular dermoid
d) Lipoma

This image is a Swelling in the scalp with no punctum
Sebaceous cyst

- Generally sebaceous cyst has punctum
- But if present in scalp and scrotum it does not have punctum.

www.medicalcoupons.in

25. What is the diagnosis ?



a) Lipoma
b) Encephalocele
c)Cystic hygroma
d) Lymphadenopathy

Encephalocele-protrusion of brain and membranes covered by skin and due to Failure of neural tube to close completely

26. 40 year old male presented with complaints of dysphagia ,regurgitation and foul smelling mouth. On further evaluation he was diagnosed to have Zenker's diverticulum. This diverticulum is due to herniation of esophagus through the Killian's dehiscence which is located:

a) Below Superior constrictor
b) Between Inferior constrictor
c) Below cricopharyngeal muscle
d) Below upper 1/3rd of smooth muscle of oesophagus

Killian's dehiscence is weakness located between the inferior constrictor muscle

27. Which organ has highest chances of Graft rejection response?

a) Cornea
b) Gut
c)Liver
d) Skin

www.medicalcoupons.in

Graft vs Host disease

Maximum chance of Graft vs Host disease is GUT transplantation/small bowel transplantation because Donor's small bowel- presence of leucocytes in the lumen

28. In hypovolemic shock which organ should be assessed for determining under-perfusion?

a) Kidney
b) Heart
c)Lung
d) Liver

In hypovolemic shock, kidney has to be assessed for determining under perfusion
Best clinical indicator of tissue perfusion - Urine output

ORTHO JUN 2019

1. A 6-year-old child is suspected with supracondylar fracture of right hand, complaining of pain and swelling. X-ray of right elbow was not significant. What is the next best step in this case?

a) Compare with X-ray of left hand
b) Closed reduction and slab
c) Closed reduction with K wire fixation
d) Cast

Supracondylar humerus fracture:

Suspected elbow fracture in a child - Supracondylar humerus fracture
Undisplaced # of humerus → posterior fat is elevated → "Fat pad sign"
Fracture is displaced → distal fragments goes posteriorly → Extension
Common n/v involved - AIN (Anterior interosseous nerve) > Median > Radial> Ulnar (AMRU)

2. A patient presented with following serum parameters: Normal serum ALP normal PTH level and increased Ca+ and PO4. Most likely cause:

a) Vitamin D toxicity
b) Hyperparathyroidism
c) Osteoporosis
d) Osteomalacia


www.medicalcoupons.in

Bio-Chemical Changes			
	Osteoporosis	Osteomalacia	Hyperparathyroidism
Calcium	Normal	↓/ Normal	Increased
Phosphate	Normal	↑	Low (phosphateuria)
ALP	Normal	↑	-
PTH	-	↑	-

Hyper toxicity of Vit-d: ↑Ca &Phosphate levels

3. Three point bony relationship has diagnostic value in:

a) Elbow fracture
b) Monteggia fracture
c)Galeazzi fracture
d) Colles' fracture



www.medicalcoupons.in

4. Management of olecranon fracture as seen in the image is:



a) Above elbow plaster sla
b) Below elbow plaster sla b
c)Close reduction with Tension band wiring
d) Open reduction with Tension band wiring

Given X ray shows olecanon fracture (type 2)- Tension band wiring

TYPES OF OLECRANON FRACTURE:-

→ Type I : Crack without displacement of fragments.
→ Type I: Clean break with separation of fragments.
→ Type II. Comminuted fracture

TREATMENT:-

- Type I → above-elbow plaster slab in 30 degree of flexion
- Type II → open reduction and internal fixation using tension-band wiring
- Type III→ plating or tension-band wiring or excision of the fragments

www.medicalcoupons.in


5. The diagnosis of the given image



a) Scapho lunate instability
b) Lunate injury
c) Distal radio ulnar joint instability
d) Wrist dislocation

Given X ray shows scaphoid instability

- Scaphoid - Most common carpal bone to fracture
- Scapho-Lunate dissociation - Gap between scaphoid & lunate Terry Thomas sign / David letterman sign
- Terry-Thomas sign indicates scapholunate instability
- It denotes the increased gap between the scaphoid and the lunate.
- It results from tearing of the scapholunate ligament following fall on outstretched hand.
- A scapholunate gap of >3 mm is suggestive of instability.





The given x-ray shows the Terry-Thomas sign and the image shows comedian Terry-Thomas, after whom this sign has been named:

www.medicalcoupons.in

6. Diagnosis of Gout is confirmed by which test?

a) Serum Uric acid level
b) Synovial fluid analysis
c)Urine uric acid levels
d) X- ray changes

Synovial fluid test
Synovial fluid is a liquid found in your body's joints. This test looks at **whether you have urate crystals in the joint fluid**, which could mean you have gout. This is considered the most accurate test for diagnosing gout.

OPTHAL JUN 2019

1. Most common cause of Blindness in India

a) Cataract
b) Refractive error
c)Trachoma
d) Glaucoma

- Most common cause of blindness in India is cataract
- Refractive error is not a cause of blindness, it is ocular morbidity
- Trachoma is the most common infectious cause of blindness
- Glaucoma is second common cause of blindness in the world

2. In patient of head injury, who presents with headache and increased intracranial pressure, the effect on pupil is:

a) Ipsilateral mydriasis
b) Contralateral mydriasis
c) Ipsilateral miosis
d) Contralateral miosis

Effect on pupil is ipsilateral mydriasis.

- In this patient of head injury bleeding occurs, which increases Intra cranial pressure
- This increased intra cranial pressure compresses and damages the 3rd nerve
- 3rd nerve carry the parasympathetic fibers which is responsible for pupillary constriction
- So, the unopposed sympathetic fibers mediated by the 5th nerve dilate the pupil

› Hence patient comes with ipsilateral mydriasis

Hutchinson's pupil is an ipsilateral mydriasis due to 3rd cranial nerve compression by an intracranial mass lesion including intracranial bleeds.

www.medicalcoupons.in

3. Foster Kennedy syndrome is characterized by all of the following except
1. Ipsilateral optic atrophy
2. Contralateral anosmia
3. Contralateral papilledema
4.Contralateral central scotoma
5. Bilateral progressive optic neuropathy

a) 1,4,5 are correct
b) 3.4,5 are correct
c)1,3,4 are correct
d) 2,4,5 are correct

Foster-Kennedy syndrome.-

- It is associated with olfactory or sphenoidal meningioma and frontal lobe tumors.
- In this condition, there occurs pressure optic atrophy on the side of lesion and papilledema on the other side (due to raised intracranial pressure).
- Ipsilateral central scotoma
- Ipsilateral anosmia

4. A 42-year-old man presented to the emergency department with multiple knife wounds, one of which penetrated the right eye creating a 207mm full thickness laceration of the upper lid and a 12?mm superonasal full-thickness scleral laceration. The laceration extended radially back towards the optic nerve with prolapse of the intraocular contents. On admission, the patient underwent Evisceration of his right eye ball .Out of the given choices, Which layer of eyeball is removed during evisceration?

a) Middle and inner
b) Outer and middle
c)Outer and inner
d) All the layers of eyeball

Evisceration means Removal of the viscera


Evisceration
The contents of the eye are removed, but the outer layer of the eyeball (sclera) is left intact.

Enucleation
The entire eye, including the globe, is removed but the orbital contents are left in place.

Exenteration
The contents of the eye socket are removed, including the muscles, lacrimal gland, optic nerve and various bones of the orbit.

www.medicalcoupons.in

5. The following image shows ?



a) Bacterial keratitis
b) Fungal keratitis
c) Viral keratitis
d) Syphilitic keratitis

Viral keratitis showing dendritic ulcer caused by HSV

Fluorescein staining

- May reveal dendritic lesions (linear branching patterns with terminal bulbs)
- If the lesion enlarges, it can form a geographic ulcer

EPITHELIAL KERATITIS →DENDRITIC ULCER (Loss of corneal sensation) →GEOGRAPHICAL ULCER

Dendritic ulcer

- Finger like projections
- Feathery margins
- Satellite lesions may be seen
- Hypopyon

6. Drug contraindicated in keratitis

a) Tear drops
b) Steroids
c) Cycloplegics
d) Timolol

Drug contraindicated in corneal ulcer is steroids because it causes perforations.

www.medicalcoupons.in

TREATMENT IN CORNEAL ULCERS

Bacterial	Topical 4th generation fluoroquinolones (DOC) -Moxifloxacin -Gatifloxacin
Fungal	Topical Natamycin (DOC)
Viral	Topical Acyclovir / Famciclovir /Valacyclovir
Acanthamoeba	PHMB (Poly Hexa Methylene Biguanide) (DOC) -Propamidine -Chlorhexidine

7. Which of the following is seen retinitis pigmentosa?

a) Arteriolar attenuation
b) Neovascularization
c)Retinal artery thrombosis
d) Papilledema

The classic clinical triad of RP is **arteriolar attenuation**, retinal pigmentary changes (could be either hypopigmentation and/or hyperpigmentation in form of bone-spicule and pigment clumpings), and waxy disc pallor.

8. Not used for color vision testing:

a) Holmgren wool test
b) Ishihara chart
c)FM 100 Hue test
d) Ames test

Ames test is not used for color vision testing


Holmgren wool test
- It is a very ancient test used for color vision testing
- Not used anymore

Ishihara chart
- Most commonly used for color vision testing

FM 100 hue test
- Used for color vision testing

9. Where is the lesion that produces the visual field defect shown in the image ?

www.medicalcoupons.in



a) Craniopharyngioma
b) Meningioma
c)Pituitary adenoma
d) Temporal lobe lesion

SUPERIOR QUADRANTANOPIA / PIE IN THE SKY

- Inferior fibres are damaged
- Inferior fibres pass through temporal lobe
- ALL PIE IN THE SKY ARE SEEN IN OPPOSITE TEMPORAL LOBE LESIONS

LESIONS



10. Which of the following field defects occurs due to a lesion at the optic chiasma?

a) Bitemporal hemianopia
b) Unilateral blindness
c) Pie on the floor
d) Bilateral homonymous macular defect


BITEMPORAL HEMIANOPIA

- Nasal fibres gets damaged

www.medicalcoupons.in

- Caused by optic chiasmal lesion → pituitary adenoma (mc)

LESIONS



Disc/optic nerve lesions	Same sided monocular blindness
Chiasmal lesions	Bitemporal Hemianopia
Optic tract lesions (mc)	Incongruous Homonymous hemianopia
Lt. Geniculate body lesions	Opposite sided Homonymous hemianopia
Meyer's loop (Temporal lobe)	Pie in the sky
Baum's loop (parietal lobe)	Pie on the Floor
Occipital lobe lesion	Congruous Homonymous hemianopia with macular sparing

11. In which of the following condition there is ipsilateral 3rd nerve palsy with contralateral hemiplegia and facial palsy of upper motor neuron?

a) Weber syndrome
b) Terson syndrome
c) Millard-Gubler syndrome
d) Foville's syndrome

Weber's syndrome
- Ipsilateral 3rd nerve Palsy
- Contralateral Hemiplegia
- Facial palsy of upper motor neuron

12. All among the following are incorrect statements regarding lens induced glaucoma except :

www.medicalcoupons.in

a) Phacomorphic glaucoma is an open angle glaucoma
b) In phacolytic glaucoma lens capsule is intact.
c) Phacomorphic glaucoma has a deep anterior chamber
d) Mature cataract is an example of phacolytic glaucoma

In phacolytic glaucoma lens capsule is intact. There are micropores through which lens proteins leak.

Phacolytic glaucoma is caused by an inflammatory process caused by the leakage of lens material through the capsule of a mature cataract. The released lens material is composed of altered lens protein, macrophages, and other inflammatory cells that lead to trabecular meshwork obstruction and precipitate glaucoma.

13. A 28-year-old man reported within 6 weeks after sustaining penetrating accidental injury to his left eye with wooden stick On examination there is marked diminution of vision in the RE associated with pain, redness and photophobia for 1?week. The attending doctor diagnosed it as case of sympathetic ophthalmitis and the first sign he can see on a fundus examination is ?

a) Circumcorneal congestion
b) Hypopyon
c) Retrolental flare
d) Iris nodule

First sign in sympathetic ophthalmitis : retrolental flare

Sympathetic ophthalmitis (Pan uveitis):

- B/L granulomatous uveitis
- Caused by trauma to 1 eye
- Injured eye → Exciting eye
- other eye → Sympathizing eye
- Accommodation failure in good eye
- Earliest sign: Retrolental flare [vitreous flare]
- Mutton fat KP'S

www.medicalcoupons.in

DERMA JUNE 2019

1. **A 25-year-old female presents with history of fever and oral ulcers and has developed erythematous lesions on her face. Comment on the diagnosis.**



a) SLE
b) Dermatomyositis
c)Melasma
d) Rosacea

Erythematous lesions given in this image is a feature of SLE - Butterfly rashes

- Fixed erythema
- Naso labial folds are spared

Systemic Manifestation

- Fever
- Joint pain
- Oral ulcer
- Photosensitivity

Melasma

- Brownish
- Malar area and Nose

Rosacea

- Erythematous lesion + Telangiectasia
- Convex areas

Dermatomyositis → Heliotrope rash → Violaceous → eye

www.medicalcoupons.in

2. **The following lesion was noticed in a patient with history of involuntary weight loss. What is the diagnosis?**



a) **Acanthosis nigricans**
b) Leser-Trélat sign
c)Actinic keratosis
d) Intertriginous Candida

Acanthosis nigricans is a **skin condition that causes a dark discoloration in body folds and creases**. It typically affects the armpits, groin and neck. Acanthosis nigricans is a skin condition characterized by areas of dark, velvety discoloration in body folds and creases. The affected skin can become thickened

Causes

Benign → obesity (mc), Endocrine disorders
Malignant → Gastric adenocarcinoma

3. **Which is not correct about the lesion shown?**



www.medicalcoupons.in

a) The lesions are infectious to seronegative individuals
b) Can be associated with meningoencephalitis
c) Bilaterally symmetrical dermatomal vesicular eruption
d) Geniculate ganglion is involved in Ramsay Hunt syndrome

Fluid filled lesions grouped together along particular dermatome and unilateral which is seen in Herpes Zoster

Herpes zoster
Caused by Varicella zoster virus
Meningoencephalitis can be due to varicella zoster virus
In Ramsay hunt syndrome, painful grouped fluid filled lesions →vesicles →Herpes zoster oticus.

4. The shown instrument is:



a) Single cutting razor
b) Double cutting razor
c) Humby's knife
d) Finochietto knife

Humby's knife – In 3rd degree burn → used for skin grafting
Uses/ Indication:
1 . Burn patients
2. Vitiligo surgery

5. Most appropriate Management to reduce recurrence of keloid:

a) Intralesional steroids
b) Surgery
c)Cryotherapy
d) Electrocoagulation

www.medicalcoupons.in

SURGERY:

Treatment of keloid:

→ Silicon dressing
→ Mechanical compression
→ Intralesional steroids [Doc- keloids]

6. An infant presents infection. with the condition as shown below after a pharyngeal The shown condition is:



a) Staphylococcal scalded skin syndrome
b) Epidermolysis bullosa simplex
c)Impetigo contagiosa
d) Pemphigus foliaceus

Staphylococcal scalded skin syndrome (SSSS), also known as **Ritter von Ritterschein disease** (in newborns), Ritter disease, and staphylococcal epidermal necrolysis, encompasses a spectrum of superficial blistering skin disorders caused by the exfoliative toxins of some strains of Staphylococcus aureus.

7. The shown penile lesion is:



a) Genital wart
b) Pearly penile papule
c) Balanitis
d) Sebaceous gland prominence

www.medicalcoupons.in

Pearly papules :

→ Asymptomatic
→ Seen in uncircumcised patients
→ Etiology unknown
→ Not infective
→ Normal lesions
→ No treatment required

8. The show lesions are ?



a) Molluscum contagiosum
b) Chicken pox
c)Pemphigus
d) Common warts

Molluscum contagiosum is **a skin infection caused by the** virus Molluscum contagiosum. It produces benign raised bumps, or lesions, on the upper layers of your skin. The small bumps are usually painless. They disappear on their own and rarely leave scars when they're left untreated.

www.medicalcoupons.in


RADIO JUNE 2019

1. Comment on the diagnosis of CT chest shown below.




a) Aortic dissection
b) Pulmonary embolism
c)Cardiac myxoma
d) Aortic aneurysm

NORMAL CT



Aortic dissection



www.medicalcoupons.in

Aortic dissection:

- M/C site of aortic dissection is Ascending aorta
- Intimal flap with double lumen is seen
- IOC for acute aortic dissection
  → CT angiogram (stable pt.)
  → TEE (unstable pt.)
- In Chronic aortic dissection: IOC: MRI

2. **A 55 year old male patient presented with complaints of dyspnea with congestive heart failure . The clinician wants to know that whether its HF with preserved EF or reduced EF. Which of the below investigation is used for calculation of ejection fraction:**

   a) MUGA
   b) SPECT using thallium 201
   c)PET myocardial perfusion imaging
   d) Sestamibi scan with pharmacological stress

Nuclear Imaging of heart:

- MUGA scan (Multi-gated acquisition scan):
- Tc 99m labeled RBC IS used
- Evaluate ventricular function / ejection fraction- Cardiac ventriculography (but most accurate investigation for ventricular function is -MRI)
- Tc99m labeled RBC is very sensitive investigation for detecting minimal lower GIT bleeding (as low 0.1 ml per minute)

3. **Refer to the image given below and give the correct diagnosis?**



www.medicalcoupons.in

a) Diaphragmatic hernia
b) Intestinal obstruction
c)Pleural effusion
d) Pneumonia

Diaphragmatic hernia is **a birth defect where there is a hole in the diaphragm** (the large muscle that separates the chest from the abdomen). Organs in the abdomen (such as intestines, stomach, and liver) can move through the hole in the diaphragm and upwards into a baby's chest.


4. Maximum radiation exposure :

a) X-ray abdomen
b) Chest X ray
c)IV pyelography
d) Barium Enema


Barium Enema- Sequential/ multiple imaging, covers larger area. therefore comparatively maximum radiation exposure.

X-ray abdomen- Single spot film, therefore comparatively less radiation exposure.
Chest X ray- Single spot film, therefore comparatively less radiation exposure.
IV pyelography- Sequential/ multiple imaging but covers less area, therefore. comparatively less radiation
exposure.
PET CT - Has highest radiation exposure.


5. In the shown lateral view the structure marked with arrow head is?



a) Esophagus
b) Pulmonary artery
c) Trachea
d) Left atrium



6. A patient presented with complaints with dyspnea. The shown X- ray is suggestive of:-



a) Consolidation
b) Exudative Pleural Effusion
c)Pneumothorax
d) Hydropneumothorax

Chest X-ray with pleural effusion

- Obliteration of CP angle
- Distinct border
- Concave upward curve called

ELLIS CURVE

- 150-200 ml on PA view
- 5 ml lateral view

www.medicalcoupons.in

- 15-20 ml on lateral decubitus view
- Damoiseau-Ellis line is a clinical sign, that refers to **upper limit of dullness to percussion caused by** pleural effusion. It has characteristic shape of curved line with highest point at middle axillary line

ENT JUNE 2019

1. **Narrowest part of airway in infant:**

   a) Supraglottis
   b) Glottis
   c) Subglottis
   d) Epiglottis

Adult VS pediatric airway
In adult larynx is cylindrical and the narrowest part is glottic (vocal cords).
In case of paediatric larynx is funnel shape. And the narrowest part is subglottic.

2. **A 15 year old male patient presented with nasal mass reaching upto cheek and causing unilateral obstruction with intermittent epistaxis. Most likely diagnosis is:**

   a) Angiofibroma
   b) Nasal polyp
   c)Nasopharyngeal Carcinoma
   d) Inverted Papilloma

Angiofibroma is a **benign (noncancerous) nasal cavity tumor** that almost exclusively affects adolescent boys. It also may be referred to as juvenile nasopharyngeal angiofibroma (JNA).
Presentation
M/C- Profuse recurrent epistaxis
Nasal obstruction, check swelling.
O/E →Reddish polypoid mass
Diagnosis - CECT-Anterior bowing of posterior maxillary wall- ANTRAL/HOLMANN MILLER SIGN
→Biopsy is Highly contraindicated
RX → Endoscopic excision

3. **HIGH TRACHEOSTOMY is done in which of the following?**

www.medicalcoupons.in

a) Laryngeal papilloma
b) Carcinoma larynx
c)Bilateral vocal folds palsy
d) Laryngomalacia

Tracheostomy → Emergency/ Planned in hospital (between- 2nd and 3rd tracheal ring)
High tracheostomy is done in carcinoma larynx (Between 1st and 2nd tracheal Ring)

4. ALL ARE PARTS OF HYPOPHARYNX EXCEPT

a) Posterior pharyngeal wall
b) Pyriform Sinus
c)Post cricoid area
d) Vallecula

- Hypopharynx is also know laryngopharynx.
- Component

1. Posterior pharyngeal wall
2.Post cricoid area
3. Pyriform fossa

- Vallecula is part of oropharynx.It is space between epiglottis and Posterior 1/3rd of tongue

5. A 3 yrs old child presented with profound Sensorineural deafness, not responding to hearing aids, with intact auditory nerve function. Which of the following procedure is preferred in this patient?

a) Hearing Aids
b) BAHA
c) Cochlear implant
d) Auditory Brainstem Implant

INDICATIONS FOR COCHLEAR IMPLANTS

- Bilateral severe to profound H.L. that is not adequately treated with standard hearing aids.
- Congenital H.L./ prelingual deafness
- Acquired H.L./ post lingual deafness
- Severe hearing loss that can be aided and that deteriorates to profound loss in childhood, adolescence, or adulthood (prelingual).

6. A patient presented with 2 days history of fever and difficulty in swallowing. On examination there was a swelling seen in oropharynx pushing tonsil to midline and swelling over upper part of neck. What is the most likely diagnosis:

www.medicalcoupons.in

a) Quinsy
b) Parapharyngeal abscess
c)Retropharyngeal abscess
d) Tonsillitis

In case of parapharyngeal space abscess this abscess grow and form a swelling on lateral wall of oropharynx and it will also push the tonsil to midline.
Parapharyngeal space- lateral boundaries-
1 . Parotid gland (Deep lobe)
2. Ramus of mandible
3. Median pterygoid muscle
In case of parapharyngeal space abscess this abscess grow and form a swelling on lateral wall of oropharynx and it will also push the tonsil to midline.

- When question says there is swelling in oropharynx pushing tonsil to midline it doesn't means quinsy or tonsillitis
- Retropharyngeal abscess will be seen in midline

7. Deep lobe of parotid gland swelling presents at

a) Below the ear lobe
b) Cheek
c)Temporal lobe
d) Parapharyngeal Space

Parapharyngeal / lateral pharyngeal space Boundaries
- Superiorly - skull bone
- Lateral - deep lobe of parotid gland ,Ramus of mandible, Median pterygoid muscle
- Medial - oropharynx with superior constrictor muscle, Palatine tonsil
- Inferior- hyoid bone.

8. Which of the following muscles is the lifesaving muscle of larynx that pulls back vocal cords in order for breathing to take place

a) Cricothyroid
b) Posterior Cricoarytenoid
c) Lateral Cricoarytenoid
d) Transverse Arytenoid

Posterior cricoarytenoid muscle is regarded as Safety muscle of Larynx.
Posterior cricoarytenoid muscle of larynx is the only muscle which abducts true vocal cords and opens the rima glottidis. This muscle is supplied by recurrent laryngeal branch of vagus nerve. Injury to this nerve results in hoarsness of voice.

www.medicalcoupons.in

9. Rodent ulcers are which type of malignancy

a) Squamous cell carcinoma
b) Basal cell carcinoma
c)Malignant Melanoma
d) Adenocarcinoma

Rodent ulcers (basel cell carcinoma)- most common skin malignancy
Seen in - sun: exposed area (hand and face)(UV-B)
This is locally invasive and metastasis is rare.
Dx →wide local excision

10. What is the most common cause of deafness in a patient of Nasopharyngeal Carcinoma:

a) Metastasis to Temporal Bone
b) Middle ear Infection
c) Serous otitis media
d) Infiltration of Tumour middle ear

Nasopharyngeal carcinoma
Due to obstruction of eustachian tube there is serous otitis media and leads to hearing loss.
M/C site of origin → Fossa of Rosenmuller
M/C type → Squamous cell carcinoma
M/C Presentation →cervical lymphadenopathy (Neck Mass)

11. Most Common Presenting Complaint of Nasopharyngeal Carcinoma

a) Cervical lymphadenopathy
b) Hearing Loss
c) Airway Obstruction
d) Epistaxis

Nasopharyngeal carcinoma
Due to obstruction of eustachian tube there is serous otitis media and leads to hearing loss.
M/C site of origin → Fossa of Rosenmuller
M/C type → Squamous cell carcinoma
M/C Presentation →cervical lymphadenopathy (Neck Mass)

12. Structures not seen in indirect laryngoscopy?

a) Anterior Commissure
b) Posterior Commissure
c) Epiglottis
d) False Vocal Cord

Anterior commissure can't be seen because of bulge of anterior wall of larynx which hides the anterior commissure

13. Killian's Dehiscence is seen at the level of:

a) Superior Constrictor
b) Middle constrictor
c) Inferior Constrictor
d) Thyroepiglottis

The Killian dehiscence is a triangular-shaped area of weakness in the muscular wall of the pharynx, **between the transverse and oblique bundles of the inferior pharyngeal constrictor**. It is the most common site of pharyngeal diverticula.

- **Killian dehiscence also known as "Gateway of tears"**
- M/C site for esophageal tear (perforation)

14. Earliest symptoms of vocal fold growth is?

a) Cough
b) Change of voice
c) Fatigue voice
d) Pain

Vocal fold growth earliest symptom is change of voice or hoarseness.

15. Parotid gland is supplied by which nerve?

a) Facial Nerve
b) Glossopharyngeal
c) Vagus Nerve
d) Hypoglossal Nerve

Nerve supply to parotid gland- Parasympathetic innervation to the parotid comes from the glossopharyngeal nerve (CNIX)and causes secretion of saliva. Parasympathetic innervation to the submandibular glands is provided by the superior salivatory nucleus via the chorda tympani, a branch of the facial nerve, that becomes part of the trigeminal nerve's lingual nerve prior to synapsing on the submandibular ganglion.

www.medicalcoupons.in

FMGE DEC 2019

ANATOMY DEC 2019

1. Which of the following is the safety muscle of tongue?

a) Genioglossus
b) Hyoglossus
c) Palatoglossus
d) Styloglossus

The genioglossus arises from the mandible and protrudes the tongue. It is also known as the tongue's "safety muscle" since it is the only muscle that propels the tongue forward. The hyoglossus, arises from the hyoid bone and retracts and depresses the tongue.

2. Trigone of bladder is derived from

a) Distal part of Paramesonephric duct
b) Distal part of Mesonephric duct
c) Ventral part of Mesonephric duct
d) Ventral part of Urogenital sinus

Embryologically, the trigone of the bladder is derived from **the caudal end of mesonephric ducts**, which is of mesodermal origin (the rest of the bladder is endodermal).


3. Identify the marked structure;



a) Anterior superior illac spine
b) Anterior inferior iliac spine
c) Iscial tuberosity
d) Pubic symphysis

www.medicalcoupons.in

The anterior superior iliac spine is an important bony surface landmark and is **the prominence is the most anterior part of the ilium**. It can be palpated at the lateral end of the inguinal fold. ... It is also the start point for the measurement of true leg length (anterior superior iliac spine to medial malleolus).



4. Which muscle tendon is stretched in patellar tendon reflex?

a) Biceps femoris
b) Quadriceps femoris
c) Addactur magnus
d) Quadratus

The knee-jerk reflex, also known as the patellar reflex, is a simple reflex that causes the contraction of **the quadriceps muscle** when the patellar tendon is stretched.

www.medicalcoupons.in

5. Celiac artery is seen at which vertebral level?

a) T9
b) T10
c) T12
d) L2

The celiac trunk, also known as the celiac artery, is a short vessel that arises from the aorta and passes below the median arcuate ligament, just as the aorta enters the abdomen at the level of the **T12 vertebra**.

6. Highest point of iliac crest is seen at;

a) L3
b) L4
c) S1
d) S2

The top of the iliac crests also marks the level of the **fourth lumbar vertebral body (L4)**, above or below which lumbar puncture may be performed.

7. 45-year-old policeman comes from duty with complain of pain in right leg. On examination, pulse in right popliteal fossa was absent and pain in right buttock, whereas in left side it was normal. The block is at which level?

a) Femoral
b) Iliac
c) Aortoiliac
d) Popliteal

Aortoiliac occlusive disease is a **type of vein condition in the legs**. It occurs when the iliac artery that brings blood to your legs becomes narrow or blocked by plaque. Symptoms can include pain, numbness, or cramping in the lower limbs, gangrene in the feet, and erectile dysfunction (ED) in men.

8. Lacrimal gland is supplied through parasympathetic system. It is supplied by which ganglion?

a) Ciliary ganglia
b) Otic ganglia
c) Pterygopalatine
d) Submandibular

www.medicalcoupons.in

The postganglionic axons are the output of **the pterygopalatine ganglion** project to the lacrimal gland and supply the parasympathetic innervation of the lacrimal gland

9. The structure which doesn't pass through the parotid gland?

a) Facial nerve
b) ICA
c) Branch of ECA
d) Retromandibular vein

The facial nerve and its branches pass through the parotid gland, as does the **external carotid artery and retromandibular vein**. The external carotid artery forms its two terminal branches within the parotid gland, maxillary and superficial temporal artery

10. Protraction of neck is due to which of the following muscle?

a) Serratus anterior
b) Pectoralis major
c) Pectoralis minor
d) Lateral pterygoid

11. Middle meningeal artery is branch of;

a) Maxillary artery
b) Mandibular
c) Facial artery
d) Ophthalmic artery

Middle meningeal artery primarily is derived as a branch from **the maxillary artery** and then passes through the foramen spinosum into the middle cranial fossa

Five branches, each of which enters a bony foramen:

- deep auricular artery (enters squamotympanic fissure)
- anterior tympanic artery (enters squamotympanic fissure)
- middle meningeal artery (enters foramen spinosum)
- accessory meningeal artery (enters foramen ovale)
- inferior alveolar artery (enters mandibular foramen)


www.medicalcoupons.in

12. The muscle which is required to open Eustachian tube;


a) Tensor veli palatini muscle
b) Salpingo pharengeous
c) Tensor tympani
d) Levator veli palatini

The eustachian tube opens upon swallowing or yawning by contraction of **the tensor veli palatini muscle**.

13. The highlighted muscle is supplied by which nerve?




a) Radial nerve
b) Ulnar nerve
c) Median nerve
d) Musculocutaneous nerve



www.medicalcoupons.in


The musculocutaneous nerve innervates the three muscles of the anterior compartment of the arm: the **coracobrachialis, biceps brachii, and brachialis**. It is also responsible for cutaneous innervation of the lateral forearm.

14. Identify the encircled structure;



a) Pterion
b) Glabella
c) Asterion
d) Nasion

The pterion is the **H-shaped formation of sutures on the side of the calvarium** representing the junction of four skull bones: the greater wing of the sphenoid bone, squamous portion of the temporal bone, frontal bone, parietal bone.




15. What is the name of marked artery?



www.medicalcoupons.in

a) **Posterior Tibial Artery**
b) Deep Phalangeal artery
c) Lateral planter
d) Medial planter



The posterior tibial artery (PTA) is **one of the 2 branches of the tibioperoneal trunk in the lower leg** and provides oxygenated blood to the posterior compartments of the leg and plantar surface of the foot. It is accompanied by the posterior tibial vein, along its course.

PATHO DEC 2019

16. Most common tumour of mediastinum?

a) Thymoma
b) Lymphoma
c) **Neurogenic tumor**
d) Neuroblastic tumor

The most common mediastinal masses are **neurogenic tumors** (20% of mediastinal tumors), usually found in the posterior mediastinum, followed by thymoma (15–20%) located in the anterior mediastinum. Lung cancer typically spreads to the lymph nodes in the mediastinum.

17. Most common type of Embryonic neural tumor;

a) Fibrous astrocytoma
b) Neuroglioma
c) Ependymoma
d) **Medulloblastoma**

www.medicalcoupons.in

Medulloblastomas. The most common type of embryonal tumor, these **fast growing cancerous** brain tumors start in the lower back part of the brain, called the cerebellum. The cerebellum is involved in muscle coordination, balance and movement.

18. Which of the following is not a component of SIRS (Systemic Inflammatory Response Syndrome)?

a) Fever >38 degrees or hypothermia < 36 degree,
b) Tachycardia >90 beats /min
c) Tachypnea>25 breaths / min
d) leucocytosis >12*10^9/l or leucopoenia <4*10^9/l

Systemic Inflammatory Response Syndrome (SIRS) is the occurrence of at least two of the following criteria:

→ fever >38.0°C or hypothermia <36.0°C,
→ tachycardia >90 beats/minute,
→ tachypnea >20 breaths/minute,
→ **leucocytosis** >12*10^9/l or leucopoenia <4*10^9/l

19. Which of the following has Autosomal Recessive inheritance?

a) Hereditary spherocytosis
b) Sickle cell anaemia
c) Osteogenesis imperfecta
d) VWD

Autosomal Recessive inheritance Examples:

- Albinism (oculocutaneous)
- Ataxia telangiectasia
- Crigler-Najjar syndrome type
- Congenital adrenal hyperplasia
- Cystic fibrosis
- Deafness (some forms)
- Dubin-Johnson syndrome
- Epidermolysis bullosa (some forms)
- Fanconi syndrome
- Friedreich's ataxia
- Galactosaemia
- Gaucher's disease
- Glycogen storage disease
- Haemochromatosis
- Homocystinuria

www.medicalcoupons.in

- Hurler's syndrome (mucopolysaccharidosis I)
- Infantile polycystic kidney disease
- Laurence-Moon-Biedl syndrome
- Phenylketonuria
- Sickle cell disease
- Tay-Sachs disease
- B-Thalassemia
- Wilson's disease

20. Which of the following is an Autosomal Dominant disease?

a) Von Willebrand disease
b) Wilson's disease
c) Cystic fibrosis
d) Friedrichs ataxia

Mnemonic for Autosomal Dominant disorders: Very Powerful **DOMINANT** Humans

- **V**on willebrand disease / **V**on hippel-lindau
- **P**seudo-hypoparathyroidism
- **D**ystrophia myotonica
- **O**steogenesis imperfecta / **O**sler-weber-rendu
- **M**arfan syndrome
- **I**ntermittent porphyria
- **N**eurofibromatosis
- **A**chondroplasia / **A**dult polycystic kidney disease
- **N**oonan syndrome
- **T**uberous sclerosis
- **H**ypercholesterolemia
- **H**untington's disease
- **H**ypertrophic obstructive cardiomyopathy
- **H**ereditary spherocytosis
- **H**ereditary non polyposis coli
- **H**ereditary hemorrhagic telangiectasia

21. A patient has MCV < 80, MCH < 23. Which type of anemia shall be classified?

a) Normocytic hypochromic
b) Microcytic hypochromic
c) Hyperchromic macrocytic
d) Normocytic normochromic

www.medicalcoupons.in

- Erythrocytes that have a normal size or volume (normal MCV) are called **normocytic**,
- When the MCV is high, they are called **macrocytic.**
- When the MCV is low, they are termed **microcytic.**
- Erythrocytes containing the normal amount of hemoglobin (normal MCHC) are called **normochromic.**
- When the MCHC is abnormally low they are called **hypochromic**, and when the MCHC abnormally high, **hyperchromic.**

	MCV	MCHC
Female:	80-95 fl	30-34 gHb/100ml
Male:	80-95 fl	30-34 gHb/100ml

22. Which among the following ions is important in cell injury?

a) **Calcium**
b) Sodium
c) Potassium
d) Chloride

Calcium ions have been proposed as mediators of the functional consequences of plasma membrane injury. An influx of extracellular calcium ions across a damaged permeability barrier and down a steep concentration gradient may convert potentially reversible injury into irreversible injury.

23. Which type of anaemia is seen in vitamin B12 deficiency?

a) Microcytic, hypochromic
b) Macrocytic Hypochromic
c) Normocytic normochromic
d) **Macrocytic, Normochromic**

Macrocytic - Normochromic Anemias

- **Pernicious anemia**

→ Caused by a lack of intrinsic factor from the gastric parietal cells
→ Required for vitamin B12 absorption
→ Results in **vitamin B12 deficiency**

www.medicalcoupons.in

24. Fertilized egg cells are;

a) Totipotent
b) Pluripotent
c) Multipotent
d) None of above

Totipotent stem cells can differentiate into embryonic and extra-embryonic cell types. These cells are produced from the fusion of an egg and sperm cell, and can construct a complete, viable organism. Cells produced by the first few divisions of the fertilized egg are also totipotent.

25. 16-year-old girl with short height, amenorrhea with widely spaced nipple. karyotyping is;

a) 45 X0
b) 46 XO
c) 47 XXX
d) 46XY

Turner syndrome (TS), also known as 45 X , or 45XO , is a genetic condition in which a female is partly or completely missing an X chromosome. Turner syndrome, a condition that affects only females, can cause a variety of medical and developmental problems, including short height, failure of the ovaries to develop and heart defects.

Signs of Turner syndrome at birth or during infancy may include:

- Wide or web like neck
- Low-set ears
- Broad chest with widely spaced nipples
- High, narrow roof of the mouth (palate)
- Arms that turn outward at the elbows
- Fingernails and toenails that are narrow and turned upward
- Swelling of the hands and feet, especially at birth
- Slightly smaller than average height at birth
- Slowed growth
- Cardiac defects
- Low hairline at the back of the head
- Receding or small lower jaw
- Short fingers and toes

www.medicalcoupons.in

26. Which type of necrosis is seen in brain?

a) Coagulative
b) Liquefactive
c) Fat
d) Fibrinoid

Type of Necrosis	Site
Coagulative Necrosis	Heart, spleen & kidney
Liquefactive (or) Colliquative necrosis	Brain, abscess cavity
Fat necrosis	Pancreas, breast
Fibrinoid necrosis	Arteries in hypertension, peptic ulcer, etc.
Caseous necrosis	In the centre of foci of tuberculus infection

27. Following heart sound is almost always pathological;

a) Loud S1
b) Loud S2
c) Systolic murmur
d) S4

A S4 heart sound is almost always pathologic. Heart sounds can be described by their intensity, pitch, location, quality and timing in the cardiac cycle. Intensity: Heart sounds can be described as increased in intensity (loud), decreased in intensity (soft) or absent.

SPM DEC 2019

28. Koplik spots are seen in;

a) Mumps
b) Measles
c) Rubella
d) Varicella

Koplik spots are seen with **measles**. They are small, white spots (often on a reddened background) that occur on the inside of the cheeks early in the course of measles.

www.medicalcoupons.in

29. NITI AAYOG Represents;

a) Nutritional institute for transforming India
b) National Institute for Transforming India
c) Newer Initiative transforming India
d) Nutrition Intake to India

National Institution for Transforming India, better known as NITI Aayog, was formed via a resolution of the Union Cabinet on 1 January 2015. NITI Aayog is the premier policy think tank of the Government of India, providing directional and policy inputs.

30. On republic day, a camp was organized and people were screened for Hypertension by checking BP and for diabetes by checking their BMI and Blood sugar level, which level of prevention is this?

a) Primary
b) Primordial
c) Secondary
d) Tertiary

There are 3 levels of disease prevention:
Primary Prevention - trying to prevent yourself from getting a disease.
Secondary Prevention - **trying to detect a disease early and prevent it from getting worse**.
Tertiary Prevention - trying to improve your quality of life and reduce the symptoms of a disease you already have.

31. A person comes with right eye 6/60 and left eye 3/60. He shall be categorized into which type of blindness?

a) Manifest
b) Social
c) Economical
d) No blindness

Types of Blindness:


- **Economic blindness:** Inability of a person to count fingers from a distance of 6 meters or 20 feet .
- **Social blindness:** Vision 3/60 or diminution of field of vision to 10°
- **Manifest blindness:** Vision 1/60 to just perception of light
- **Absolute blindness:** No perception of light
- **Curable blindness:** That stage of blindness where the damage is reversible by prompt management e.g. cataract

www.medicalcoupons.in

- **Preventable blindness:** The loss of blindness that could have been completely prevented by institution of effective preventive or prophylactic measures e.g. xerophthalmia, trachoma and glaucoma
- **Avoidable blindness:** The sum total of preventable or curable blindness is often referred to as avoidable blindness.

32. Recycling, tubing and recyclable catheters will be disposed in which bag according to biomedical waste management?

a) **Red**
b) Blue
c) Yellow
d) White



33. Main focus of UNICEF is on;

a) **Child health**
b) Nutritional health
c) Mental health
d) Social health

UNICEF is mandated by the United Nations General Assembly to **advocate for the protection of children's rights**, to help meet their basic needs and to expand their opportunities to reach their full potential.

34. Subacute Sclerosing Panencephalitis is a rare and dangerous complication of;

www.medicalcoupons.in

a) Measles
b) Mumps
c) Rubella
d) Varicella

Subacute sclerosing panencephalitis (SSPE) is a very rare, but fatal disease of **the central nervous system** that results from a **measles** virus infection acquired earlier in life. SSPE generally develops 7 to 10 years after a person has measles, even though the person seems to have fully recovered from the illness.

35. Spermicidal cream which is used in contraceptives;

a) Gossypol
b) Nonoxynol-9
c) Centchroman
d) Clomiphene

Nonoxynol-9 is the primary chemical in spermicides to inhibit sperm motility. Active secondary spermicidal ingredients can include **octoxynol-9, benzalkonium chloride and menfegol**. The most effective strength spermicide contains **at least 100 mg of nonoxynol-9 per dose**.

36. Kyoto Protocol is for;

a) Water sanitization
b) Food and nutrition
c) To reduce greenhouse gas emission
d) Water chlorination

Kyoto Protocol operationalizes **the United Nations Framework Convention on Climate Change by committing industrialized countries and economies in transition to limit and reduce greenhouse gases (GHG) emissions in accordance with agreed individual** targets.

37. Sustainable Development Goals is Designed by;

a) UNICEF
b) United Nations
c) WHO
d) Government of India

The Sustainable Development Goals (SDGs) or Global Goals are a collection of 17 interlinked global goals designed to be a "blueprint to achieve a better and more

www.medicalcoupons.in

sustainable future for all". The SDGs were set up in 2015 by **the United Nations General Assembly** and are intended to be achieved by the year 2030.

38. Which of the following is the chemoprophylaxis of choice in a person who is on a journey to endemic malarial region?

a) Chloroquine
b) Quinine
c) Lumefantrine
d) Pyrimethamine

Atovaquone/proguanil (Malarone), doxycycline, and mefloquine are the drugs of choice for malaria prevention in most malaria-endemic regions. **Chloroquine** (Aralen) may be used safely in all trimesters of pregnancy, and mefloquine may be used safely in the second and third trimesters of pregnancy.

39. Worms, bacteria and bugs attached in clothing of a person. This is;

a) Infection
b) Infestation
c) Contamination
d) Contagion

contamination is the act or process of contaminating; pollution; defilement; taint; also, that which contaminate while **infestation** is the presence of a large number of pest organisms in an area of field, on the surface of a host or anything that might contact a host, or in the soil.

40. Target couple is;

a) Couple with 3 children
b) Couple using contraception
c) Currently married couple where the wife is in reproductive age (15-49 year)
d) Couple that is eligible for practicing family planning

Target couples :
The term target couples was applied to couples who have had 2-3 living children, or even newly married couples to make them accept the idea of family planning from the earliest possible stage.

41. Yellow fever vaccine is valid for;

a) Life long
b) 10 years
c) 5 years
d) 20 years

www.medicalcoupons.in

As of 11 July 2016, a certificate of vaccination against yellow fever (ICVP) is valid for **the lifetime of the person vaccinated**. Prior to 2016, yellow fever vaccination certificates were valid for a period of 10 years, commencing 10 days after the date of vaccination.

42. In Africa, vaccination was given in an endemic outburst area for a specific disease but due to shortage of vaccines, mass chemoprophylaxis treatment was given to people who were not diseased. Which type of prevention is this?

a) Specific protection
b) Health promotion
c) Early diagnosis and treatment
d) Rehabilitation

SPECIFIC PROTECTION:
Idea is to avoid disease altogether. Currently available interventions include:

- Immunization
- Use of specific nutrients
- Chemoprophylaxis
- Protection against occupational hazards
- Protection against accidents
- Protection from carcinogens
- Avoidance of allergen
- Better environmental conditions e.g. control of air and noise pollution
- Control of consumer product quality and safety of foods, drugs, cosmetics.

43. Patients are categorized on the basis of chances of survival in Disaster management.

a) Triage
b) Tagging
c) Mitigation
d) Surge capacity

Triage is the process of quickly examining sick or injured people, for example after an accident or a battle, so that those who are in the most serious condition can be treated first.

www.medicalcoupons.in

44. Vision 2020 - which was added recently in India?

a) Cataract
b) Childhood blindness
c) Glaucoma
d) Refractive errors and low vision

Glaucoma is now regarded as a sort of "missing link" and VISION 2020 India successfully advocated for the inclusion of glaucoma in the next phase by the Indian government.

45. Diseases spread by the shown vector is all except;



a) Kala azar
b) Oroya Fever
c) Chagas disease
d) Cutaneous leishmania

Sandfly (Phlebotomus) Diseases Transmitted

- Leishmaniasis
- Kala Azar
- Oriental Sore (Cutaneous Leishmaniasis)
- Sandfly fever

46. SARS is caused by novel variant of;

a) Pox virus
b) Toga virus
c) Coronavirus
d) Lyssa virus

Corona viruses are a large family of viruses which may cause illness in animals or humans. In humans, several coronaviruses are known to cause respiratory infections ranging from the common cold to more severe diseases such as Middle East Respiratory Syndrome (MERS) and Severe Acute Respiratory Syndrome (SARS)

www.medicalcoupons.in

47. N95 respirator is used for?

a) Airborne Precautions
b) Droplet Precautions
c) Contact Precautions
d) None of the Above

N95 respirators and surgical masks are examples of **personal protective equipment** that are used to protect the wearer from **Airborne particles** or from liquid contaminating the face.

48. Which of the following is not a measure of dispersion?

a) Mean
b) Range
c) Variance
d) Standard deviation

Measures of dispersion describe the spread of the data. They include the **range, interquartile range, standard deviation and variance**. The range is given as the smallest and largest observations.

49. Anthracosis is caused by;

a) Silica
b) Asbestos
c) Coal dust
d) Sugar cane waste

Anthracosis is defined in Dioline as, "the asymptomatic, milder type of pneumoconiosis as caused by **the accumulation of carbon in the lungs due to repeated exposure to air pollution or inhalation of smoke or coal dust particles**".

50. Breast feeding should be continued for minimum how many months?

a) 6 months
b) 3 months
c) 12 month
d) 5 months

The World Health Organization recommends that all babies be exclusively breastfed for **6 months**, then gradually introduced to appropriate foods after 6 months while continuing to breastfeed for 2 years or beyond. Stopping breastfeeding is called weaning.

www.medicalcoupons.in

51. Components of Kuppuswamy scale-

a) Housing, Livestock, land
b) Income, Occupation, Education
c) Income, Housing, Education
d) Income, Livestock, Housing


The modified Kuppuswamy scale is commonly used to measure SES in urban and rural areas. This scale was devised by Kuppuswamy in 1976 and consists of a **composite score which includes the education and occupation of the Family Head along with income per month of the family**, which yields a score of 3–29.

52. Mode of transmission to bacteria through bacteriophage is by?

a) Transformation
b) Transduction
c) Translation
d) Conjugation


The transfer of genes from one bacterial strain to another by a bacteriophage is called **transduction** and can occur in a generalized or specific manner.

53. Identify the parasite as shown in photo;



a) Fasciola hepatica
b) Schistosoma japonicum
c) Schistosoma mansoni
d) Schistosoma hematobium

www.medicalcoupons.in



54. A patient presented with following findings what will be the media used to diagnose this condition?



a) **Loeffler's serum slope**
b) Lownstein Johnson
c) Karry blair
d) Chocolate agar

Loeffler's serum slope:
The medium contains horse serum, beef extract, dextrose, and proteose peptones which together supply the complex nitrogenous substances and nutrients necessary to support the growth of Corynebacterium diphtheriae.
Diphtheria destroy healthy tissues in the respiratory system. The **dead tissue forms a thick, gray coating** that can build up in the throat or nose. This thick gray coating is called a **"pseudomembrane"**

55. Identify the parasite as shown in this photo;

www.medicalcoupons.in



a) Brugia Malayi
b) Wucheria bancrofti
c) Tinea solium
d) Tinea saginatum

Brugia malayi is a filarial (arthropod-borne) nematode (roundworm), one of the three causative agents of lymphatic filariasis in humans. Lymphatic filariasis, also known as elephantiasis, is a condition characterized by swelling of the lower limbs. The two other filarial causes of lymphatic filariasis are Wuchereria bancrofti and Brugia timori, which both differ from B.malayi morphologically, symptomatically, and in geographical extent.

56. Most sensitive test for microfilariae;

a) DEC challenge test
b) Immunofluroscence test
c) Thick films
d) Membrane filter concentration technique

Membrane filtration and **the Knott concentration method** are more sensitive than routine thick blood smears for microfilaria detection, because they can be used to concentrate microfilariae from 1 mL or more of blood.

57. What is the causative agent for malignant otitis externa?

a) Pseudomonas
b) Staph aureus
c) Streptococcus
d) Influenza

www.medicalcoupons.in

Malignant external otitis (MEO) is an infection that affects the external auditory canal and temporal bone. The causative organism is usually **Pseudomonas aeruginosa**, and the disease commonly manifests in elderly patients with diabetes

58. Smallest cestode among the following;

a) H. nana
b) Schistosoma
c) T. saginata
d) T. Solium


Hymenolepis nana, the dwarf tapeworm, is the smallest and a common tapeworm in humans worldwide. H. nana infection occurs more frequently in warm climates and temperate zones such as Asia, Central and South America, and Eastern Europe.

59. Which antibody is seen in recent infection?

a) IgM
b) IgG
c) IgE
d) IgD

Immunoglobulin M (IgM) is one of several isotypes of antibody (also known as immunoglobulin) that are produced by vertebrates. IgM is **the largest antibody**, and it is the first antibody to appear in the response to initial exposure to an antigen.

60. A patient who was recently got dentature fixed, found with oral thrush. It bleeds on scraping. Causative agent for this condition



a) Diphtheria
b) Candida
c) Strep Mutans
d) Staph Aureus

www.medicalcoupons.in

Oral thrush (mouth thrush)

- Also called: oral candidiasis
- An infection in which the fungus Candida albicans accumulates in the mouth.
- It's common in babies, people with immune deficiency and those who use steroid sprays for asthma.
- Oral thrush causes white lesions on the tongue or inner cheeks
- Unsweetened yoghurt or an anti-fungal medication can help.

61. Bacterial vaginosis is caused by all except-

a) Gardenerella
b) Mobilincus
c) Proteus
d) Coccobaccillus

Proteus mirabilis is a common pathogen responsible for **complicated urinary tract infections (UTIs)** that sometimes cause bacteremia. Most cases of P. mirabilis bacteremia originate from a UTI; however, the risk factors for bacteremia and mortality rates from P. mirabilis UTI have not been determined. Proteus species also **produce urease**, which has been shown to be associated with an increased risk of pyelonephritis and upper UTIs. Proteus species also hydrolyze urea to ammonia, thereby alkalinizing the urine. Through the production of urease and ammonia, Proteus can produce an environment where it can survive
Whereas BV is linked to **an imbalance of "good" and "harmful" bacteria that are normally found in a woman's vagina**. Having a new sex partner or multiple sex partners, as well as douching, can upset the balance of bacteria in the vagina. This places a woman at increased risk for getting BV.

62. Kala azar cannot occur at an altitude of;

a) 400 meters
b) 100 meters
c) 600 meters
d) 200 meters

A disease caused by infection with Leishmania parasites.
Visceral leishmaniasis is spread by Sandfly bites. This type of leishmaniasis affects the internal organs, usually the spleen, liver and bone marrow.
Some people have no symptoms. For others, symptoms may include fever, weight loss and swelling of the spleen or liver.
Medication exists to kill the parasites. If left untreated, severe cases are typically fatal.
Kala-azar or Visceral leishmaniasis (VL) is a disease of low altitude (approximately **500 meters** mean sea level)

63. A bacterium which can sustain >100 degree comes is classified as;

a) Thermophilic
b) Mesophilic
c) Cycrophilic
d) Halophilic


A thermophile is **an organism—a type of extremophile—**that thrives at relatively high temperatures, between 41 and 122 °C (106 and 252 °F). Many thermophiles are archaea, though they can be bacteria. ... The enzymes in thermophiles function at high temperatures.

64. The causative agent for duodenal ulcer is;

a) Shigella
b) E.Coli
c) H. Pylori
d) Lactobacilli

Helicobacter pylori infection is the most common cause of duodenal ulcer disease, yet duodenal ulcer is an uncommon outcome of H. pylori infection


65. Identify the microorganism;



a) Influenza virus
b) Corona virus
c) Rhea virus
d) Rabies virus

Influenza is a viral infection that attacks your respiratory system — your nose, throat and lungs. Influenza is commonly called the flu, but it's not the same as stomach "flu" viruses that cause diarrhea and vomiting.

www.medicalcoupons.in



66. Which of the following is cardio protective?

a) HDL
b) LDL
c) VLDL
d) CHYLOMICRON

High-density lipoprotein cholesterol (HDL-C) has been identified as a powerful independent negative predictor of cardiovascular disease.

67. Gluconeogenesis is inhibited by?

a) Glucagon
b) Insulin
c) Cholecystokinin
d) 5 alpha hydroxylase synthase

Insulin is a key hormone that inhibits gluconeogenesis, and insulin resistance is a hallmark of type 2 diabetes.

68. Which structure of Protein is not denatured after heating upto 100 degree?

a) Primary
b) Secondary
c) Tertiary
d) Quaternary

Primary structure, such as the sequence of amino acids held together by covalent peptide bonds, is not disrupted by denaturation.

69. Which of the following is present in skeletal muscle?

www.medicalcoupons.in

a) **GLUT 4**
b) GLUT 5
c) GLUT 7
d) GLUT 2

GLUT4 expression is highest in **adipose tissue and skeletal muscle**, but GLUT4 is also found in other organs such as brain, kidney and intestine

70. Synovial fluid contains–

a) **Hyaluronic acid**
b) Chondroitin sulphate
c) Keratan sulphate
d) Dermatan sulphate

Synovial fluid is produced by the synovium and is composed of **water, inorganic salts and macromolecules, hyaluronic acid, lubricin and aggrecans**, which contribute to the boundary lubrication.

71. Melanin is derived from which amino acid?

a) **Tyrosine**
b) Tryptophan
c) Phenylalanin
d) Alanine

Melanin is a highly irregular heteropolymer consisting of monomeric units derived from the enzymatic oxidation of the **amino acid tyrosine**. The process of melanin formation takes place in specialized acidic organelles (melanosomes) in melanocytes.

72. The bases which are present in human DNA?

a) adenine-guanine-thiamine- uracil
b) adenine-guanine- cytosine-uracil
c) **adenine-guanine-cytosine-thiamine**
d) None of the above

The bases in DNA are **adenine (A), thymine (T), guanine (G) and cytosine (C)**. RNA consists of four nitrogenous bases: **adenine, cytosine, uracil, and guanine**.

73. Keshan's disease is due to deficiency of

www.medicalcoupons.in

a) Selenium
b) Zinc
c) Copper
d) Iron

Keshan disease (KD) is a dilated cardiomyopathy closely related with a diet deficient **in the mineral selenium**. It is named for the northeastern Chinese county Keshan, where the disease prevalence is high because of selenium deficient soil.

74. Type IV complex of ETC is inhibited by;

a) Oligomycin
b) Antimycin
c) Cyanide
d) CO2

Cyanide binds to Fe^{3+} in heme-containing proteins. This inhibits the terminal cytochrome complex IV of the electron transport chain. The blocklock of complex IV by cyanide depletes ATP culminating in cell death.

75. Zinc is cofactor of which enzyme?

a) Carbonic anhydrase
b) Lysyl oxidase
c) Carboxylase
d) Kinase

Carbonic anhydrase is a metalloenzyme that requires **Zinc** as a cofactor and is involved in diverse biological processes including pH regulation, CO2 transfer, ionic exchange, respiration, CO2 photosynthetic fixation, and stomatal closure.

76. Allopurinol inhibits which enzyme?

a) Xanthine oxidase
b) Kinase
c) Lysyl oxidase
d) Carboxylase

Allopurinol was effective in inhibiting **xanthine oxidase activity** in vivo as measured by the dramatic reduction of uric acid production.

77. 1g of protein yields how much energy?

a) 1 Kcal
b) 4 Kcal
c) 8 Kcal
d) 2 Kcal

www.medicalcoupons.in

1 gram of protein (1 **gram times 4 calories** = 4 calories from protein)

78. Ammonia formed in brain, is converted into;

a) Glutamine
b) Glycine
c) Cyteine
d) Urea

Much of the waste nitrogen produced in tissues such as skeletal muscle and brain is packaged as **glutamine**. Glutamine synthase catalyzes the conversion of ammonia and glutamate to glutamine

79. Most sensitive screening test for vitamin A deficiency;

a) Retinol < 10
b) Beta carotene < 10
c) Bitot spot
d) Night blindness

According to the World Health Organization (WHO). **One of the first signs of vitamin A deficiency is night blindness**, Young children and pregnant women are most commonly affected. Vitamin A testing is **used to help diagnose a** vitamin A deficiency in people with signs and symptoms, such as night blindness, or in people who are at risk of vitamin A deficiency.

80. Which enzyme level is tested in thiamine deficiency?

a) PDH
b) Pyruvate kinase
c) Transketolase
d) Kinase

In conjunction with whole blood or erythrocyte **transketolase activity** preloading and postloading, a thiamine loading test is the best indicator of thiamine deficiency. An increase of more than 15% in enzyme activity is a definitive marker of deficiency.

81. Most abundant amino acids found in collagen;

a) Glycine
b) Cysteine
c) Valine
d) Lysine

www.medicalcoupons.in

The triple-helical structure of collagen arises from an unusual abundance of three amino acids: **glycine**, proline, and hydroxyproline. These amino acids make up the characteristic repeating motif Gly-Pro-X, where X can be any amino acid.

82. Which is a specific test for uric acid for gout;

a) Synovial uric acid
b) Plasma uric acid
c) Uric acid in urine
d) Serum uric acid

Joint fluid test (Synovial fluid, also known as joint fluid, is a **thick liquid located between your joints)**. Your doctor may use a needle to draw fluid from your affected joint. Urate crystals may be visible when the fluid is examined under a microscope.

83. HCG is secreted by?

a) Syncytiotrophoblast
b) Yolk sac
c) Cytotrophoblast
d) Placenta

hCG is secreted by **the syncytiotrophoblast** which originates from fused and differentiated cytotrophoblast cells

84. Lady presents with infertility and diagnosed with bilateral cornual block on hysterography. What is the next step?

a) IVF
b) Tuboplasty
c) Laproscopy and hysteroscopy
d) USG

85. Amsel criteria is used in the diagnosis of ?

a) **Bacterial vaginosis**
b) Candidiasis
c) Trichomoniasis
d) Chlymadia infection

The Amsel criteria are in the diagnosis of **bacterial vaginosis**. It is considered particularly helpful during situations in which the diagnostician's microscopic experience, availability of microscope tools, or time may be a limiting factor. There are four parameters used to determine the presence or absence of BV.

www.medicalcoupons.in

86. Testosterone secreted by;

a) Leydig cell
b) Sertoli cell
c) Theca lutein
d) Granulosa cell

The main hormone secreted by **the testes** is testosterone, an androgenic hormone. Testosterone is secreted by cells that lie between the seminiferous tubules, known as the Leydig cells.

87. After a normal delivery in a 27-year-old female, placenta is still attached with uterus. Most common complication which can occur due to forceful traction of cord?

a) Haemorrhage
b) Uterine inversion
c) Uterine rupture
d) Placenta retraction

Controlled cord traction may result in complications such as uterine inversion, particularly if traction is applied before the uterus has contracted sufficiently, and without applying effective counter-pressure to the uterine fundus

88. A patient with 37 weeks of gestation, came to hospital without antenatal check-up and presented with onset of labour. On examination, the mother is Hep B positive, what management should be given to neonate?

a) Hep b vaccine only
b) Hep B vaccine+ IG
c) First IG then HEP B vaccine after 1 month
d) Only IG

Infants born to HBsAg-positive mothers should receive **HBIG (0.5 mL) intramuscularly (IM)** once they are physiologically stable, preferably within 12 hours after birth. HB vaccine, either plasma-derived (10 *gmg per dose) or recombinant (5 *gmg per dose), should be administered IM in three doses of 0.5 mL each.

89. A 32-year lady with uterine fetal death after normal vaginal delivery continues passing of urine from vagina. What is the most probable diagnosis?

a) Bladder rupture
b) Vesicovaginal fistula
c) Urge incontinence
d) Stress incontinence

www.medicalcoupons.in

A vesicovaginal fistula is **an abnormal opening that forms between the bladder and the wall of the vagina**. A fistula can be a complication after vaginal or bladder surgery. The only way to repair this opening is through surgery, which is very often successful.



90. A 16 years old girl with acute bleeding per vaginal was presented to the clinic. What is the immediate management?

a) Uterine ablation
b) Uterine artery block
c) Progesterone
d) Hysteroscopy examination

400mg oral Progesterone SR (compounded) daily, and sometimes higher, has been tried to stop bleeding. Occasionally it works well, and the main side effect is drowsiness. After bleeding is controlled, progesterone is dosed cyclically at normal physiologic doses.

91. A 67 years old female with HTN and diabetes comes with heavy bleeding per vaginal. What is the next management?

a) Uterine ablation
b) Uterine artery block
c) Hysterectomy
d) Laser

Indications of Hysterectomy:

- Fibroids 30%
- Endometriosis
- Uterine prolapse
- Cancer of the uterus, cervix, or ovaries
- Vaginal bleeding, DUB 20%
- Uncontrollable PPH.

www.medicalcoupons.in

92. A lady with amenorrhea and galactorrhea;

a) **Pituitary adenoma**
b) Adrenal hyperplasia
c) 70 alpha hydroxylase deficiency
d) None of above

The most common pathologic cause of galactorrhea is a **pituitary tumor**. Other causes include hypothalamic and pituitary stalk lesions, neurogenic stimulation, thyroid disorders, and chronic renal failure.

93. A 35 years old with 36 week pregnancy comes with history of 5 convulsions at home. Diagnosis made by doctor is eclampsia. Next Management;

a) Only Labetalol
b) Only MgSO4
c) **MgSO4 + Labetalol**
d) Clonidine

labetalol and **hydralazine intravenous infusion** regimens are well tolerated and effective in controlling severe hypertension in pregnant women with severe preeclampsia in combination with magnesium sulfate.

94. A 34 years old lady with 4 children, after 5th normal vaginal delivery, excessive bleeding after placenta removal is due to;

a) Estrogen
b) Progesterone
c) FSH and LH
d) **PPH**

After the placenta is delivered, these contractions help put pressure on the bleeding vessels in the area where the placenta was attached. If the uterus does not contract strongly enough, these blood vessels bleed freely. This is the most common cause of **postpartum hemorrhage**.

95. 32 years old lady with twin pregnancy dichorionic diamniotic, first baby breech presentation and second baby cephalic presentation. What is the management?

a) **C – Section**
b) Assisted breech
c) Instrumental delivery
d) Normal vaginal delivery

www.medicalcoupons.in

Indications of caesarean section in Twins

Obstetrics	Specific for twins
• Placenta previa • Severe preeclampsia • Post C/S • Cord prolapse of the 1st baby • Abnormal uterine contraction • Contracted pelvis	• Both the fetus or even the 1st fetus with non-cephalic presentation. • Conjoined twins • Collision of both heads at the brim preventing engagement of either head. • Monochorionic twins with Twin to twin transfusion syndrome (TTTS)

96. A lady with 36-week pregnancy with previous C section comes with low BP, tachycardia and on USG fluid present in peritoneum. What is diagnosis and next management?

a) **Uterine scar rupture with Laparotomy**
b) Laparoscopy
c) Ectopic pregnancy and abortion
d) None of the above

A complete uterine rupture is **a tear through the thickness of the uterine wall at the site of a prior cesarean incision**. It is a potentially life threatening condition for both the mother and/or the baby and requires immediate surgical intervention. Treatment of uterine rupture is **immediate laparotomy with cesarean delivery** and, if necessary, hysterectomy.

97. 32 years old lady presented with pregnancy, mild bleeding and pain. On examination uterus is tender and fetal heart sound is absent. What is diagnosis?

a) **Abruptio placenta**
b) Fetal distress
c) Ectopic pregnancy
d) Placenta Previa

Placental abruption occurs **when the placenta separates from the inner wall of the uterus before birth**. Placental abruption can deprive the baby of oxygen and nutrients and cause heavy bleeding in the mother

98. A lady with 12-week pregnancy with bleeding and on examination vagina is normal, Internal OS is closed and live birth sign on USG and fundal height is 13 weeks. Diagnosis;

www.medicalcoupons.in

a) **Threatened abortion**
b) Complete abortion
c) Incomplete abortion
d) Inevitable abortion

Threatened Abortion

- It is a clinical entity where the process of abortion has started but has not progressed to a state from which **recovery is impossible.**

→ C/o - **Slight bleeding**
→ O/E - Uterus **- corresponds to gestational age**, C - Internal **os is closed**
→ USG - **live fetus** with subchorionic haemorrhage

99. A newly married couple, the woman is having irregular menstruation. What is the contraceptive of choice?

a) Barrier method
b) OCP
c) Calendar method
d) Progesterone only pills

Oral contraceptive pills -
These are hormonal birth control pills, with some amount of estrogen and progestin to preserve the pregnancy. Low dose oral contraceptive pills are low on hormone contraception. They have less amount of estrogen & progestin as compared to traditional pills. It is advised by doctors to newly married women.

100. The contraceptive which is contraindicated in DVT;

a) OCP
b) Barrier method
c) Billing's method
d) IUD

COCs, combined injectable contraceptives, and combined patch/vaginal ring contraceptives should all be avoided in those with a history of DVT/PE, acute DVT/PE, DVT/PE while receiving established anticoagulant therapy, major surgery with prolonged immobilization, or known thrombophilia

101. Absolute contraindication of IUCD;
a) Acute PID
b) PCOD
c) Breast cancer
d) Anemia

www.medicalcoupons.in

Several absolute contraindications to IUDs are recognized, including: **acute or chronic pelvic inflammatory disease (PID); known or suspected pregnancy**, abnormal uterine bleeding; confirmed or suspected malignancy of the genital tract; and congenital uterine abnormalities or fibroids distorting the cavity in any manner

102. A female came with 36 weeks of gestation, gravid 2 and parity 1, with pain and clinical findings were hematuria, fetal HR was not detected. what is the diagnosis and treatment?

a) Placenta previa and Suction and evacuation
b) Abruption placenta and CS
c) Obstructed labour and CS
d) Abruptio placenta and observation



- ✓ Labour is said to be obstructed when despite good uterine contractions, there is arrest of progress due to mechanical factors causing obstruction to delivery.
- ✓ The uterine contractions are normal (power is normal)
- ✓ There is a **problem in passage or passenger**

Management of obstructed labor-

- **Management of dehydration-** by giving i: v fluids
- **Antibiotics** are given to prevent infection
- **Most important step is to relieve obstruction**

The mode of relieving obstruction is **based on fetal viability**

Viable fetus

- Immediate cesarean

Dead fetus

- Perform a **destructive procedure and relieve the obstruction**
- If the obstetrician **is not well versed with destructive procedures then immediate cesarean**

www.medicalcoupons.in

- Immediate cesarean even in a dead baby because even a **dead baby causing** obstruction can **lead to rupture or severe infections**

103. Angle of cut in episiotomy

a) 45 degrees mediolateral right
b) 30
c) 60
d) 15

In a mediolateral episiotomy, the incision begins in the middle of the vaginal opening and extends down toward the buttocks at a 45-degree angle.

104. Investigation of choice for endometriosis

a) CT
b) MRI
c) USG
d) Laproscopy

Consider **laparoscopy** to diagnose endometriosis in women with suspected endometriosis, even if the ultrasound was normal. ureter, consider a pelvic ultrasound or MRI before an operative laparoscopy. laparoscopic surgery for endometriosis should perform a systematic inspection of the pelvis

105. A multigravida woman in labour room, after delivery and placenta removal, uncontrolled bleeding was seen. What is the most common cause of PPH in this woman?

a) Atonic
b) Traumatic PPH
c) Clotting factor deficiency
d) None of the above

Atonic PPH

- It is the most common cause of primary PPH accounting for 90% of cases.
- The bleeding occurs as the blood vessels are not obliterated by contraction and retraction of uterine muscle fibres

o The causes of atonicity are
- Grand multipara
- Malnutrition/Anemia

www.medicalcoupons.in

- Previous H/o atonic PPH
- Antepartum hemorrhage
- Over distended uterus due to multiple pregnancy, hydramnios and macrosomia
- Uterine malformations or fibroid uterus
- Precipitate labour and prolonged labour
- Mismanaged third stage of labour
- Inadvertent use of oxytocin
- Use of general / epidural anesthesia especially halothane

106. What is the dose of mifepristone in emergency contraception?

a) 600 mg
b) 200 mg
c) 400 mg
d) 100 mg

Mifepristone is a **highly effective and well-tolerated emergency contraceptive** when given in a dose of 600 mg within 72 h of unprotected coitus. We assessed whether the same effectiveness can be achieved with lower doses of mifepristone (50 mg and 10 mg) and a longer postcoital treatment period (120 h)

107. Double decidual sign is seen in;
a) Uterine gestational sac
b) Pseudo gestational sac
c) Threatened Abortion
d) Miscarriage

The double decidual sac sign (DDSS) is a useful feature on **early pregnancy ultrasound** to confirm an early intrauterine pregnancy (IUP) when the yolk sac or embryo is still not visualized.
It consists of the **decidua parietalis (lining the uterine cavity) and decidua capsularis (lining the gestational sac)** and is seen as two concentric rings surrounding an anechoic gestational sac.

108. Most common organism causing pelvic inflammatory diseases

a) Neisseria meningitis
b) GardenellaVaginalis
c) Bacteroids
d) Chalamydia

Most often, PID is caused by infection from two common STIs: **gonorrhea and chlamydia**. The number of women with PID has dropped in recent years. This may be because more women are getting tested regularly for chlamydia and gonorrhea.

www.medicalcoupons.in

109. Late rupture of tubal pregnancy is seen in;

a) Interstitium
b) Isthmus
c) Ampula
d) Fimbriae

An ectopic pregnancy occurs when a fertilized egg grows outside of the uterus. Almost all ectopic pregnancies—more than 90%—occur in a fallopian tube. As the pregnancy grows, **it can cause the tube to burst** (rupture). A rupture can cause major internal bleeding. Very late rupture of tubal pregnancy is seen when the implantation is at Interstitium.

110. A woman post coitus after 96 hours, best contraceptive of choice is

a) OCP
b) IUCD
c) Mifepristone
d) Progesterone only pills

Insertion of a copper IUD should be performed as soon as possible after unprotected or inadequately protected sexual intercourse. It is effective when **placed up to 5 days after sexual** intercourse and, in some studies, was used up to 10 days afterward without failure 18.

111. Delusion is;
a) Disorder of thinking
b) Disorder of memory
c) Disorder of perception
d) Disorder of affect

Delusional disorder, previously called paranoid disorder, is a type of serious mental illness called a **psychotic** disorder. People who have it can't tell what's real from what is imagined. Delusions are the main symptom of delusional disorder. They're unshakable beliefs in something that isn't true or based on reality

112. 25-year-old male had RTA after that he is fearful of driving vehicle and going to that place since 6 weeks and he is having flashbacks of incidents and he is not going for his office also. What is diagnosis?

a) Post-traumatic stress disorder
b) Acute stress
c) Depression
d) Acute panic disorder

Posttraumatic stress disorder (PTSD) is a **psychiatric disorder** that may occur in people who have experienced or witnessed a traumatic event such as a natural

www.medicalcoupons.in

disaster, a serious accident, a terrorist act, war/combat, or rape or who have been threatened with death, sexual violence or serious injury.

113. 20-year-old female scared of elevator since birth. She prefers stairs instead elevator. Diagnosis

a) Agoraphobia
b) Specific phobia
c) Social phobia
d) Acrophobia

Fear of places and situations that might cause panic, helplessness or embarrassment.
Agoraphobia is an anxiety disorder that often develops after one or more panic attacks.
Symptoms include fear and avoidance of places and situations that might cause feelings of panic, entrapment, helplessness or embarrassment.
Treatments include talk therapy and medication.

114. A female was brought to psychiatry after 5 weeks of delivery she appeared to fearfulness irritability and getting suicidal thoughts. diagnosis is

a) Post-partum blues
b) Postpartum depression
c) Postpartum psychosis
d) Schizophrenia

Depression that occurs after childbirth.
Those who develop postpartum depression are at greater risk of developing major depression later on in life.
Symptoms might include insomnia, loss of appetite, intense irritability and difficulty bonding with the baby.
Untreated, the condition may last months or longer. Treatment can include counselling, antidepressants or hormone therapy.

115. A person is unable to find pleasure in doing old activities, which he used to enjoy earlier. Mostly he is suffering from;

a) Alexithymia
b) Depression
c) Anhedonia
d) Derailment

www.medicalcoupons.in

An inability to experience pleasure from activities usually found enjoyable.


COMMON CAUSES
Lack of pleasure can have causes that aren't due to underlying disease. Examples include overwork, recent tragedy, financial problems, bad weather and boring activities.

116. A clinician asked the patient "how is your mood", patient replied "up and down", what could be the diagnosis?

a) Neologism
b) Tangentiality
c) Clanging
d) Derailment

Tangentiality is **the tendency to speak about topics unrelated to** the main topic of discussion. While most people engage in tangentiality from time to time, constant and extreme tangentiality may indicate an underlying mental health condition, particularly schizophrenia

117. Identify the case;



a) Anterior uveitis
b) Glaucoma
c) Optic neuritis
d) Keratoconus

Anterior uveitis is **an inflammation of the middle layer of the eye**. This middle layer includes the iris (colored part of the eye) and adjacent tissue, known as the ciliary body.

118. Keratometer is used to assess;
a) Curvature of cornea
b) Diameter of cornea
c) Thickness of cornea
d) Curvature of lens


www.medicalcoupons.in

A keratometer. This device **measures the curvature of the anterior corneal surface based on the power of a reflecting surface**. It does this by measuring the size of an image reflected from 2 paracentral points and utilizes doubling prisms to stabilize the image enabling more accurate focusing.

119. Internal hordeolum is due to inflammation of-

a) Zeis gland
b) Meibomian glands
c) Mohl gland
d) Lacrimal gland

usually caused by Staphylococcus that infects the eyelash hair follicle. The external hordeolum is caused by a blockage of the sebaceous (Zeis) glands or sweat (Moll) glands. The blockage occurs at the lash line and presents as a painful red swollen area that develops into a pustule. The internal hordeolum is caused by a blockage of the Meibomian glands, and the pustule forms on the inner surface of the eyelids. Hordeola may present on both the upper and the lower eyelids.



120. Identify the clinical condition;



a) Herpes simplex keratitis
b) Bacterial keratitis
c) Mycolic keratitis
d) Vernal keratoconjuctivitis

HSV (Herpes Simplex Virus) keratitis is **an infection of the cornea**—the clear dome that covers the colored part of the eye—that is caused by HSV. The infection usually heals without damaging the eye, but more severe infections can lead to scarring of the cornea or blindness. The hallmark of HSV keratitis is the **presence of multiple small branching epithelial dendrites on the surface of the cornea**, although often times it first presents as a coarse, punctuate epithelial keratitis, which may be


www.medicalcoupons.in

mistaken for a viral keratitis

121. Fleischers ring is seen in;

a) Keratoconus
b) Keratomalacia
c) Keratoglobus
d) Trachoma

Fleischer rings are **indicative of keratoconus**, a degenerative corneal condition that causes the cornea to thin and change to a conic shape.



122. Phlyctenular conjunctivitis seen due to

a) Post fungal infection
b) Allergic
c) Post protozoal
d) Post tuberculoid

Phlyctenular keratoconjunctivitis is an **eye disorder** that involves an immune reaction(allergy) of the cornea (the clear layer in front of the iris and pupil) and conjunctiva (the membrane that lines the eyelids and covers the white of the eye) to bacteria. Small, yellow-gray, raised bumps form on the eye. Phlyctenular keratoconjunctivitis is not an infection. It is an **immune reaction to bacteria** that are on or around the eye. Some of the bacteria that trigger this disorder are staphylococci, and rarely, tuberculosis, and Chlamydia. This disorder is more common among children.

123. Chronic granulomatous inflammation in upper lid (painless swelling);

a) Internal Hordeolum
b) External hordeolum
c) Chalazion
d) Trachoma

A chalazion is a **small, slow-growing lump or cyst that develops within the eyelid**. They are not usually painful and rarely last longer than a few weeks. A chalazion can develop when a meibomian gland at the edge of an eyelid becomes blocked or inflamed. These glands produce oil that lubricates the surface of the eye.

www.medicalcoupons.in

124. In myopia image is formed at;

a) Anterior to retina
b) Posterior to retina
c) On retina
d) None

In a myopic eye, the image of a distant object is formed **in front of the retina** and not at the retina itself. This defect may arise due to (i) excessive curvature of the eye lens, or (ii) (ii) elongation of the eyeball. This defect can be corrected by using a concave lens of suitable power.

125. 20-year-old male complaints of repeatedly changing in glasses, may result into;

a) Keratoconus
b) Pathological myopia
c) Glaucoma
d) Cataract

We must keep in mind that keratoconus is a progressive condition. Unfortunately, an eye with keratoconus is a changing eye and there is no accurate way to predict how fast it will change and how severe it will progress. It is our hope that eventually the condition will slow down and will not worsen beyond a certain point- but do we take our chances as to when we hope that might happen. Even when the vision is best corrected with glasses or contact lenses, it is important to remember that keratoconus can still progress and worsen. It is not uncommon that a prescription that worked wonderfully on a year, will suddenly be not as sharp the next year. Regularly scheduled eye examinations are vital to ensuring clear and comfortable vision.

126. Find out the clinical condition as shown in photo;



a) Keratoconus
b) Limbus
c) Keratomalacia
d) None

www.medicalcoupons.in

A condition in which the clear tissue on the front of the eye (cornea) bulges outward. With keratoconus, the clear, dome-shaped tissue that covers the eye (cornea) thins and bulges outward into a cone shape. Its cause is unknown.
Symptoms first appear during puberty or the late teens and include blurred vision and sensitivity to light and glare.
Vision can be corrected with glasses or contact lenses early on. Advanced cases may require a cornea transplant.

127. Visual loss due to cerebral degeneration is related to which artery?

a) Posterior cerebral artery
b) Middle cerebral artery
c) Internal carotid artery
d) Inferior cerebral artery

Radiographic angiography of the posterior circulation. Bilateral vertebral arteries fuse to form the **basilar artery**. The ultimate branches of the basilar artery form the posterior cerebral arteries (PCAs, arrows), which supply the visual cortex. Infarctions in the PCA territories can cause cortical vision loss.

128. Appropriate treatment for the mild congenital ptosis;

a) LPS Resection
b) Antibiotics and hot compression
c) Tarsal fracture
d) Wedge resection of conjunctiva

Resection of LPS is **the most physiological approach to the treatment of ptosis**. About 70% to 80% ptosis repairs are done by this technique. This is the operation of choice in all cases with fair to good (more than 4 mm.) levator function.

129. Vision 2020, which disease has added recently for India?


a) Cataract
b) Childhood blindness
c) Glaucoma
d) Refractive errors and low vision

VISION2020 aims to eliminate avoidable blindness in the world by 2020 and targets the world's leading causes of avoidable visual impairment: **cataract, trachoma, onchocerciasis**, childhood blindness (including vitamin A deficiency), and refractive error and low vision.

www.medicalcoupons.in

Glaucoma is now regarded as a sort of "missing link" and VISION 2020 India successfully advocated for the inclusion of glaucoma in the next phase by the Indian government.

130. Diabetic patient sudden loss of vision, 6/60 vision, fundus picture is given here. What is the diagnosis?



a) CRVO
b) CRAO
c) Haemorrhage
d) Hypertensive retinopathy

Central retinal vein occlusion (CRVO) is a common retinal vascular disorder. Clinically, CRVO presents with variable visual loss; the fundus may show retinal hemorrhages, dilated tortuous retinal veins, cotton-wool spots, macular edema, and optic disc edema

131. Advantage of contact lens over normal glasses;
a) Prismatic effect will less
b) Prevention from UV rays
c) Decrease inflammation
d) Decrease infection

Contact lenses **move with your eye**, allow a natural field of view, have no frames to obstruct your vision and greatly reduce distortions. Unlike glasses, they do not fog up or get splattered by mud or rain. Contact lenses are excellent for sports and other physical activities. since the contact lens are extremely close to the eye. they reduce the prismatic effect which is normal while wearing spectacles due to the distance between the glasses and the eye

132. Which of the following drug causes postural hypotension commonly?
a) ACE inhibitor
b) Beta blocker
c) Alpha blocker
d) Angiotensin receptor blockers

Alpha blockers are a **type of blood pressure medication**. They lower blood pressure by preventing a hormone called norepinephrine from tightening the muscles

in the walls of smaller arteries and veins. As a result, the blood vessels remain open and relaxed. This improves blood flow and lowers blood pressure. When you start taking an alpha blocker, you might develop **low blood pressure and dizziness**, which can make you faint when you rise from a sitting or lying position.

133. Mechanism of action of allopurinol is

a) Xanthine oxidase inhibition
b) Recombinant uricase
c) Increase uric acid excretion
d) Decrease chemotaxis

Mechanism of Action
Allopurinol and oxypurinol both **inhibit xanthine oxidase**, an enzyme in the purine catabolism pathway that converts hypoxanthine to xanthine to uric acid.

134. Which of the following drugs follow zero order kinetics?
a) Phenytoin
b) Aspirin
c) Morphine
d) Propranolol

Zero order kinetics is **a way of describing how the body uses and breaks down some medicines**. While the rate at which the body eliminates most drugs is proportional to the concentration administered, known as first order kinetics, drugs that work by zero order kinetics work at a predictable, constant rate

List of drugs following zero-order kinetics

- Phenytoin, Phenylbutazone.
- Warfarin.
- Heparin.
- Ethanol.
- Aspirin & other salicylates.
- Theophylline, Tolbutamide.
- Salicylates.

135. Which of the following is an oral direct thrombin inhibitor?
a) Dabigatran
b) Rivaroxaban
c) Warfarin
d) Lepirudin

Dabigatran is a potent, synthetic, reversible, non-peptide thrombin inhibitor. The

www.medicalcoupons.in

inhibition of thrombin results in decreased formation of fibrin and **reduces thrombin-stimulated platelet aggregation** and thus prevents the formation of thrombi.

136. Therapeutic drug monitoring is done for;

a) Phenytoin
b) Heparin
c) Aspirin
d) Metformin

Therapeutic drug monitoring of phenytoin is necessary to ensure **therapeutic and nontoxic levels**. Hypoalbuminemia, renal failure, and interactions with other highly protein-bound drugs (e.g., valproic acid) alter protein binding of phenytoin.

137. A lady has taken medication for ameioliasis infection. She drank alcohol in a party. She has nausea vomiting and dizziness. Which anti amoebic drug could have led to interaction with alcohol to produce these symptoms?

a) Metronidazole
b) Nitazoxanamide
c) Paromomycin
d) Diloxanide

Metronidazole (Flagyl) is a common and effective treatment for bacterial vaginosis. A known adverse effect of metronidazole (Flagyl) is a drug-drug reaction that occurs when metronidazole and alcohol are taken together. Symptoms of this disulfiram-like reaction include **severe nausea and vomiting**.

138. Most potent opioid is;

a. Fentanyl
b. Morphine
c. Pentazocine
d. Pethidine

Fentanyl is a powerful synthetic opioid analgesic that is similar to morphine but is **50 to 100 times more potent**. In its prescription form it is prescribed for pain, but fentanyl is also made illegally.

139. Most cardiotoxic anti-cancer drug among the following;

a) Anthracyclines
b) Cyclophosphamide
c) Lmatinib
d) Tamoxifen

Anthracyclines, such as doxorubicin and idarubicin, remain an important class of

www.medicalcoupons.in

chemotherapeutic agents. Unfortunately, their efficacy in treating cancer is limited by a cumulative dose-dependent cardiotoxicity, which can cause irreversible heart failure.
This progressive cardiotoxicity usually occurs **after the completion of treatment with anthracyclines**, and may become apparent within one year of the completion of treatment (early onset chronic cardiotoxicity) or many years after chemotherapy has been completed (late onset chronic cardiotoxicity).

140. Which of the following is a carbonic anhydrase inhibitor?

a) **Acetazolamide**
b) Hydrochlorthiazide
c) Furosemide
d) Mannitol

Carbonic anhydrase inhibitors are a **medication used in the management and treatment of glaucoma, idiopathic intracranial hypertension**, altitude sickness, congestive heart failure, and epilepsy, among other diseases. Carbonic anhydrase inhibitors are considered part of the diuretic class of medications.
Acetazolamide, dichlorphenamide, and methazolamide are carbonic anhydrase inhibitors.

141. Which of the following is the chemoprophylaxis of choice in a person who is on a journey to endemic malarial region?

a) **Chloroquine**
b) Quinine
c) Lumefantrine
d) Pyrimethamine

Malaria chemoprophylaxis concerns prescribing healthy individuals medication for an infection they have an unknown chance of getting. Sensible use of malaria chemoprophylaxis is a balance between the risk of infection and death, and the risk of side effects.
Chloroquine phosphate is **used to prevent and treat malaria**. It is also used to treat amebiasis. Chloroquine phosphate is in a class of drugs called antimalarials and amebicides. It works by killing the organisms that cause malaria and amebiasis.

142. All of the following drugs are used for prophylaxis of migraine except;

a) Propranolol
b) Amitriptyline
c) **Flutamide**
d) Flunarizine

www.medicalcoupons.in

Drugs of first choice
Propranolol, metoprolol, bisoprolol
Flunarizine*
Topiramate
Valproic acid
Drugs of second choice
Amitriptyline
Methysergide*
Pizotifen*
Low-dose aspirin
Magnesium
Lisinopril
Candesartan
Alternative therapies (drugs of third choice)
Acupuncture
Lamotrigine (only with severe aura symptoms)
Gabapentin
Zonisamide
Venlafaxine
Riboflavin
Feverfew

143. Antidote for opioid poisoning;

a) Naloxone
b) Pethidine
c) Flumazenil
d) Physostigmine

Opioid use can lead to death due to the effects of opioids on the part of the brain which regulates breathing. An opioid overdose can be identified by a combination of three signs and symptoms.

- pinpoint pupils;
- unconsciousness; and
- difficulties with breathing.

Death following opioid overdose is preventable if the person receives basic life support and the timely administration of the drug naloxone. Naloxone is an antidote to opioids that will reverse the effects of an opioid overdose if administered in time. Naloxone has virtually no effect in people who have not taken opioids.

144. Monoclonal antibody used in cancer is

a) Rituximab
b) Cisplatin
c) 5-fluorouracil
d) Methotrexate

Rituximab is a chimeric monoclonal antibody targeted against CD20 which is a surface antigen present on B cells. Therefore, it acts by **depleting normal as well as pathogenic B cells** while sparing plasma cells and hematopoietic stem cells as they do not express the CD20 surface antigen.

www.medicalcoupons.in

145. A patient after valve replacement will require follow up treatment with

a) Warfarin
b) ACE inhibitors
c) Beta blockers
d) Thiazides

To help prevent thromboembolism, people who receive mechanical heart valves must take **anticoagulant medications (anticlotting drugs)** for the rest of their lives. All patients with mechanical prosthetic valves **require lifelong VKA (eg, warfarin)** anticoagulation. Additional indications for anticoagulation, such as atrial fibrillation (AF), may require a higher goal INR than for the mechanical valve alone

146. All of the following can result in gynecomastia except;

a) Liver failure
b) Spironolactone
c) Digoxin
d) Aromatase inhibitors

Gynecomastia can be **caused by an imbalance between the sex hormones testosterone and oestrogen**. Oestrogen causes breast tissue to grow. While all men produce some oestrogen, they usually have much higher levels of testosterone, which stops the oestrogen from causing breast tissue to grow.
Aromatase inhibitors work by **blocking the enzyme aromatase**, which turns the hormone androgen into small amounts of estrogen in the body. This means that less estrogen is available to stimulate the growth of hormone-receptor-positive breast cancer cells.

147. Hydrochlorothiazide works by inhibiting

a) Na+ Cl pump in late DCT
b) Na+ Cl pump in early DCT
c) Na+ K+ 2CT pump in descending limb of loop of henle
d) Na+K+2CT pump in ascending limb of loop of henle

Hydrochlorothiazide acts on the distal convoluted tubules and inhibits **the sodium chloride co-transporter system**. This action leads to a diuretic action that lowers blood pressure, but there is also a potassium loss in the urine.

148. The drug of choice for hyperthyroidism in third trimester of pregnancy is;

a) Carbimazole
b) Prophylthyouracil
c) Sodium iodide
d) Radioactive iodine

PTU(propylthiouracil) **Propylthiouracil** is the drug of choice for hyperthyroidism in

www.medicalcoupons.in

first trimester of pregnancy. It has a side effect of hepatotoxicity and hence due to this reason DOC for hyperthyroidism in second and third trimester is Carbimazole

149. Mechanism of action of exenatide in diabetes mellitus is

a) It is analogue of GLP released from gut and increase glucose dependant insulin secretion
b) It is DPP-4 inhibitor and result in decreased breakdown of GLP
c) It inhibits SGLT-2 and cause glucosuria
d) It is amylin analogue and decrease glucagon

Exenatide is a GLP-1 receptor agonist released from the gut and **acts to increase glucose-dependent insulin secretion from pancreatic beta cells**, suppress glucagon secretion, delay gastric emptying, and reduce food intake. The binding of the drug to pancreatic GLP-1 receptors mediates these actions

150. Which anti-asthma drug is avoided with erythromycin?

a) Salbutamol
b) Ipratropium
c) Theophylline
d) Terbutaline

Administration of **allopurinol**, oral contraceptives, cimetidine, erythromycin, quinolone antibiotics, fluvoxamine, zileuton, and zafirlukast increase theophylline levels.
Using erythromycin together with theophylline **increase the effects of theophylline**. High serum theophylline levels **increase the risk of toxicity**. Toxic effects include nausea and vomiting, headaches, gastric discomfort, diuresis, insomnia, cardiac arrhythmias, behavioral disturbances, and epileptic seizures

151. McNaughton's rule;

a) IPC 82
b) IPC 84
c) IPC 87
d) IPC 85

Section 84 IPC embodies McNaughton rules as follows: "**Nothing is an offence which is done by a** person who, at the time of doing it, by reason of unsoundness of mind, is incapable of knowing the nature of the act or that he is doing what is either wrong or contrary to the law.

152. A farmer with pinpoint pupils, increased secretions and urination. What is the most likely diagnosis?

a) Organophosphate poisoning
b) Opioid poisoning
c) Alcohol poisoning

www.medicalcoupons.in

d) Atropine poisoning

Organophosphate poisoning is poisoning due to organophosphates (OP's).
Organophosphates are used as insecticides, medications, and nerve agents.
Symptoms include **increased saliva and tear production**, diarrhea, vomiting, small pupils, sweating, muscle tremors, and confusion.
The mainstays of medical therapy in organophosphate (OP) poisoning include **atropine, pralidoxime (2-PAM), and benzodiazepines (eg, diazepam)**.
Initial management must focus on adequate use of atropine.

153. Posthumous birth;
a) Child born after death of father
b) After death of mother during delivery
c) After death of both parents
d) None

A posthumous birth is the birth of a child after the death of a biological parent. A person born in these circumstances is called a posthumous child or a posthumously born person

154. Poisoning caused by bullet retained in body is;
a) Lead
b) Phosphorus
c) Nitro cellulose
d) Iron

Lead poisoning is an unusual complication of gunshot wounds that occurs when retained **lead bullet** fragments are in contact with body fluids capable of solubilizing lead. The epidemic of violence by gunfire may result in increasing numbers of lead poisoning cases from this exposure.

155. After injury hair bulbs are seen crushed under microscopy. Probable type of wound;
a) Abrasion
b) Laceration
c) Stab injury
d) None

Lacerations
- These wounds are commonly known as 'gashes, tears or cuts' of the skin
- **It is an open wound.**
- Breach in the epidermis, dermis or underlying tissue as a result of blunt force

Features
- **Tissues not cleanly divided and** is **bridged across base of the wound.**

- Foreign material may be found as well
- **Hair and hair bulbs are crushed**
- **Margins are irregular, ragged and uneven and their ends are pointed or blunt**
- **Lacerations produced without excessive skin crushing may have relatively regular sharp margins.**
- Bruising is seen either in the skin or subcutaneous tissue around the wound
- Hair bulbs are crushed
- Hair and epidermal tags may be driven deeply into the wound
- Less haemorrhage
- Presence of foreign matter in the wound
- Shape and size of the wound does not correspond to the weapon used

156. A patient was found to have pink coloured skin and mucosa, bitter almond smell from breath and frothy discharge. Cause of death in this patient;

a) Cyanide poisoning
b) CO poisoning
c) H2S poisoning
d) Dhatura poisoning

Cyanide can be a colorless gas, such as hydrogen cyanide (HCN) or cyanogen chloride (CNCl), or a crystal form such as sodium cyanide (NaCN) or potassium cyanide (KCN). Cyanide sometimes is described as having a “bitter almond” smell **Cyanide prevents the cells of the body from using oxygen**. When this happens, the cells die. Cyanide is more harmful to the heart and brain than to other organs because the heart and brain use a lot of oxygen.

157. Entry wound kind of depression will be seen in

a) Inversion
b) Eversion
c) Pond’s fracture
d) Depression

Entrance wounds: point where the bullet enters the biological target

- General features:
  - Abrasion ring: a reddish brown abrasion on the skin surrounding the gunshot wound of entrance
  - Greasy rim or bullet wipe is a dirt collar due to the lubricant carried on the bullet
  - Inverted edges: due to the bullet movement inside the body
  - Smaller than bullet caliber (usually): due to the elastic recoil of the affected tissues

www.medicalcoupons.in

158. A lady comes to OPD after fall from scooty. Her vitals are stable. She is having continuous, clear watery discharge from **nose after 2 days. These are most likely the feature of;**

a) CSF rhinorrhoea
b) Acute respiratory infection
c) Middle cranial fossa fracture
d) Rhinitis

Cerebrospinal fluid (CSF) rhinorrhea is **a condition where the fluid that surrounds the brain leaks into the nose and sinuses**. Head trauma, surgery, or even birth defects can make a hole in the membranes that hold this fluid. It then leaks into your nose or ear, causing a watery, runny nose

159. Office headache is due to inflammation of which sinus?

a) Frontal
b) Maxillary
c) Ethmoidal
d) Sphenoid

Frontal sinusitis is inflammation or infection of the sinuses located just behind the eyes and in the forehead. The sinuses are a system of connected hollow cavities in the face that contain air and a thin layer of mucus. All sinuses produce mucus that moisturizes the airways and drains into the nasal passages.
If the frontal sinuses are inflamed or infected, they cannot drain mucus efficiently, and this can make breathing difficult. It can also lead to a feeling of increased pressure around the eyes and forehead.
Since most cases of acute frontal sinusitis are caused by a viral infection, treatment would be to recommend **taking a nasal spray or decongestant** to decrease inflammation, assist with mucus drainage, and relieve pressure in the frontal sinuses

160. Adam's apple in Male is due to;

a) Thyroid cartilage
b) Cricoid cartilage
c) Hyoid cartilage
d) Epiglottis cartilage

The Adam's apple is made up of **the thyroid cartilage**. Cartilage is the same tissue that makes up your nose, ears, and windpipe (i.e., trachea). The voice box and windpipe have several kinds of cartilage, which work together to make sure your airways stay clear and you're able to speak.

161. Most prominent and larger air cell of ethmoidal sinus?

a) Agger nasi cell
b) Haller cell
c) Onodi cell

www.medicalcoupons.in

d) **Bulla ethmoidalis**

Bulla ethmoidalis

> **This is the largest and most prominent anterior ethmoid air cell. It is related laterally to the lamina papyracea. It may fuse with the skull base superiorly and with the lamella basalis posteriorly**

162. Most common presentation of nasopharyngeal carcinoma;

a) **Neck lymph node**
b) Blockage of nose
c) Loss of smell
d) Blood tinged discharge

The four most common symptoms in order of frequencies were **cervical lymphadenopathy, epistaxis, hearing loss and diplopia**. 80.8% of the patients presented with cervical lymphadenopathy and about 85% of the patients presented in the advanced stages.

Other signs and symptoms may include

- Swelling of the neck.
- Persistent headaches.
- Nasal congestion (a blocked nose)
- Facial pain.
- Nosebleeds.
- Changes in hearing.
- Ringing in the ears.
- Many people have no symptoms

163. EBV is associated with;

a) **Nasopharyngeal carcinoma**
b) Esophageal cancer
c) Gastric cancer
d) Renal carcinoma

EBV-associated diseases

- Nasopharyngeal carcinoma (NPC) ...
- EBV-associated gastric cancer...
- Breast cancer. ...
- HL and NHL. ...
- Post-transplant lymphoproliferative carcinoma (PTLD) ...
- Infectious mononucleosis (IM) ...
- Chronic active EBV infection (CAEBV) ...
- EBV-associated hemophagocytic lymphohistiocytosis (EBV-HLH)

www.medicalcoupons.in

164. All of the following is true about Meniere's disease except;

a) Idiopathic
b) Low frequency hear loss
c) Increase in endolymph
d) Destroying the cochlea is mandatory

An inner ear disorder that causes episodes of vertigo (spinning).
Ménière's disease usually starts in one ear, but later may involve both. Smoking, infections or a high-salt diet may worsen the disease.
Symptoms include a spinning sensation (vertigo), hearing loss, ear ringing (tinnitus) and ear pressure. The vertigo may cause severe nausea and imbalance. Hearing loss may become permanent.
Drugs for motion sickness or nausea may help manage symptoms.

165. Investigation used for CSF rhinorrhoea are all except

a) CT cisternogram
b) Skull X-ray
c) beta2 transferrin
d) Nasal endoscopy

Diagnosing a CSF leak includes **an analysis of the nasal fluid for a protein called beta-2 transferrin** which is most only found in cerebrospinal fluid. X-ray,CT and MRI scans may also be requiring to determine the location and severity of the leakage.

166. Treatment of choice in nasopharyngeal carcinoma;

a) Chemotherapy
b) Radiotherapy
c) Chemo given with radiation
d) Surgery

The main treatment for NPC is **radiation therapy**. It is often given in combination with chemotherapy. This approach may be called concomitant chemoradiotherapy. Surgery for NPC is occasionally used, mainly to remove lymph nodes after chemoradiotherapy or to treat NPC that has come back after initial treatment.

167. A 6-year-old boy came to hospital with complaints of sore throat and difficulty in swallowing, his left tonsils were pushed medially. what will be the diagnosis?

a) Peritonsillar abscess
b) Retropharyngeal abscess
c) Ludwig's angina
d) Parapharyngeal abscess

www.medicalcoupons.in


A parapharyngeal abscess is a deep neck space abscess of the parapharyngeal space, which is lateral to the superior pharyngeal constrictor muscle and medial to the masseter muscle. This space is divided by the styloid process into anterior and posterior compartments. A parapharyngeal abscess is a deep neck abscess. Symptoms include **fever, sore throat, odynophagia, and swelling in the neck down to the hyoid bone**. Diagnosis is by CT. Treatment is antibiotics and surgical drainage.

168. A young boy came to OPD with the complaints of difficulty in breathing. On examination, bilateral polyps were found. On aspiration, bleeding was seen. What will be the initial management?

a) I and D
b) Antihistaminic
c) Conservative
d) Septoplasty

Corticosteroids. **Oral and topical nasal steroid administration** is the primary medical therapy for nasal polyposis. Antihistamines, decongestants, and cromolyn sodium provide little benefit. Immunotherapy may be useful to treat allergic rhinitis but, when used alone, does not usually resolve existing polyps.

169. Identify condition in the shown image?




a) Impetigo contagiosa
b) Contact dermatitis
c) Molluscum contagiosum
d) HSV

Nonbullous impetigo, or impetigo contagiosa, is **caused by Staphylococcus aureus or Streptococcus pyogenes**, and is characterized by honey-colored crusts on the face and extremities. Impetigo primarily affects the skin or secondarily infects insect bites, eczema, or herpetic lesions.
A highly contagious skin infection that causes red sores on the face.
Impetigo mainly affects infants and children.
The main symptom is red sores that form around the nose and mouth. The sores rupture, ooze for a few days, then form a yellow-brown crust.
Antibiotics shorten the infection and can help prevent spread to others.

www.medicalcoupons.in

170. Exclamation mark hair pattern is seen in



a) Alopecia areata
b) Androgenic alopecia
c) Anagen effluvium
d) Telogen effluvium

Sudden hair loss that starts with one or more circular bald patches that may overlap.
Alopecia areata occurs when the immune system attacks hair follicles and may be brought on by severe stress.
The main symptom is hair loss.
Treatment may address any underlying conditions and includes topical scalp medication.
The characteristic diagnostic finding of alopecia areata is the exclamation point hair. These can be found in areas of hair loss and are **short broken off hairs** that are narrower closer to the scalp and therefore mimic an exclamation point. In some cases a biopsy is necessary for diagnosis.

171. What is the name of this appearance which is seen in pellagra?



a) Burn
b) Castle neck appearance
c) Shawl sign
d) Gottron's papule

www.medicalcoupons.in

The Casal collar or Casal necklace is a clinical sign in which there is **an erythematous pigmented skin rash in the distribution of a broad collar** (dermatomes C3 and C4). It is seen in patients with pellagra, as a result of vitamin b3 deficiency. The sign is named after Gaspar Casal


172. A lady with 50% burns of dermis & subcutaneous tissue came to emergency department. Burns will be classified as;

a) 1st degree
b) 2nd degree superficial
c) 2nd degree deep
d) 3rd degree burn

Classification	Characteristics
First degree	Partial thickness burns. ✸ Characterized by erythema (localized redness). ✸ Appear sunburn-like. ✸ Are not included when calculating burn size. ✸ Usually heal by themselves.
Second degree	Partial thickness burns. ✸ Part of skin has been damaged or destroyed. ✸ Have blisters containing clear fluid. ✸ Pink underlying tissue. ✸ Often heal by themselves.
Third degree	Full thickness burns. ✸ Full skin has been destroyed. ✸ Deep red tissue underlying blister. ✸ Presence of bloody blister fluid. ✸ Muscle and bone may be destroyed. ✸ Require professional treatment.
Fourth degree	Full thickness burns. ✸ Penetrate deep tissue to fat, muscle, bone. ✸ Require immediate professional treatment.

www.medicalcoupons.in

173. Identify the abnormal condition given below;



a) Testicular torsion
b) Hydrocele
c) Varicocele
d) Testicular atrophy

A twisting of the male organ that makes hormones and sperm (testicle).
When the testicle rotates (testicular torsion), it twists the cord supplying blood to the loose bag of skin (scrotum) beneath the penis. This may occur after vigorous activity, a minor injury to the testicles or sleep.
Sudden, severe pain and swelling in the testicle are symptoms.
Surgery is required. Treated promptly, the testicle can often be saved. A longer wait may affect fertility.

174. 20 year old female diagnosed previously with adenocarcinoma stomach and on examination following is seen;



a) Sister marry joseph nodule
b) Ulcer
c) Infected umbilical hernia
d) Irish node

Sister Mary Joseph nodule or Sister Mary Joseph Sign refers to a palpable nodule bulging into the umbilicus as a result of metastasis of a malignant cancer in the pelvis or abdomen. Gastrointestinal malignancies account for about half of the underlying sources (gastric, colonic, pancreatic cancer), gynecologic (ovarian, uterine cancer), unknown primary tumors and rarely bladder or respiratory malignancies cause umbilical metastasis.

175. Rodent ulcer is
a) Basal cell cancer
b) Squamous cell ca
c) Rhinophyma

d) Adenocarcinoma

Rodent ulcers are **a type of skin cancer**. Also known as basal cell carcinoma, rodent ulcers occur in the lower most layer of your skin. They usually develop in skin areas exposed to the sun. Rodent ulcers are a very slow growing cancer that usually does not spread to other parts of your body.

176. Most common oral cancer;

a) Squamous cell ca
b) Adenocarcinoma
c) Transition cell ca
d) Mucoepidermoid

Squamous cell carcinoma is the most common type of mouth cancer, accounting for 9 out of 10 cases. Squamous cells are found in many areas of the body, including the inside of the mouth and in the skin. Less common types of mouth cancer include: adenocarcinoma, which is cancers that develop inside the salivary glands.

177. Distal ileum was removed in a 20-year-old girl. Which absorption deficient will be seen?

a) Iron
b) Bile salts
c) Folic acid
d) Copper

The main function of the ileum is to **absorb vitamin B_{12}, bile salts**, and whatever products of digestion were not absorbed by the jejunum. ... They absorb fatty acid and glycerol, the products of fat digestion.

178. 62-year-old female has kidney stone and treated with PCNL. After 2 days she again comes to OPD with chills and fever. What is the complication?

a) Bacterial sepsis
b) Acute pancreatitis
c) Splenic injury
d) Ureteric stricture

Bacterial sepsis is a **life-threatening condition** that arises when the body's response to an infection injures its tissues and organs. Sepsis is caused by conditions that threaten the functional integrity of the host such as microbial invasion. **The signs and symptoms of sepsis can include a combination of any of the following:**

- confusion or disorientation,
- shortness of breath,
- high heart rate,
- fever, or shivering, or feeling very cold,

www.medicalcoupons.in

- extreme pain or discomfort, and.
- clammy or sweaty skin.

179. Cellulitis is

a) Nonsuppurative and noninvasive
b) Suppurative and noninvasive
c) Nonsuppurative and invasive
d) Suppurative and invasive

Nonpurulent cellulitis includes **rapidly spreading superficial cellulitis** and erysipelas; typically involves groups A, B, C, and G beta-hemolytic streptococci and, occasionally, methicillin-susceptible Staphylococcus aureus (MSSA); these infections are diagnosed clinically

180. identify the condition as shown in image;



a) Port wine stain
b) Strawberry nevus
c) Erythema multiforme
d) Exanthem subitem

A port-wine stain is **a type of birthmark**. It got its name because it looks like maroon wine was spilled or splashed on the skin. Though they often start out looking pink at birth, port-wine stains tend to become darker (usually reddish-purple or dark red) as kids grow.
Port-wine stains are caused by **an abnormal formation of tiny blood vessels in the skin**. In rare cases, port-wine stains are a sign of Sturge-Weber syndrome or Klippel-Trenaunay-Weber syndrome.

181. 28-year-old alcoholic patient walks to hospital with the complaints of binge vomiting, chest pain, fever and pneumomediastinum. The most probable condition;

www.medicalcoupons.in

a) **Boerrhave syndrome**
b) Tension pneumo thorax
c) PUD perforation
d) Mallory weis tear

Effort rupture of the esophagus, or Boerhaave syndrome, is **a spontaneous perforation of the esophagus that results from a sudden increase in intraesophageal pressure combined with negative** intrathoracic pressure (eg, severe straining or vomiting).
It is induced by **increased esophageal pressure followed by straining**, which can result from retching, vomiting, weightlifting, childbirth, or defecation

182. Abdominal pain, fever and jaundice. This triad is known as;
a) **Charcot's triad**
b) Saint's triad
c) Virchow triad
d) Renault's triad

Charcot's cholangitis triad is **the combination of jaundice; fever, usually with rigors; and right upper quadrant abdominal pain**. It occurs as a result of ascending cholangitis (an infection of the bile duct in the liver).

183. 40 years old patient with femur fracture with pulmonary infiltration and respiratory distress
a) **Fat embolism**
b) Pulmonary embolism
c) Air embolism
d) Obstruction

A disruption to blood supply caused by fat globules in a blood vessel.
A fat embolism occurs when fat globules are released into the bloodstream. It's most commonly associated with a trauma, such as a bone fracture.
Symptoms, if present, typically occur 24 to 72 hours after the trauma. They include shortness of breath, confusion and a rash.

184. Which patient has better prognosis in breast cancer?
a) **Luminal A**
b) Luminal B
c) Patient who has triple test negative
d) Patient who has triple test positive

Luminal A breast cancer is hormone-receptor positive (estrogen-receptor and/or

www.medicalcoupons.in

progesterone-receptor positive), HER2 negative, and has low levels of the protein Ki-67, which helps control how fast cancer cells grow. Luminal A cancers are **low-grade**, tend to grow slowly and have the best prognosis

185. 33-year man presented with fever and pain in upper right hypochondrium after food intake. Investigation of choice,

a) CECT
b) Ultrasound
c) MRI
d) HRCT

Ultrasonography is the most common test used in the emergency department for the diagnosis of biliary colic and acute cholecystitis. This imaging modality may be diagnostic for biliary disease, help exclude biliary disease, or may reveal alternative causes of the patient's symptom

186. Long term diabetic patient with blisters walked barefoot few miles on hot sand. He presented with this clinical condition. What is the most probable diagnosis?



a) Diabetic foot
b) Burn
c) Necrotizing Fasciitis
d) Elephantiasis

A serious bacterial infection that destroys tissue under the skin.
Flesh-eating disease occurs when bacteria enter the body through a break in the skin. People with a weakened immune system can be at greater risk of developing this condition.
The condition spreads quickly. Symptoms include blisters, fever, fatigue and pain worse than a person would expect based on the wound's appearance.
Treatment involves immediate delivery of IV antibiotics. Surgical removal of dead or infected tissue from the wound is often required.

187. A man was presented to emergency department with head injury after an accident with vehicle. Investigation of choice should be;

www.medicalcoupons.in

a) CECT
b) MRI
c) NCCT
d) MRI

Noncontrast head CT is **the most useful study for the evaluation of patients after trauma, focal neurologic symptoms, change in mental status, and severe headaches** to rule out life-threatening pathology. Clinicians in all fields must be able to readily identify a number of different abnormalities.

188. ALVARADO score 2 defines:
a) Temperature
b) Leukocytosis
c) Tenderness in left iliac fossa
d) Migratory pain

Alvarado score	
Feature	**Score**
Migration of pain	1
Anorexia	1
Nausea	1
Tenderness in right lower quadrant	2
Rebound pain	1
Elevated temperature	1
Leucocytosis	2
Shift of white blood cell count to the left	1
Total	**10**

189. A patient of Varicose veins came to hospital; intern was on duty. Which test he shall perform to rule out the DVT?
a) Brodie Trendelenberg test
b) Perthe's test
c) Thomas test
d) Ober test

The **Brodie–Trendelenburg percussion test** is a medical test to determine valvular incompetence in superficial veins. A finger is placed over the lower (distal) part of the vein being examined. The upper (proximal) part of the vein is then tapped (percussed). If the impulse is felt by the finger placed at the lower end, it indicates incompetence of valves in that vein.

www.medicalcoupons.in

190. Early investigation of choice for genital TB

a) Early Morning Urine Sample
b) Urine sample with giemsa staining
c) Endometrial ablation
d) Incision and curettage

Early morning urine (EMU) is accepted as a **useful method for improving the concentrations** of a number of antigens and analytes found in human urine. The strategy has been shown to improve the yield of urine culture for TB diagnosis

Urine is easily obtainable and can be tested using either a **lateral flow assay (LFA)** for TB-lipoarabinomannan (TB-LAM) or the Xpert MTB/RIF assay (Xpert; Cepheid, Sunnyvale, CA). Tuberculosis-LAM is emerging as the key urine-based TB diagnostic

191. True about crohn's disease except

a) Recurrence is more common
b) Rectum is involved
c) Fissures are formed
d) Transmural

A chronic inflammatory bowel disease that affects the lining of the digestive tract. Crohn's disease can sometimes cause life-threatening complications.
Crohn's disease can cause abdominal pain, diarrhoea, weight loss, anaemia and fatigue. Some people may be symptom-free most of their lives, while others can have severe chronic symptoms that never go away.
Crohn's disease cannot be cured. Medications such as steroids and immunosuppressants are used to slow the progression of disease. If these aren't effective, a patient may require surgery. Additionally, patients with Crohn's disease may need to receive regular screening for colorectal cancer due to increased risk.

192. Patient underwent gastrectomy, after eating within 20 minutes sweating, diarrhoea, what could be the cause?

a) Hyperglycemia
b) Early dumping syndrome
c) Late dumping syndrome
d) Hypoglycemia


Early dumping syndrome happens when **the sudden influx of food into your intestine causes a lot of fluid to move from your bloodstream into your intestine** as well. This extra fluid causes diarrhea and bloating. Your intestines also release substances that speed your heart rate and lower your blood pressure.

193. In an accident case, after the arrival of medical team, all should be done in early management except;

www.medicalcoupons.in

a) Stabilization of cervical vertebrae
b) Check BP and Pulse
c) Check Respiration
d) Glasgow coma scale


The **Glasgow Coma Scale** (GCS) is used to objectively describe the extent of impaired consciousness in all types of acute medical and trauma patients. The scale assesses patients according to three aspects of responsiveness: eye-opening, motor, and verbal responses.



194. Patients are categorized on the basis of chances of survival in disaster management is;

a) Triage
b) Mitigation
c) Tagging
d) Surge capacity

www.medicalcoupons.in



195. Risk factors of oesophageal carcinoma are all except

a) GERD
b) Betel chewing
c) Oxalate food
d) Spicy food

Multiple risk factors are associated with the development of esophageal adenocarcinoma. These include **Barrett's esophagus**, acid peptic disorders, motor disorders of the esophagus, other malignancies, medications, environmental exposures, diet, and nutrition.

196. Left kidney transplantation will be placed in which place?

a) Iliac fossa
b) kidney fossa
c) Retroperitoneal space
d) Left lower quadrant

Traditionally **the right iliac fossa is the standard fossa** for a kidney transplantation procedure and the **left iliac fossa is the preferred site for simultaneous kidney-pancreas transplantation**.

197. For severe malnutrition MAC will be less than-

a) 14.5
b) 13.5
c) 12.5
d) 11

Mid-arm circumference of < 11cm indicates severe malnutrition in infants from 1-6 months of age. Physiopathological principle of arm circumference measurement in children aged 1-5 years and the relationship with severity of malnutrition

www.medicalcoupons.in

198. 3 months old baby with fever and respiratory rate 60, irritable, and well feed, no stridor, no chest indrawing, no convulsions. Diagnosis-

a) Pneumonia
b) Severe pneumonia
c) Very sever pneumonia
d) No Pneumonia

SIGNS	CLASSIFY AS	IDENTIFY TREATMENT (Urgent pre-referral treatments are in bold print.)
• Any general danger sign or • Chest indrawing or • Stridor in calm child.	SEVERE PNEUMONIA OR VERY SEVERE DISEASE	➤ Give first dose of an appropriate antibiotic. ➤ Refer URGENTLY to hospital.
• Fast breathing	PNEUMONIA	➤ Give an appropriate oral antibiotic for 5 days. ➤ Soothe the throat and relieve the cough with a safe remedy. ➤ Advise mother when to return immediately. ➤ Follow-up in 2 days.
No signs of pneumonia or very severe disease.	NO PNEUMONIA: COUCH OR COLD	➤ If coughing more than 30 days, refer for assessment. ➤ Soothe the throat and relieve the cough with a safe remedy. ➤ Advise mother when to return immediately. ➤ Follow-up in 5 days if not improving.


199. Number of primary teeth;

a) 20
b) 24
c) 16
d) 32

At birth people usually have **20 baby** (primary) teeth, which start to come in (erupt) at about 6 months of age. They fall out (shed) at various times throughout childhood. By age 21, all 32 of the permanent teeth have usually erupted.

200. All of the following are types of Primary headache except

a) Migraine
b) Cluster
c) Tension
d) Temporal arteritis



www.medicalcoupons.in

201. Which is associated with alcohol related cardiomyopathy

a) Dilated
b) Restrictive
c) Hypertrophic
d) None

Alcoholic cardiomyopathy is a type of dilated cardiomyopathy typically found in people with alcohol use disorder. Due to the direct toxic effects of alcohol on heart muscle, the heart is unable to pump blood efficiently, leading to heart failure.

202. 40-year-old male presents with fever and abdominal pain and diagnosed with HIV and TB. How will you give treatment?

a) ATT and AIDS treatment simultaneously.
b) First ATT and then ART
c) ATT only
d) First ART and then ATT

WHO guidelines recommend starting ART **between 2 and 8 weeks** of TB treatment in patients with CD4 < 200 cells/µL and during continuation phase in those with CD4 counts 200-350 cells/µL . In patients with very low CD4 counts there is greater urgency to start ART.

203. Patient with abdominal pain, IgA deposition, haematuria and purpura over leg as given in image. Diagnosis;



a) IgA nephropathy
b) HSP
c) Buerger's disease
d) PSGN

A disorder causing inflammation and bleeding in the small blood vessels. Henoch-Schönlein purpura affects the small blood vessels of the skin, joints, intestines and kidneys. It's most common before the age of seven but can affect anyone.
Symptoms include reddish-purple spots on the lower extremities, swollen and sore joints, abdominal pain or bloody urine.
The condition is often treated with steroids.

www.medicalcoupons.in

204. Patient presents with diarrhoea, abdominal pain and finding on upper GI endoscopy is shown in the photo. What is the pathology?



a) Antigliadin antibody
b) Antimitochondrial antibody
c) Anticentromere antibody
d) Antitopoisomerase antibody

A gliadin antibodies test is used to help **doctors diagnose celiac disease**. Celiac disease is an autoimmune disorder in which the body's immune system mistakenly thinks that gluten — a protein in wheat, barley, rye, and oats

205. Almost odourless Stool seen in-
a) Upper GI bleeding
b) Bacillary Dysentery
c) Ameobiasis
d) Cholestatis stool

Character	Amoebic dysentery	Bacillary dysentery
Macroscopic		
Number	6-8 motions a day	Over 10 motions a day
Amount (Volume)	Relatively copious	Small amount
Appearance and Amount	Blood mucus, semi formed	Blood mucus, mainly water
Odour	Offensive (fishy odour)	Odourless
Colour	Dark red (altered blood)	Bright red (fresh blood)
Reaction	Acidic	Alkaline
Consistency	Not adherent to the container	Adherent to the container

206. Hyperkalemia means more than
a) 4.5 meq/l
b) 5.5 meq/l
c) 7.5 meq/l
d) 10.5 meq/l

www.medicalcoupons.in

A typical potassium level for an adult falls between 3.5 and 5.0 millimoles per liter (mmol/L). Hyperkalemia occurs when levels go **above 5.5 mmol/L**. A reading above 6.5 mmol/L can cause heart problems that require immediate medical attention

207. Female presents with galactorrhea and amenorrhea and diagnosed with Pituitary adenoma and now complains of loss of vision which will be-

a) Bitemporal hemianopia
b) Binasal hemianopia
c) Homonymous hemianopia
d) Superior temporal quadrantopia

Characteristically lesions at the level of the optic chiasm produce a bitemporal hemianopia. Pituitary adenomas, which grow upward from the pituitary stalk, compress the chiasm from below, which preferentially involves the inferior, nasal, and macular nerve fibers.

208. Women presented with on and off headache. On lateral x-ray suprasellar calcifications are seen and she has heteronymous hemianopia. Diagnosis;

a) Craniopharyngioma
b) Pituitary macroadenoma
c) Astrocytoma
d) None

Craniopharyngiomas are **benign tumors that grow near the pituitary gland**. They can develop as solid tumors or cysts (hollow sacs filled with fluid). Approximately 10 percent to 15 percent of pituitary tumors are craniopharyngiomas.
Craniopharyngioma is **a rare type of noncancerous (benign) brain tumor**. Craniopharyngioma begins near the brain's pituitary gland, which secretes hormones that control many body functions. As a craniopharyngioma slowly grows, it can affect the function of the pituitary gland and other nearby structures in the brain

209. Which is not a contraindication for OPV?

a) Leukaemia
b) Diarrhoea
c) Malignancy
d) Immunocompromised

False Contraindications

- diarrhea.
- minor upper respiratory illnesses with or without fever.
- mild to moderate local reactions to a previous dose of vaccine.
- current antimicrobial therapy.
- the convalescent phase of an acute illness.

www.medicalcoupons.in

210. Unconscious patient, most common cause of respiratory obstruction;

a) Tongue fall
b) Head injury
c) Respiratory center block
d) Hyoid fracture

The tongue is the most common cause of upper airway obstruction, a situation seen most often in patients who are comatose or who have suffered cardiopulmonary arrest. Other common causes of upper airway obstruction include edema of the oropharynx and larynx, trauma, foreign body, and infection.

211. Hypoxic hypoxia seen in

a) CO poisoning
b) HCN poisoning
c) Ischemia
d) AV shunt

A pulmonary shunt often occurs when the alveoli fill with fluid, causing parts of the lung to be unventilated although they are still perfused. Intrapulmonary shunting is the main **cause** of hypoxemia (inadequate blood oxygen) in pulmonary edema and conditions such as pneumonia in which the lungs become consolidated.

212. A 32 years' lady presents with shoulder tip pain and diagnosis is pan coast tumour and presents with meiosis. Diagnosis-

a) Horner syndrome
b) Thoracic outlet syndrome
c) Aberrant right subclavian artery
d) Upper trunk of brachial plexus injury

A disrupted nerve pathway on one side from the brain to the face and eye.
Horner's syndrome is the result of another medical condition, such as a stroke, tumour or spinal cord injury.
Symptoms include a smaller pupil, drooping eyelid and little or no sweating on the affected side.
There's no specific treatment for Horner's syndrome, but treatment for the underlying cause may restore normal nerve function.

213. A 85 year old patient was brought to the ER, BP: 180/100, right hemiparesis was seen. What is the next best step in management?

a) Reduce BP
b) NCCT
c) MRI
d) Aspirin 300mg and anticoagulants

Noncontrast head CT (NCCT) is **the standard radiologic test for patients presenting with acute stroke**. Early ischemic changes (EIC) are often overlooked

www.medicalcoupons.in

on initial NCCT. We determine the sensitivity and specificity of improved EH detection by a standardized method of image evaluation

214. Central obesity is seen in;

a) Cushing's disease
b) Chron's disease
c) Conns disease
d) Celiac disease

Central obesity in Cushing's syndrome. Fat tissue redistribution in CS leads to central obesity **and metabolic complications**. Visceral obesity is associated to altered adipokine secretion further contributing to insulin resistance, inflammation and fat accumulation.

215. Earliest finding in the diagnosis of diabetic nephropathy;

a) Microalbuminuria
b) Exudates
c) Hematuria
d) Macroalbuminuria

Microalbuminuria, urine albumin-**to-creatinine ratio >30 and less than <300 mg/g** has widely been accepted as biomarker of early diabetic kidney disease (DKD).

216. Reversible dementia seen in

a) Alzheimer's disease
b) Hypothyroidism
c) Subacute demyelination of spinal cord
d) Pick's disease

Hypothyroidism is one of the most important causes of potentially reversible dementia. The present case demonstrates a relatively rapid remission of cognitive and noncognitive symptoms within a period of a few months.

217. All are included in celiac sprue diet except:

a) Wheat
b) Barley
c) Rye
d) Quinoa

Celiac disease, sometimes called celiac sprue or gluten-sensitive enteropathy, is **an immune reaction to eating gluten**, a protein found in wheat, barley and rye. If you have celiac disease, eating gluten triggers an immune response in your small intestine.

www.medicalcoupons.in

218. All of the following are causes of hypercalcemia except

a) Hypervitaminosis of vit D
b) Hypeparathyroidism
c) Thiazides
d) Acute pancreatitis

Causes of hypercalcemia

- **Overactive parathyroid glands (hyperparathyroidism).** This most common cause of hypercalcemia can stem from a small, noncancerous (benign) tumor or enlargement of one or more of the four parathyroid glands.
- **Cancer.** Lung cancer and breast cancer, as well as some blood cancers, can increase your risk of hypercalcemia. Spread of cancer (metastasis) to your bones also increases your risk.
- **Other diseases.** Certain diseases, such as tuberculosis and sarcoidosis, can raise blood levels of vitamin D, which stimulates your digestive tract to absorb more calcium.
- **Hereditary factors.** A rare genetic disorder known as familial hypocalciuric hypercalcemia causes an increase of calcium in your blood because of faulty calcium receptors in your body. This condition doesn't cause symptoms or complications of hypercalcemia.
- **Immobility.** People who have a condition that causes them to spend a lot of time sitting or lying down can develop hypercalcemia. Over time, bones that don't bear weight release calcium into the blood.
- **Severe dehydration.** A common cause of mild or transient hypercalcemia is dehydration. Having less fluid in your blood causes a rise in calcium concentrations.
- **Medications.** Certain drugs — such as lithium, used to treat bipolar disorder — might increase the release of parathyroid hormone. Thiazides have several metabolic effects contributing to higher serum calcium levels, but **increased renal tubular reabsorption of calcium resulting in reduced urine calcium excretion** is the most likely cause
- **Supplements.** Taking excessive amounts of calcium or vitamin D supplements over time can raise calcium levels in your blood above normal.

219. Lady exposed to cold, extremities turn to blue associated with which antibody;

a) Anti scl 70
b) ANA
c) anti ssb
d) anti ssa

Raynaud's disease causes smaller arteries that supply blood flow to the skin to narrow in response to cold or stress. The affected body parts, usually fingers and toes, might turn white or blue and feel cold and numb until circulation improves, usually when you get warm
In a patient presenting with Raynaud phenomenon, a positive ANA test (even in the

absence of other symptoms) **warrants more frequent follow-up, urinalysis and** perhaps referral to a rheumatologist.

220. PR interval in ECG means

a) Reflects conduction through AV node
b) Ventricular Depolarisation
c) Ventricular Repolarisation
d) Atrial repolarization

The PR interval is **the time from the onset of the P wave to the start of the QRS complex**. It reflects conduction through the AV node

221. Lining of ventricles

a) Ependymal cells
b) Oligodendrocytes
c) Schwann cells
d) Astrocytes

Ependymal cells are mostly known as the cell type lining the brain ventricles. As non-neuronal cells in the brain and derived from neuroectoderm, they are clearly defined as a subtype of glial cells.

222. A patient is having continuous tremors and he has tendency to fall. Lesion is most commonly seen in;

a) Substantia nigra
b) Globus pallidus
c) Caudate
d) Putamen

When the neurons in the substantia nigra are damaged in large numbers, **the loss of dopamine prevents normal function in basal ganglia and causes the motor symptoms of PD: tremor, rigidity, impaired balance, and loss of spontaneous movement**.

223. Which type of gout is seen in a patient who is on treatment of CML?

a) Secondary gout
b) Pseudogout
c) Acute gout
d) Primary gout

Secondary gout refers to the **presence of a recognized cause or precipitating factor**, such as lymphoma (especially following chemotherapy), the excessive use of alcohol, or the use of diuretics. **an acquired rise in the serum urate that is directly**. **related to another disorder**.

www.medicalcoupons.in

224. Vasopressin works on the principle of;

a) Hyposmolality
b) Hyperosmolality
c) Hypovolemia
d) Hypervolemia

Vasopressin neurons in the supraoptic nucleus of the hypothalamus sense blood osmolarity. When osmolarity increases (e.g., after a meal, especially a salty one) vasopressin is secreted, **stimulating the kidney to reabsorb more water, reducing urinary volume**.

225. Which is not a component of brown sequard syndrome?

a) C/L pain
b) C/L temperature
c) C/L loss of proprioception
d) I/L paralysis

Brown-Séquard syndrome is a neurological syndrome resulting from hemisection of the spinal cord. It manifests with **weakness or paralysis and proprioceptive deficits on the side of the body ipsilateral to the lesion and loss of pain and temperature sensation on the contralateral side**.


226. Absent a waves in JVP is a feature of;

a) Atrial fibrillation
b) Atrial flutter
c) Ventricular tachycardia
d) Ventricular fibrillation

WAVEFORM	CAUSE
Cannon (irregular, large) *a* waves	Complete heart block
Absent *a* waves	Atrial fibrillation
Dominant *v* waves	Tricuspid regurgitation
Prominent *x* descent	Acute cardiac tamponade Constrictive pericarditis
Absent *x* descent	Atrial fibrillation
Sharp *y* descent	Severe tricuspid regurgitation Constrictive pericarditis
Slow *y* descent	Tricuspid stenosis Right atrial myxoma

227. 20-year-old male came to clinic and following findings were seen, what will be the diagnosis?

www.medicalcoupons.in

a) AML
b) ALL
c) CML
d) CLL

Acute myeloid leukemia (AML) is a cancer of the myeloid line of blood cells, characterized by the rapid growth of abnormal cells that build up in the bone marrow and blood and interfere with normal blood cell production.
Signs and symptoms of acute myelogenous leukemia include:

- Fever.
- Bone pain.
- Lethargy and fatigue.
- Shortness of breath.
- Pale skin.
- Frequent infections.
- Easy bruising.
- Unusual bleeding, such as frequent nosebleeds and bleeding from the gums.

228. What is the sequence of jaw thrust method?

a) Head tilt, chin lift, jaw thrust
b) chin lift, head tilt, jaw thrust
c) jaw thrust, chin lift, head lift
d) head lift, jaw thrust, chin lift

Head tilt, chin lift, jaw thrust is the correct sequence of jaw thrust method

229. What is the surgeon performing here?



a) Systolic pressure measurement on dorsalis pedis
b) Nerve conduction
c) Muscle tone
d) None

www.medicalcoupons.in

The surgeon is measuring systolic pressure measurement on dorsalis pedis

230. Endotracheal tube in the oesophagus is best assessed by,

a) CO2 Exhalation
b) Chest wall movement
c) Auscultation
d) chest tube passes through vocal cords

Capnography is the most reliable method to confirm endotracheal tube placement in emergency conditions in the prehospital setting.
Capnography is **the monitoring of the concentration or partial pressure of carbon dioxide (CO_2) in the respiratory gases**. Its main development has been as a monitoring tool for use during anesthesia and intensive care.

231. Identify the condition




a) Osteosarcoma
b) Ewings sarcoma
c) Osteoid osteoma
d) Osteochondroma

Osteosarcoma is **a type of bone cancer that begins in the cells that form bones**. Osteosarcoma is most often found in the long bones — more often the legs, but sometimes the arms — but it can start in any bone. In very rare instances, it occurs in soft tissue outside the bone.
The sunburst appearance occurs **when the lesion grows too fast and the periosteum does not have enough time to lay down a new layer and instead the Sharpey's fibers stretch out perpendicular to the bone**. It is frequently associated with osteosarcoma but can also occur with other aggressive bony lesions

www.medicalcoupons.in


232. What is the diagnosis of the given image?



a) Extradural haemorrhage
b) Subdural haemorrhage
c) Subarachnoid haemorrhage
d) Intracerebral haemorrhage

An extradural haematoma is **a collection of blood in the 'potential' space between the skull and the outer protective lining that covers the brain** (the dura mater). It usually occurs because of a head injury. It is a serious condition and emergency treatment is needed. An operation to remove the haematoma may be needed.
In almost all cases, extradural hematomas are seen on CT scans of the brain. They are typically **bi-convex (or lentiform) in shape**, and most frequently beneath the squamous part of the temporal bone. EDHs are hyperdense, somewhat heterogeneous, and sharply demarcated

233. PA view of chest X-ray is given here. What is the diagnosis?



a) Right Pneumothorax with right tracheal shift
b) Right Pneumothorax with left tracheal shift
c) Left Pneumothorax with left tracheal shift
d) Left Pneumothorax with right tracheal shift

What if one side of the chest cavity has an increase in pressure (such as in the case of a pneumothorax) the trachea will **shift towards the opposing side**. The trachea is the tube that carries air from the throat to the lungs.

www.medicalcoupons.in

234. A boy was riding bicycle, he fell down forward and injury to shoulder was seen, which nerve injury will be seen in the condition as given in X-ray?



a) Radial
b) Ulnar
c) Median
d) Axillary

all nerves of the brachial plexus are at risk of injury during glenohumeral dislocation, the most commonly injured is **the axillary nerve**

235. Post-menopausal women, fell down in washroom. What is the most common fracture she may suffer?



a) Colles fracture
b) Smith fracture
c) Monteggia fracture
d) Galeazzi fracture

the Colles fracture is defined as a **distal radius fracture with dorsal comminution**, dorsal angulation, dorsal displacement, radial shortening, and an associated fracture of the ulnar styloid. The term Colles fracture is often used eponymously for distal fractures with dorsal angulation

236. Football player with knee injury diagnosed as medial collateral ligament injury. which structure is most commonly associated with this type of injury?

www.medicalcoupons.in

a) Anterior cruciate ligament
b) Medial meniscus
c) Posterior Cruciate Ligament
d) Lateral lemniscus

A common injury in which forceful twisting causes certain tissue in the knee to tear. A meniscus tear occurs in the rubbery knee cartilage that cushions the shinbone from the thighbone. The meniscus can tear with forceful twisting or rotation of the knee.
Pain, swelling, stiffness and difficulty extending the knee are symptoms.
Treatment includes rest, ice, pain relievers and physiotherapy. Less commonly, surgery may be required.

237. Avascular necrosis of femur head is due to which of the following artery?

a) Medial circumflex femoral artery
b) Lateral circumflex femoral artery
c) Obturator artery
d) Profunda femoris artery

The medial circumflex femoral artery (internal circumflex artery, medial femoral circumflex artery) is an artery in the upper thigh that helps supply blood to the neck of the femur. Damage to the artery following a femoral neck fracture may lead to avascular necrosis (ischemic) of the femoral neck/head.

238. The shown apparatus is used for;



a) Ankle knee stabilizer
b) Patella tendon bearing brace
c) Thomas splint
d) None of the above

www.medicalcoupons.in

the brace was considered **to off-load tibia, fibula, and foot bones by transmitting the weight bearing loads via lateral upright**

239. Initial finding seen in knee TB

a. Reduction of joint space
b. Narrowing of joint space
c. Degeneration of cartilage
d. None of the above

The **Phemister triad** was named after American orthopedic surgeon Dallas Burton Phemister (1882–1951). The three features are juxta-articular osteopenia or osteoporosis, peripheral osseous erosions, and gradual narrowing of joint spaces—classically seen with tuberculous arthropathy.

240. A female presents with facial hair, secondary amenorrhea, increased levels of androgen and deranged LH: FSH ratio. What is the diagnosis?

a) PCOS
b) Endometriosis
c) Androgen insensitivity
d) Testicular feminization syndrome

A hormonal disorder causing enlarged ovaries with small cysts on the outer edges. The cause of polycystic ovary syndrome isn't well understood, but may involve a combination of genetic and environmental factors.
Symptoms include menstrual irregularity, excess hair growth, acne and obesity. Treatments include birth control pills to regularise periods, medication called metformin to prevent diabetes, statins to control high cholesterol, hormones to increase fertility and procedures to remove excess hair.

241. A child comes with the complaints of bed wetting. Treatment of choice-

a) Bed alarm
b) Behavioral therapy
c) Counselling
d) Anti-depressants

Bed alarm is the treatment of choice for nocturnal enuresis

242. Which is a product of purine metabolism?

a) Uric acid
b) Calcium
c) Phosphate
d) None of the above

www.medicalcoupons.in

Uric acid is the end product of purine metabolism in humans. Two-thirds of uric acid in the human body is normally excreted through the kidney, whereas one-third undergoes uricolysis (decomposition of uric acid) in the gut.

243. Giant cell pneumonia is associated with:

a) CMV
b) Herpes Simplex
c) Herpes Zoster
d) Measles

Giant cell pneumonia: A deadly but fortunately rare complication of measles that tends to strike children who are immunodeficient from leukemia or AIDS. The lung tissue shows multinucleated giant cells lining the alveoli (air sacs) of the lungs. Also known as Hecht's pneumonia.

244. Investigation of choice for the following



a) X Ray
b) CT
c) MRI
d) HRCT

coxa vara usually presents with a limp, a leg length difference, and limited ability to bring the thigh out to the side. **X-rays** are used to diagnose coxa vara. Depending on what caused the condition and how severe the deformity, reconstructive surgery may be needed to improve the hip alignment and biomechanics.

245. A 35-year-old anemic female patient came for family planning, she has 2 living children and history of ectopic pregnancy what is your next step:

www.medicalcoupons.in

a) Lippe's Loop
b) Hysterectomy
c) T Piece
d) Progestasert

The Progestasert system combines the advantageous features of IUDs and oral minidose progestogen preparations. An internal device continuously delivers progesterone for 1 year to the uterine lumen and endometrium. A T-shaped Progestasert which releases 65 mcg/day has been selected for wide scale clinical use.

246. Reduviid bug **transmits which disease:**

a) Lyme disease
b) Chagas Disease
c) Anaplasmosis
d) Powassan Virus Disease


Chagas disease, also known as American trypanosomiasis, is caused by the parasite Trypanosoma cruzi. It is transmitted to humans via the reduviid bug (the "kissing bugs"), and is therefore characterized as a zoonotic disease.

247. Ectopic pregnancy lasts longest at which site:

a) Ampulla
b) Isthmus
c) Infundibular
d) Interstitium

An ectopic pregnancy occurs when a fertilized egg grows outside of the uterus. Almost all ectopic pregnancies—more than 90%—occur in a fallopian tube. As the pregnancy grows, **it can cause the tube to burst** (rupture). A rupture can cause major internal bleeding. Very late rupture of tubal pregnancy is seen when the implantation is at Interstitium.

248. A female came with complain of severe abdominal pain since morning. She mentions the pain didn't subside on medications. She has a history of amenorrhea for 2 months. Following was viewed by the surgeon. What is your diagnosis:



a) **Ectopic Pregnancy**
b) Chocolate Cyst
c) Ovarian Cyst
d) None of the above

A pregnancy in which the fertilised egg implants outside the uterus.
The fertilised egg can't survive outside the uterus. If left to grow, it may damage nearby organs and cause life-threatening loss of blood.
Symptoms include pelvic pain and vaginal bleeding.
To prevent complications, treatment is required. In the early stages, medication may be sufficient. Later stages require surgery.

All the following leads to renal artery stenosis except:

a) **Polyarteritis nodosa**
b) Fibromuscular dysplasia
c) Atherosclerosis
d) Takayasu Arteritis

the two main causes of renal artery stenosis include:

- **Buildup on kidney (renal) arteries.** Fats, cholesterol and other substances (plaque) can build up in and on your kidney artery walls (atherosclerosis). As these deposits get larger, they can harden, reduce blood flow, cause kidney scarring and eventually narrow the artery. Atherosclerosis occurs in many areas of the body and is the most common cause of renal artery stenosis.
- **Fibromuscular dysplasia.** In fibromuscular dysplasia, the muscle in the artery wall grows abnormally, often from childhood. The renal artery can have narrow sections alternating with wider sections, giving a bead-like appearance in images of the artery.

249. After cotrimoxazole administration patient develops hyperpigmentation over back as shown below. What is your diagnosis?



www.medicalcoupons.in

a) **Fixed Drug Eruption**
b) DRESS Syndrome
c) Toxic Epidermal Necrolysis
d) Steven Johnson Syndrome

Fixed drug eruption (FDE) is a **distinctive type of cutaneous drug reaction** that characteristically recurs in the same locations upon reexposure to the offending drug.
Fixed drug eruption typically presents as a **single (or small number of) well-defined, round or oval red or violaceous patch or plaque which may blister or ulcerate**. It is usually asymptomatic but can be itchy or painful.

250. **A young male presents with recurrent bouts of diarrhoea, two hours after drinking milk. Diagnosis**?

a) Gastrocolic Reflex
b) **Lactase deficiency**
c) Inflammatory Bowel Disease
d) IBS

The inability to fully digest sugar (lactose) in dairy products.
Lactose intolerance is usually caused by a deficiency of an enzyme in the body called lactase.
Symptoms can include abdominal cramps, bloating and diarrhoea.
Treatment focuses on avoidance of dairy products, use of lactose-free products or the use of lactase supplements.

251. Cotrimoxazole is given for management in all of the following except?

a) **Actinomycosis**
b) Toxoplasma Gondii
c) Pneumocystis Jiroveci
d) Isospora Belli

In most cases of actinomycosis, antimicrobial therapy is the only treatment required, although surgery can be adjunctive in selected cases. **Penicillin G** is the drug of choice for treating infections caused by actinomycetes

252. Which of the following has a late diastolic murmur?

a) Atrial regurgitation
b) **Atrial Myxoma**
c) Pulmonary regurgitation
d) Pulmonic Stenosis

www.medicalcoupons.in

Late diastolic (presystolic) murmurs start after S2 and extend up to S1 and have a crescendo configuration. They can be associated with AV valve narrowing. They include **mitral stenosis, tricuspid stenosis, myxoma, and complete heart block**.

253. A 3-year-old patient is having loose motions with weight loss. Small intestinal biopsy was done. Which of the following tests is best suited for clinical diagnosis of the patient?

a) Anti-Mitochondrial AB
b) Anti-Gliadin AB
c) Anti-SCL 70 AB
d) Anti-Microsomal AB

A gliadin antibodies test is used to help **doctors diagnose celiac disease**. Celiac disease is an autoimmune disorder in which the body's immune system mistakenly thinks that gluten — a protein in wheat, barley, rye, and oats

254. A 40-year old smoker patient has been diagnosed with infra-renal aorta aneurysm of 4 cm. What is the best treatment?

a) Urgent Resection and Anastomosis
b) Intraluminal Graft Placement
c) Endoscopy
d) Periodic Size Measurement

255. A 30-year old patient is having weakness due to secretory diarrhoea. Endoscopy shows large duodenal ulcer. Probable diagnosis is?

a) Gastrinoma
b) Primitive Neuroectodermal Tumour
c) Carcinoid Syndrome
d) Autoimmune Gastritis

A gastrinoma is a **gastrin-secreting tumor** that can occur in the pancreas, although it is most commonly found in the duodenum
The signs and symptoms of gastrinomas are related to increased gastric acid production, this is caused by the excess gastrin hormone that is produced by the tumour cells. Most patients experience **abdominal pain, heartburn, nausea, diarrhoea, blood in the vomit and/or bleeding from the back passage**.

256. Patient presents with MI, STK (streptokinase) was given. What should be checked in order to monitor toxicity?

a) Bleeding Time
b) Fibrin degradation products
c) Platelet Count
d) Thrombin Time

www.medicalcoupons.in

Careful monitoring for arrhythmia is recommended during and immediately following administration of **Streptase**, Streptokinase, for acute myocardial infarction. Occasionally, tachycardia and bradycardia have been observed.

257. In which of the following conditions JVP increases inspiration?

a) Pericardial Effusion
b) Cardiac Tamponade
c) Aortic Dissection
d) Constrictive Pericarditis

Elevated jugular venous pressure (JVP) is present in virtually all patients that are not hypovolemic. Pericardial constraint **results in the inability of the right heart to accommodate inspiratory abdominal venous return**, translating to an inspiratory increase in the JVP (Kussmaul's sign)

258. What is the best treatment for a 35-year old AIDS positive patient with Cryptosporidium Diarrhoea?

a) Nitazoxanide
b) Metronidazole
c) Parmomycin
d) Cotrimoxazole

Nitazoxanide, paromomycin, and azithromycin have activity against Cryptosporidium. A 3-day course of nitazoxanide oral suspension has been approved by the FDA for the treatment of cryptosporidiosis-related diarrhea in adults and in children older than 12 months who do not have HIV infection.

259. Which of the following is Monosynaptic reflex?

a) Withdrawal Reflex
b) Crossed Extensor Reflex
c) Peristaltic Reflex
d) Stretch Reflex

The monosynaptic stretch reflex, or sometimes also referred to as the muscle stretch reflex, deep tendon reflex, is a reflex arc that **provides direct communication between sensory and motor neurons innervating the muscle**. ... This contraction allows the muscle to resist the force that initially caused the reflex.

www.medicalcoupons.in

FMGE AUG 2020

1. **Choose the correct sequence of muscle contraction in ejaculation**
   a) **Ischiocavernosus contraction → urethralis→ Bulbospongiosus**
   b) Ischiocavernosus contraction → Bulbospongiosus → urethralis
   c) Bulbospongiosus → urethralis → Ischiocavernosus
   d) Bulbospongiosus Ischiocavernosus → urethralis

During the emission phase, the muscles around the epididymis and ductus deferens (the tube extending from the epididymis) contract to push the sperm into the prostate and urethra. During ejaculation, the semen is expelled by strong spasmodic contractions of the bulbocavernosus muscle, which encircles the corpus spongiosum (the structure in the penis that encloses the urethra). The whole process of ejaculation is accomplished by nerve impulses received from the penis; once ejaculation is started it becomes a reflex reaction that cannot be voluntarily interrupted.

2. **Deep inguinal ring is a defect in:**
   a) External oblique muscle aponeurosis
   b) Internal oblique muscle aponeurosis
   c) Transversus abdominis muscle aponeurosis
   d) **Transversalis fascia**


The deep (internal) inguinal ring is a defect in the transversus abdominis fascia that allows the contents to enter the canal and leave the abdomen before extending obliquely, medially and inferiorly through the canal to exit at the superficial (external) inguinal ring, a defect in the external oblique fascia.

3. **Name the skin receptor responsible for detecting rapid vibration sense:**
   a) Meissner corpuscle
   b) **Paccinian corpuscle**
   c) Merkel cell
   d) Ruffini's corpuscle

The Pacinian corpuscle or Vater-Pacinian corpuscles or Lamellar corpuscles in the skin and fascia detect rapid vibrations of about 200–300 Hz.

4. **Which type of joint is present at the arrow mark shown in the diagram:**

   a) Primary cartilaginous
   b) **Secondary cartilaginous**
   c) Fibrous
   d) Synovial



www.medicalcoupons.in

Secondary cartilaginous joint

- symphysis pubis between the right and left pubic bones.
- manubriosternal joint between the sternal body and the manubrium.
- intervertebral discs.
- sacrococcygeal symphysis.

5. **A lesion in left sided motor nucleus of facial nerve leads to:**

a) Right sided loss of wrinkles on forehead and right side corner of mouth goes down

b) Right sided loss of wrinkles on forehead and right side corner of mouth goes up

c) Left sided loss of wrinkles on forehead and left side corner of mouth goes down

d) Left sided loss of wrinkles on forehead and right side corner of mouth goes up

For example, a left corticobulbar lesion results in paralysis of the muscles that control the lower right quadrant of the face. By contrast, a lower motor neuron lesion to the facial motor nucleus results in **paralysis of facial muscles** on the same side of the injury.

6. Structure developing in dorsal mesogastrium:

a) Lesser omentum

b) Falciform ligament

c) Spleen

d) Liver

The liver develops in the ventral mesogastrium; **the spleen** develops in the dorsal mesogastrium. The liver grows rapidly, pressing against the body wall, and obliterating these layers of peritoneum. These changes produce this almost separate pocket behind the stomach, the lesser sac.

7. A patient developed the following deformity, after overhead weight-lifting exercises. Which of the following muscle is involved?



a) Latissimus dorsi

b) Trapezius

c) Serratus anterior

d) Supraspinatus

www.medicalcoupons.in

Scapular winging is almost always caused by damage to one of three nerves that control muscles in your arms, back, and neck: the long thoracic nerve, which controls the **serratus anterior muscle**.

8. Contralateral homonymous hemianopia with macular sparing occurs due to lesion at the level of?

a) Optic nerve
b) Optic tract
c) Optic radiation
d) Orbital cortex

Homonymous hemianopsia can be congenital, but is usually caused by brain injury such as from stroke, trauma,[5] tumors, infection, or following surgery.

Vascular and neoplastic (malignant or benign tumours) lesions from the optic tract, to visual cortex can cause a contralateral homonymous hemianopsia. Injury to the right side of the brain will affect the left visual fields of each eye. The more posterior the cerebral lesion, the more symmetrical (congruous) the homonymous hemianopsia will be.

9. Posterior tibial artery pulse is felt:

a) Between medial malleolus and medial border of tendo-calcaneus in inversion
b) Between medial malleolus and medial border of tendo-calcaneus in eversion
c) Between medial malleolus and lateral border of tendo-calcaneus in inversion
d) Between medial malleolus and lateral border of tendo-calcaneus in inversion

The posterior tibial artery enters the sole of the foot by passing deep to the flexor retinaculum. Its pulsation can be felt **midway between the medial malleolus and the medial border of the tendocalcaneus**. The pulsation of the peroneal artery is felt in front of the lateral malleolus at its medial border.

10. Patient underwent removal of left submandibular salivary gland, and lingual nerve was damaged. Which of the following statements is NOT correct?

a) Sublingual gland secretions are reduced
b) Previous 2 / 3rd tongue taste sensation is lost
c) Sensation at the floor of mouth lost
d) Tongue deviated to left side

www.medicalcoupons.in

If your lingual nerve sustains an injury, you'll most likely experience any of these nerve damage symptoms: Changed sensation in the tongue, chin, or lower lip areas (similar to sensations you feel when your oral cavity is numbed for a dental procedure or as the anesthesia slowly wears off) **Altered ability to taste**. If your lingual nerve sustains an injury, you'll most likely experience any of these nerve damage symptoms: Changed sensation in the tongue, chin, or lower lip areas (similar to sensations you feel when your oral cavity is numbed for a dental procedure or as the anesthesia slowly wears off) **Altered ability to taste**.

11. Ductus arteriosus develops from which pharyngeal arch artery?

a) 3
b) 4
c) 5
d) 6

Ductus arteriosus itself arises from the **left 6th pharyngeal arch artery**.

12. Which enzyme deficiency is seen in TaySach's disease?

a) Hexosaminidase A
b) Glucocerebrosidase
c) G-6-PD
d) Hexosaminidase A & B

Tay-Sachs disease is a rare, neurodegenerative disorder in which deficiency of an enzyme (hexosaminidase A) results in excessive accumulation of certain fats (lipids) known as gangliosides in the brain and nerve cells.

13. An Alcoholic patient presented with mental confusion, gait problem and ophthalmoplegia. Which vitamin deficiency is seen?

a) B1
b) A
c) B6
d) B12

Wernicke-Korsakoff syndrome is a condition caused by a deficiency of **thiamine or vitamin B1**. Patients classically present with a clinical triad of ophthalmoplegia, altered mental status, and ataxia.

www.medicalcoupons.in

14. A female patient with macrocytic anaemia, Homocystinuria & normal methylmalonate levels. Which vitamin therapy can help in this?

a) Folate plus Vitamin B12

b) Folate

c) Vitamin A

d) Vitamin C

Vitamin B12 or B9 (commonly called folate) deficiency anaemia occurs when **a lack of vitamin B12 or folate causes the body to produce abnormally large red blood cells** that cannot function properly. Red blood cells carry oxygen around the body using a substance called haemoglobin. Additional testing with homocysteine and MMA determinations may help distinguish between B12 and folate deficiency states. In folate deficiency, homocysteine levels **are elevated and MMA levels are normal**. In vitamin B12 deficiency, both homocysteine levels and MMA levels are elevated.

15. Macrocytic anaemia case with normal methyl-malonyl with increased level of homocysteine. What is the diagnosis?

a) B9 deficiency

b) B12, 9 deficiency

c) Iron deficiency

d) Anaemia of chronic ds

Vitamin B12 or B9 (commonly called folate) deficiency anaemia occurs when **a lack of vitamin B12 or folate causes the body to produce abnormally large red blood cells** that cannot function properly. Red blood cells carry oxygen around the body using a substance called haemoglobin. Additional testing with homocysteine and MMA determinations may help distinguish between B12 and folate deficiency states. In folate deficiency, homocysteine levels **are elevated and MMA levels are normal**. In vitamin B12 deficiency, both homocysteine levels and MMA levels are elevated.

16. A heart patient suddenly has chest pain. Which isoenzyme of LDH enzyme is elevated?

a) LDH -1

b) LDH -5

c) LDH -2

d) LDH -4

It is known that the ratio of **isoenzyme 1** to total lactate dehydrogenase (LDH1) in serum is increased in all patients with acute myocardial infarction within 24 h of the infarct.

www.medicalcoupons.in

17. Cofactor for Mitochondrial SOD (Superoxide Dismutase):

a) Mn

b) Cu

c) Zn

d) Mb

Manganese superoxide dismutase (MnSOD), an **important primary antioxidant enzyme** located in mitochondria, is essential for the removal of superoxide radicals constantly being generated by the electron transport chain (ETC).

18. Nitric oxide is synthesized from:

a) L-Hydroxy proline

b) Arginine

c) D-Histidine

d) Cysteine

Nitric oxide (NO), synthesized from the amino acid l-arginine by NO synthases, is a unique type of transmitter in the nervous system.

19. Which of the following is involved in protein folding?

a) Chaperone

b) Ubiquitin

c) Proline

d) Arginine

Chaperones are a **group of proteins that have functional similarity and assist in protein** folding. They are proteins that have the ability to prevent non-specific aggregation by binding to non-native

20. Which of the following is not given in a case of fatty liver disease?

a) Choline

b) Ethanol

c) Folic acid

d) Methionine

Recent studies indicate that additional effects of ethanol impair fat oxidation as well as stimulate lipogenesis. The effects of ethanol on lipid metabolism result from **inhibition of PPAR-□ and stimulation of SREBP-1**, resulting in metabolic remodelling of the liver toward a fat-storing, rather than fat oxidizing organ.

21. 16 Years old girl was kidnapped. Police caught the accused and brought to the police station. After some time, the accused was found dead in the station. Inquest in this case should be done by?

www.medicalcoupons.in

a) Executive Magistrate.
b) Judicial Magistrate.
c) Police.
d) No inquest needed.

Death under police custody case is inquested by Judicial Magistrate, also he inquests
Death in a psychiatric hospital.
Death in jail.
Death as a result of police shooting.

22. **Dowry Death inquest is according to which CrPC?**

a) 174
b) 175
c) 176
d) 304-B

According to **Section 304-B**, to make out a case of dowry death, a woman should have died of burns or other bodily injuries or "otherwise than under normal circumstances" within seven years of her marriage.


23. **Identify the fingerprint pattern shown in image**



a) Loop
b) Whorl
c) Arch
d) Composite

www.medicalcoupons.in

24. Identify the test that shows the following crystals.



a) Teichman
b) **Takayama**
c) Florence
d) Barberios

Takayama test is **a confirmation test used to detect blood spots**. Based on the research results this test can still be used to identify dried blood spots on clothing aged 20 years, and able to detect positively the presence of blood with the formation of pink crystals.

25. Which of the following is a true statement about dying declaration?

a) **If victim survives it is as such inadmissible**
b) Oath is administered while recording it
c) Ideal person to record DD is the attending doctor
d) Alteration of terms / phrases of dying person is allowed

A dying declaration is considered credible and trustworthy evidence based upon the general belief that most people who know that they are about to die do not lie. ... A person who makes a dying declaration **must**, however, be competent at the time he or she makes a statement, otherwise, it is inadmissible.

26. In regard to a fetal death classification, a test was done to determine the cause which includes dipping the lung in to the water. Identify the test.

a) Foderes
b) Ploquets
c) Hydrostatic

www.medicalcoupons.in

d) Wredins

The hydrostatic test, or floatation test, has historically been used to determine whether a newborn infant has breathed. It is performed by placing the lungs, still attached to the heart, in water and seeing whether they float.

27. A patient is brought to casualty with history of snake bite with features as shown. During course of Treatment, the patient expires. The accused snake's photo was captured by relatives as shown in image. What could be the cause of death in this case?



a) Shock
b) Respiratory paralysis
c) Circulatory failure
d) Rhabdomyolysis

The venom of common krait contains 3 major neurotoxins (α, β and κ-bungarotoxin) that can induce failure of neuromuscular transmission resulting in respiratory muscles paralysis.

28. Identify the method of solvent abuse done with use of a plastic bag applied to mouth & nose.



a) Bagging
b) Huffing
c) Gliding
d) Sniffing

www.medicalcoupons.in

"bagging" — sniffing or inhaling fumes from substances sprayed or deposited inside a plastic or paper bag

29. Sexual gratification obtained through obscene telephone calls is called as;

a) Scophophilia
b) Scoptophilia
c) Scatologia
d) Voyeurism

Scatologia, also called Coprolalia, deviant sexual practice in which sexual pleasure is obtained through the compulsive use of obscene language. The affected person commonly satisfies his desires through obscene telephone calls, usually to strangers.

30. A person was found dead in his apartment and the forensic team observed the scene and gave these noted details regarding injury that caused death. The injury over the vertebra had clean cut margins but the bones appeared to be crushed. What could be the correct option in this scenario?

a) Incised
b) Chop
c) Lacerated
d) Stab

31. Rotten egg odour is associated with

a) Arsenic
b) Hydrogen sulphide
c) Aluminium phosphide
d) Phosphorus

Hydrogen sulphide (H_2S) is a colourless gas with a characteristic odour of rotten eggs which being denser than air may pool in low areas in still conditions.

32. Minimum number of Antenatal visits required are;

a) 3
b) 4
c) 5
d) 6

www.medicalcoupons.in

For women whose pregnancies are progressing normally, WHO recommends a minimum of **four ANC visits**, ideally at 16 weeks, 24-28 weeks, 32 weeks and 36 weeks

33. A man came for health check-up after his father had a cerebrovascular accident who died last month. He was having the history of hypertension. Which type of prevention is this?

a) Primordial
b) Primary
c) Secondary
d) Specific protection


Secondary Prevention - **trying to detect a disease early and prevent it from getting worse**. Tertiary Prevention - trying to improve your quality of life and reduce the symptoms of a disease you already have

34. Mid-year population is taken on

a) 01 September
b) 01June
c) 01 April
d) 01 July

The date of the midyear population is **July 1**. Note that the midyear month is different from the census month. These differences are taken into account when calculating.

35. Identify the Statistical diagram shown below



a) Histogram
b) Bar diagram
c) Frequency polygon
d) Scatter diagram

www.medicalcoupons.in

Histogram is a diagram consisting of rectangles whose area is proportional to the frequency of a variable and whose width is equal to the class interval.

36. Identify the Logo shown below?



a) AGMARK
b) PFA standards
c) Food Standards and Safety Authority of India (FSSAI)
d) Bureau of Indian Standards (BIS)

BIS is the National Standard Body of India established under the BIS Act 2016 for the **harmonious development of the activities of standardization**, marking and quality certification of goods and for matters connected therewith or incidental thereto.

37. Regarding the School health service guidelines recommendations, which of the following statements is true?

a) Minimum area 5 sq. ft. per student
b) Minus type desk
c) Can allot 60 students per classroom
d) Light coming from front of the expressed glass

38. **Researcher was conducting study in relation of Depression associated with history of social media usage. One group had social media users with depression and another group had social media users without depression. Which type of study design is used?**

a) Cohort study
b) Case control study
c) Cross sectional study
d) Random sampling

A cross-sectional study involves looking at data from a population at one specific point in time. The participants in this type of study are selected based on particular variables of interest.

www.medicalcoupons.in

39. Delphi technique used for collective opinion of;

a) Population

b) Group

c) Community

d) Individual

The Delphi method is a process used to arrive at a group opinion or **decision by surveying a panel of experts**. ... The experts can adjust their answers each round, based on how they interpret the "group response" provided to them. The ultimate result is meant to be a true consensus of what the group thinks.

40. **A Researcher did study on staff, nurses and junior resident doctors doing yoga (Divided into male and female). Yoga shows 25% decreased risk of infections. Which is the best test to test the significance of the result?**

a) ANOVA

b) Chi square test

c) Z test

d) Correlation

chi-square test

relating to or denoting a statistical method assessing the goodness of fit between observed values and those expected theoretically. Here the researcher has 2 groups where 25% decreased risk of infections when they do yoga i.e. 25% less than the risk expected theoretically

41. A Researcher undertakes follow-up study for a certain time period duration, and tells that the people increasingly develop the disease with time, then it expressed as,

a) Incidence

b) Cumulative incidence

c) Prevalence

d) Relative risk

Cumulative incidence is calculated as **the number of new events or cases of disease divided by the total number of individuals in the population at risk for a specific time interval**. Researchers can use cumulative incidence to predict risk of a disease or event over short or long periods of time.

42. All of following are Principles of Primary Health Care Except?

a) Community participation

b) Appropriate facility

c) Intersectoral coordination

d) Equitable distribution

www.medicalcoupons.in

Five (5) Principles of Primary Health Care (PHC):

- Social equity.
- Nation-wide coverage/wider coverage.
- Self- reliance.
- Intersectoral coordination.
- People's involvement (in planning and implementation of programs)

43. In different types of graphs, Secular trend of a Disease is best represented by;

a) Line diagram
b) Bar graph
c) Box and whiskers plot
d) Stem leaf plot

The secular trend describes **the occurrence of disease over a prolonged period**, usually years; it is influenced by the degree of immunity in the population and possibly nonspecific measures such as improved socioeconomic and nutritional levels among the population and are usually represented by line diagram.

44. Which of the following depicts Severity / lethality of disease?

a) Proportional mortality rate
b) Case specific death rate
c) Case fatality rate
d) Total deaths due to that disease

Case fatality rate, also called **case fatality risk** or **case fatality ratio**, in epidemiology is the proportion of people who die from a specified disease among all individuals diagnosed with the disease over a certain period of time. Case fatality rate typically is used as a measure of disease severity and is often used for prognosis (predicting disease course or outcome)

45. In a hospital, systematic observation and recording of doctors spending time with patients care and time without patients' care is calculated for future management purposes. The type of management technique used here is?

(a) Network analysis
(b) System analysis
(c) Work sampling
(d) Decision making

www.medicalcoupons.in

Work sampling is the statistical technique used for determining the proportion of time spent by workers in various defined categories of activity (e.g. setting up a machine, assembling two parts, idle…etc.) It is as important as all other statistical techniques because it permits quick analysis, recognition, and enhancement of job responsibilities, tasks, performance competencies, and organizational work flows

46. Definition of blindness, as per WHO;

a) <1/60 in better eye
b) <1/60 in worse eye
c) <3/60 in better eye
d) <3/60 in worse eye

'Blindness' is defined as **visual acuity of less than 3/60**, or a corresponding visual field loss to less than 10°, in the better eye with the best possible correction.

47. A 8-month-old child reported to a PHC with runny nose and fever since last 3 days. Rash started on face and then covered the trunk part of body. What's the most likely diagnosis?

a) Measles
b) Chicken pox
c) Mumps
d) Rubella

Measles symptoms appear **7 to 14 days after contact** with the virus and typically include high fever, cough, runny nose, and watery eyes. Measles rash appears 3 to 5 days after the first symptoms.

48. Which of the following is NOT constrained by Time or Existing resources?

a) Vision
b) Target
c) Goal
d) Mission

Goal is not constrained by Time or Existing resources

49. Newborn Health Mission component do not include;


a) Target of infant mortality to double digit by 2030
b) Antenatal care
c) Postnatal visits
d) Care during labor

www.medicalcoupons.in

NHM has six financing components:

- NRHM-RCH Flexipool.
- NUHM Flexipool.
- Flexible pool for Communicable disease.
- Flexible pool for Non communicable disease including Injury and Trauma.
- Infrastructure Maintenance and.
- Family Welfare Central Sector component.

50. **Identify the Contraceptive shown in the Photograph given below**



a) CuT 7
b) CuT 220B
c) **CuT 380A**
d) NOVA-T

CuT 380A - It is a T-shaped device with a polyethylene frame holding 380 mm2 of exposed surface area of copper. The IUD frame contains barium sulfate thus making it radio-opaque.

51. If you are posted as a Medical officer in an area under NVBDCP, and you will be advised to spray Malathion for malaria prophylaxis. What could be the probable decision you will take regarding frequency of Malathion spray?

a) Once very fortnight
b) Once every 3 months
c) Twice every3 months
d) Once every month

www.medicalcoupons.in

Frequency of malathion spray under NVBDCP is once every 3 months.

52. A country is suspecting a severe disease. What's the probable time within which one should report it to WHO?

a) 6 hours
b) 12 hours
c) 24 hours
d) 48 hours

the probable time within which one country should report it to WHO is 24 hrs

53. Least Iron content is present in;

a) Beans
b) Milk
c) Spinach
d) Liver

Both cow's milk and breastmilk are low in iron (around **0.2-0.5 mg iron per litre**), however the iron in breastmilk is bonded to lactoferrin and offers greater bioavailability.

54. In an operation theatre the nurse spilled blood on the floor. What can be used to disinfect it?

a) Cresol
b) 70% Ethyl alcohol
c) Glutaraldehyde
d) 1% Sodium hypochlorite

Wipe the area with water and detergent until it is visibly clean. Saturate the area again with sodium hypochlorite 1% (10 000 ppm available chlorine). This is a 1:10 dilution of 5.25% sodium hypochlorite bleach, which should be prepared daily. Rinse off the tongs, brush and pan, under running water and place to dry.

55. In a train accident, there were 74 dead, 64 were severely injured, 20 moderately injured and 32 mildly injured. Color coding in Triage Highest to lowest category is

a) Green Yellow Red Black
b) Yellow Red Green Black
c) Black Yellow - Red Green
d) Red - Yellow - Green – Black

www.medicalcoupons.in

RED: (Immediate) severe injuries but high potential for survival with treatment, taken to collection point first.

YELLOW: (Delayed) serious injuries but not immediately life-threatening.

GREEN: (Walking wounded) minor injuries

BLACK: morbid patients

56. **A healthy person get in contact with an infected case and they have to separate him not more than the incubation period of disease. This is called as;**

a) Absolute quarantine

b) Isolation

c) Interruption of transmission

d) Serial interval

Absolute or complete quarantine: The **limitation of freedom of movement of those exposed to a communicable disease** for a period of time not longer than the longest usual incubation period of that disease, in such manner as to prevent effective contact with those not so exposed.

57. A male underwent Vasectomy and after 3 months his wife got pregnant. What advice should have been given to him post-vasectomy?

a) Abstinence for 3 months after the procedure

b) Usage of Barrier methods for 3 months after the procedure

c) OCPS usage for I month

d) Female condom usage for 1 month

Sperms can be seen in semen for up to 3 months' post ligation, so repeat semen analysis **after 3 months and if azoospermia recorded** then he can have unprotected intercourse

Post-procedural semen analysis at 8 to 16 weeks to document sterility.

58. First investigation for TB Diagnosis?

a) Chest X-ray

b) Sputum smear examination

c) CB-NAAT

d) Liquid culture

A definitive diagnosis of tuberculosis can only be made by culturing Mycobacterium tuberculosis organisms from a specimen taken from the patient (most often sputum,

www.medicalcoupons.in

but may also include pus, CSF, biopsied tissue, etc.). A diagnosis made other than by culture may only be classified as "probable" or "presumed". For a diagnosis negating the possibility of tuberculosis infection, most protocols require that two separate cultures both test negative

59. In ORS, Sodium is given along with Glucose for

a) Reducing secretions
b) Osmosis
c) Facilitative diffusion
d) Co-transport

Composed of glucose and sodium and other electrolytes, ORS promote sodium and thus **water absorption in part** via passive sodium-coupled glucose transport in intestinal villi.

60. A factory worker had a history of frequent exposure to groundnuts. Subsequently he develops Hepatocellular carcinoma. Most likely exposure association is

a) Ergot toxin
b) Aflatoxin
c) Sanguinarine
d) Pyrrolizidine alkaloids

Aflatoxin Alert: Moldy Nuts and Corn Increases Liver Cancer Risk 60-Times If You Have Hepatitis B. One of the biggest health threats to people living with chronic hepatitis B is a toxic, nearly invisible mold called aflatoxin found in corn, peanuts, peanut butter, almonds, Brazil nuts, walnuts and pistachios.

61. A Medical officer examines workers in mines and suspected Silicosis. What should be his comments?

a) It is reversible once we remove the cause
b) It takes 6-10 years to develop
c) Fibrotic changes in the lungs of the patient can be reversed
d) TB screening should be done at regular intervals

Long time workers in mines have high risk of TB and can be suspected to have silicosis, So TB screening should be done at regular intervals.

62. A person returning from malaria endemic country should continue chemoprophylaxis for how long?

www.medicalcoupons.in

a) 1 week
b) 15 days
c) 4 weeks
d) 4 months

All prophylactic drugs should be taken with unfailing regularity for the duration of the stay in the malaria risk area, and should be continued for **4 weeks after the last possible** exposure to infection since parasites may still emerge from the liver during this period.

63. An infant is given Buffalo milk. It is superior because;
a) High calories and high protein
b) Low calories and high protein
c) High calories and low protein
d) Low calories and low protein

Buffalo milk is **rich in vitamin A**, has a higher protein value and contains more iron, calcium and phosphorus than the cow's milk. ... It is better to introduce buffalo milk slowly.

64. Long term changes / sequelae of a disease are seen in
a) Secular trend of a disease
b) Cyclical trend of a disease
c) Disease changing its traits according to herd immunity in the population
d) Disease changing the symptoms as per seasons

Long term changes of a disease are seen in secular trend. The secular trend describes **the occurrence of disease over a prolonged period**, usually years; it is influenced by the degree of immunity in the population and possibly nonspecific measures such as improved socioeconomic and nutritional levels among the population.

65. A 5-year-old child came into OPD with fever, rashes on the body. There were rashes on the axilla and flexor surface with various macules, papules and vesicles. Rashes and blisters are in different stages and non- uniform. Most probable diagnosis is?
a) Chickenpox
b) Smallpox
c) Epstein Barr
d) Measles

www.medicalcoupons.in

Chickenpox is **an infection caused by the varicella-zoster virus**. It causes an itchy rash with small, fluid-filled blisters. Chickenpox is highly contagious to people who haven't had the disease or been vaccinated against it

66. Regular monitoring of visual acuity and fundus examination is required with the patient is being treated with which of the following DMARD?

a) Hydroxychloroquine
b) Methotrexate
c) Sulfasalazine
d) Leflunomide

All individuals who have taken hydroxychloroquine for greater than 5 years should receive **annual screening** for retinopathy. All individuals who have taken chloroquine for greater than 1 year should receive annual screening for retinopathy

67. A female presented with burning micturition and symptoms of UTI. Which fluoroquinolones is least likely to be effective?

a) Ofloxacin
b) Moxifloxacin
c) Ciprofloxacin
d) Levofloxacin

Gemifloxacin and moxifloxacin are **not effective for the treatment of UTIs** because they do not achieve adequate concentrations in the urine. 5-7 The fluoroquinolones indicated for UTI are excreted in the urine as greater than 40% unchanged drug while gemifloxacin is less than 35% and moxifloxacin is only 20%.

68. A patient with heart disease presented with breathlessness the features of the patient shown in images. Which of the following is used to treatment of oedema for this patient?





www.medicalcoupons.in

a) ACE Inhibitor
b) Na – K – 2Cl symporter inhibitor in loop of Henle
c) Na – Cl symporter inhibitor in DCT
d) Aldosterone in DCT

Furosemide is a loop diuretic (water pill) that **prevents your body from absorbing too much salt**. This allows the salt to instead be passed in your urine. Furosemide is used to treat fluid retention (edema) in people with congestive heart failure, liver disease, or a kidney disorder such as nephrotic syndrome.

69. A patient with pancreatic carcinoma is on morphine for treatment of pain. After a few days, the pain is not relieved as much as earlier. What can be the likely reason for this?

a) Decrease in absorption of morphine
b) Pain intensity has increased
c) Opioid receptors are up regulated
d) Desensitization of Opioid receptors

Chronic opioid use leads to tolerance, defined as a decrease of the drug response. It's possible to reproduce in vitro such phenomenon when cellular models expressing OR are exposed to agonists, in that situation, a decrease of signaling is observed and is designated as OR desensitization. Some reports distinguish the OR desensitization from the cellular tolerance.

70. **A female patient presented with pain and redness in great toe. Serum uric acid level is 9.7 mg/dL. The physician prescribed a drug for the treatment. Which of the following enzyme is inhibited by this drug?**

a) Xanthine oxidase
b) Thymidylate synthase
c) phosphoribosyl transferase
d) DHFR

Xanthine oxidase inhibitors are primarily used in the **clinical prevention and treatment of gout associated with hyperuricemia**. The archetypal xanthine oxidase inhibitor, Allopurinol has been shown to have other beneficial effects such as a reduction in vascular reactive oxygen species and mechano-energetic uncoupling.

71. A patient on anticancer therapy developed infection. Total leucocytes count of this patient was 2000/mm3, which of the following drug is effective?

www.medicalcoupons.in

a) Oprelvekin
b) Filgrastim
c) Erythropoietin
d) Romiplostim

Filgrastim is a haematopoietic growth factor that primarily acts to stimulate the proliferation and differentiation of neutrophil progenitor cells. Filgrastim is capable of reducing the incidence and severity of neutropenia and the complications that accompany it in patients with cancer or HIV infection.

72. A person presents with acute chest pain. Nitroglycerine was given sublingually and the pain relieved within 5 minutes. What is the mechanism of the drug?

a) Release of nitric oxide
b) Release of Endothelin
c) Calcium channel Blocker
d) Beta channel Blocker

nitroglycerin converts to nitric oxide (NO) in the body. NO then activates the enzyme guanylyl cyclase, which converts guanosine triphosphate (GTP) to guanosine 3',5'-monophosphate (cGMP) in vascular smooth muscle and other tissues. cGMP then activates many protein kinase-dependent phosphorylations, which enhances the reuptake of calcium into the sarcoplasmic reticulum, increases extracellular calcium, and opens the calcium-gated potassium channel. This ultimately results in the dephosphorylation of myosin light chains within smooth muscle fibers. This activity causes the relaxation of smooth muscle within blood vessels, resulting in the desired vasodilatory effect.

73. A 60-year old man present to OPD with low backache. His PSA levels are 100 ng/ml. Which of the following drug is indicated?

a) Somatostatin
b) Goserelin
c) Terlipressin
d) Testosterone

Goserelin is a parenterally administered, gonadotropin releasing hormone (GnRH) agonist which **causes an inhibition of estrogen and androgen production** and is used predominantly to treat prostate cancer.

74. A patient with resting tremor was diagnosed to be of parkinsonism. Which of the following therapy is not indicated?

www.medicalcoupons.in

a) Pramipexole
b) Rivastigmine
c) Dee Brain Stimulation
d) Levodopa + Carbidopa

Rivastigmine is now widely approved for the treatment of mild to moderately severe dementia in Parkinson's disease (PDD). However, since anticholinergic drugs have a role in the management of tremor in patients with Parkinson's disease (PD), concerns have been raised that the use of cholinergic drugs might worsen PD

75. A Pregnant female presented with burning micturition. A diagnosis of cystitis was made. which of the following drug will be useful for the treatment?
a) Amoxicillin
b) TMP - SMX
c) Ciprofloxacin
d) Doxycycline

Those with asymptomatic bacteriuria can be treated with a single dose of an antimicrobial to which the organism is susceptible. For those with symptomatic UTI, we recommend **amoxicillin 500 mg tid for three days**.

76. A patient has pain in facial nerve distribution while chewing and speaking, which of the following drug should be given for treatment?
a) Lamotrigine
b) Phenobarbitone
c) Haloperidol
d) Carbamazepine

Carbamazepine is a **medicine used to treat epilepsy**. It can also be taken for nerve pain caused by diabetes (peripheral neuropathy) or if you have a painful condition of the face called trigeminal neuralgia.

77. Which of the following is a true statement regarding Aliskiren?
a) It is contraindicated in diabetes mellitus
b) It is indirectly increase renin levels
c) It is given in pregnancy
d) It increases the aldosterone levels

Aliskiren is in a class of medications called **direct renin inhibitors**. It works by decreasing certain natural chemicals that tighten the blood vessels, so blood vessels relax and the heart can pump blood more efficiently

www.medicalcoupons.in

78. Which of the following drug is used as radio sensitizer in head and neck cancers?

a) Cisplatin
b) Amifostine
c) Paclitaxel
d) Doxorubicin

Paclitaxel is a radiosensitizer which **may stabilize microtubules**, block the G2/M phase of the cell cycle and thus modulate the radioresponsiveness of tumor cells

79. A 20-year-male is on low dose of oral steroid therapy for the last 10 years, now he develops progressive loss of vision in both the eyes. What is the reason?

a) Glaucoma
b) Cystoid macular edema
c) Cataract
d) Retinal detachment

Taking steroids can **cause a cataract** type called posterior sub-capsular cataracts. It causes a small, cloudy area to form underneath the eye's lens. While cataracts are a known side effect for some people when taking steroids, they're highly treatable.

80. For the treatment of acute migraine attack, a patient took some medicines. After that he develops change in colour vision and numbness in the tip of the thumb. What is likely drug responsible and its treatment?

a) Sumatriptan
b) Dihydroergotamine
c) Propranolol
d) Phenoxybenzamine

Dihydroergotamine is a severe vaso constrictor and it mimics peripheral vascular disorders when used in acute migraine attack.

81. A child presented to emergency with GTCS since 30 minutes. Airway has been secured but the child is still having seizures. What is initial treatment to control the seizures?

a) Valproate
b) Diazepam
c) Phenobarbitone
d) Levetiracetam

www.medicalcoupons.in

In the absence of intravenous access, **buccal midazolam or rectal diazepam** are therefore acceptable first- line anticonvulsants for the treatment of an acute tonic-clonic convulsion that has lasted at least five minutes.

82. A Person from non-endemic area visited India and was given prophylaxis for malaria. How long the person should continue after travelling from India?

a) 7days
b) 15days
c) 4weeks
d) 4 months

All prophylactic drugs should be taken with unfailing regularity for the duration of the stay in the malaria risk area, and should be continued for **4 weeks after the last possible** exposure to infection since parasites may still emerge from the liver during this period.

83. A child took several iron tablets. What is the antidote for poisoning?

a) BAL
b) EDTA
c) Penicillamine
d) Desferrioxamine

Desferrioxamine injection is used to remove excess iron from the body in anemia or thalassemia patients who have many blood transfusions. It is also used with other medicines to treat acute iron poisoning, especially in small children.

84. Adrenaline acts via beta 2 receptors cause bronchodilatation where histamines act on H1 receptors to cause bronchoconstriction. Which type of antagonism is this?

a) Physical
b) Chemical
c) Pharmacological
d) Physiological

Physiological antagonism describes the behaviour of a substance that produces effects counteracting those of another substance (a result similar to that produced by an antagonist blocking the action of an agonist at the same receptor) using a mechanism that does not involve binding to the same receptor.

www.medicalcoupons.in

85. A Person having a history of angina and heart rate of 90 bpm and tense personality developed BP of 150/125. The preferred drug for treatment of hypertension?

a) Enalapril
b) Amlodipine
c) Atenolol
d) Thiazides

Atenolol belongs to a group of medicines called beta blockers. It's **used to treat high blood pressure and irregular heartbeats** (arrhythmia). It can also be used to prevent chest pain caused by angina. If you have high blood pressure, taking atenolol helps prevent future heart disease, heart attacks and strokes

86. A boy fell down while playing and the nose is deviated but septum was found normal, what is the next step done?

a) Lower the edema then close reduction after 7 days
b) Closed reduction and automatically swelling decreases
c) Septo-rhinoplasty
d) Open reduction

Initially medications given to treat the edema and close reduction done after 7 days.

87. A patient underwent removal of submandibular gland and lingual nerve was damaged during surgery. Which of the following is NOT correct?

a) Rate of sublingual (or submandibular) secretion is reduced.
b) Taste sensation of Anterior 2 / 3rd of tongue is lost.
c) Sensation in floor of mouth lost
d) Tongue deviation.

If your lingual nerve sustains an injury, you'll most likely experience any of these nerve damage symptoms: Changed sensation in the tongue, chin, or lower lip areas (similar to sensations you feel when your oral cavity is numbed for a dental procedure or as the anesthesia slowly wears off) **Altered ability to taste**. but not tongue deviation.


88. Syphilis affects which part of the nose?

a) Bony septum
b) Lateral wall of nose
c) Floor of nose
d) vestibule

www.medicalcoupons.in

Syphilis and leprosy are bacterial infections that can have many health implications, including lesions and ulcers that attack **the cartilage in the nose (bony septum).** If left untreated, these infections could cause permanent damage to the nose, resulting in a saddle nose deformity.

89. A professional trumpet blower presents with left sided neck swelling. Following is the radiographic findings of the patient. What is the most likely diagnosis?



a) Laryngocele.
b) Laryngeal cyst
c) Thyroglossal cyst
d) Epiglottits

Laryngoceles can be congenital or acquired, though the majority fall into the latter category. They occur as a result of **increased intralaryngeal pressure** such as that in excessive coughing, playing blowing instruments or due to obstructing lesions within the larynx.

90. Drug used for treatment of cystitis in pregnancy?
a) Amoxicillin/ Ampicillin
b) Ciprofloxacin
c) Azithromycin
d) Sulfonamides

Those with asymptomatic bacteriuria can be treated with a single dose of an antimicrobial to which the organism is susceptible. For those with symptomatic UTI, we recommend **amoxicillin 500 mg t/d for three days**.


www.medicalcoupons.in

91. Pharyngeal pouch is located between?

a) Superior and middle constrictor
b) Middle and Inferior constrictor
c) **Thyropharyngeus and cricopharyngeus**
d) Thyroid and cricoids

Pharyngeal pouch is located between Thyropharyngeus and cricopharyngeus. A Zenker's diverticulum is an outpouching that occurs **at the junction of the lower part of the throat and the upper portion of the esophagus**. The pouch forms because the muscle that divides the throat from the esophagus, the cricopharyngeal (CP) muscle, fails to relax during swallowing.

92. Endoscopic image is given. Identify the marked structure? (Marked in YELLOW)



a) Eustachian tube opening
b) Torus tubaris
c) Fossa of rosenmuller
d) None

The fossa of Rosenmüller is **a bilateral projection of the nasopharynx just below the skull base**. It is also called the lateral pharyngeal recess or simply the pharyngeal recess. The fossa is covered by nasopharyngeal mucosa and is the most common site of origin of nasopharyngeal carcinoma

93. Hypopharyngeal carcinoma with involvement of the same side with no extension to the opposite side. What is the staging?


a) T3 NO MO
b) T4b NO MO
c) T1 NO MO
d) T4a NO MO

www.medicalcoupons.in

- N0: No regional LN
- N1: Single ipsilateral LN ≤ 3cm
- N2a: Single ipsilateral LN 3-6cm
  - b: Multiple ipsilateral LNs ≤ 6cm
  - c: Bilateral or contralateral LNs ≤ 6cm
- N3: Any LN more than 6cm

- M stage:
- Mx- cannot be assessed,
- M0- no distant metastasis,
- M1- distant metastasis

T & N STAGING IN HYPOPHARYNGEAL CANCERS



T1	Limited to 1 subsite of hypopharynx; Greatest dimension ≤ 2 cm
T2	Involves >1 subsite of hypopharynx, or adjacent site, or measures 2-4cms in greatest dimension without fixation of hemilarynx or extension to oesophagus
T3	> 4cms in greatest dimension or fixation of hemilarynx or extension to oesophagus
T4	Moderately advanced or very advanced
T4a	Moderately advanced local disease Invades thyroid/cricoid cartilage, hyoid bone, thyroid gland, oesophagus, or central compartment soft tissue
T4b	Very advanced local disease Invades prevertebral fascia, encases carotid artery, or involves mediastinal structures

94. Reservoir sign and pain near mastoid antrum is seen in which of the following condition?

a) Mastoiditis

b) Acute otitis media

c) otitis externa

d) CSOM

Mastoiditis is the result of an infection that extends to the air cells of the skull behind the ear. Specifically, it is an inflammation of the mucosal lining of the mastoid antrum and mastoid air cell system inside the mastoid process

www.medicalcoupons.in

95. All are true about the given picture. Except?



a) Its usually a retention cyst

b) Extravasation is not that common

c) It is transluminant

d) All are true

Extravasation is not common and its transluminant from the given picture. but it is not usually a retention test

96. A 2-month old baby is having watery discharge / fluid leakage from umbilicus, this is due to

a) Failure of midgut to recanalize

b) Failure of involution of allantoic duct

c) Failure of involution of omphalomesenteric duct

d) Failure of rotation of midgut

Patent urachus refers to a spectrum of umbilical disorders that result from the failure of involution of fetal embryologic tissues that connect the developing bladder to the umbilical cord. Similarly, presentation is variable and can be readily evident at birth with umbilical cord abnormalities or diagnosed later in adolescence with the complaint of persistent umbilical wetness or recurrent umbilical infection.

97. A paediatric patient presented with headache, ear discharge, fever, resp[illegible] sign is present. what is the diagnosis?

a) CSF otorrhea

b) Acute mastoiditis

c) Acute suppurative otitis media

d) Otitis externa

www.medicalcoupons.in

Mastoiditis is the result of an infection that extends to the air cells of the skull behind the ear. Specifically, it is an inflammation of the mucosal lining of the mastoid antrum and mastoid air cell system inside the mastoid process

98. A mother delivered 33 weeks old baby, via normal vaginal delivery with no complications at birth. What is the preferred method of feeding the neonate?

a) OGT feeds
b) Direct breast feeding
c) Paladai feeds
d) Parenteral nutrition

The Paladai is a cup-like utensil with a narrow tip that has been used traditionally to feed babies in India1 when the **mother cannot breast** feed. It is used to hold expressed breast milk or other types of liquid or semi-solids. The baby is held reclining while being fed.

99. A 6-year-old child presented with abdominal pain, hematemesis and palpable spleen. He had a history of exchange transfusion for neonatal jaundice. What is the probable diagnosis?

a) Budd Chiari syndrome
b) Cirrhosis of liver
c) Viral hepatitis
d) Portal vein thrombosis

Portal vein thrombosis is **blockage or narrowing of the portal vein** (the blood vessel that brings blood to the liver from the intestines) by a blood clot. Most people have no symptoms, but in some people, fluid accumulates in the abdomen, the spleen enlarges, and/or severe bleeding occurs in the esophagus.

100. A 1-year-old child presents with 1st episode of wheeze and breathing difficulty. Chest X-ray shows bilateral hyperinflation. Which of the following can be a causative organism for this case?

a) RSV
b) S. pneumoniae
c) H. influenza
d) Coxsackie virus

The early phase of RSV in babies and young children is often **mild**, like a cold. In children younger than age 3, the illness may move into the lungs and cause

www.medicalcoupons.in

coughing and wheezing. In some children, the infection turns to a severe respiratory disease.

101. True regarding buffalo milk composition when compared to human milk?

a) High calorie, high protein
b) Low calorie, low protein
c) High calorie, low protein
d) Low calorie, high fat

Buffalo milk is **rich in vitamin A**, has a higher protein value and contains more iron, calcium and phosphorus than the cow's milk. ... It is better to introduce buffalo milk slowly.

102. A 3-month old child presents with complaints of fever, cough, not able to feed and breathing difficulty. On examination, respiratory rate was 58 / min, chest retractions were present. What would be your next step of Management?

a) Severe pneumonia - Give antibiotics and refer immediately
b) Very severe pneumonia and refer urgently
c) Pneumonia -Start antibiotics and send child home
d) No pneumonia

Beta-lactam antibiotics (e.g., **amoxicillin, cefuroxime**) are preferred for outpatient management. And is referred immediately to regional hospital

103. All of the following are components of TOF except?

a) Overriding aorta
b) Sub-pulmonic stenosis
c) VSD
d) Tricuspid stenosis

The four components that make up the "tetralogy" include:

- a ventricular septal defect (VSD);
- pulmonary stenosis (sub-valvar, valvar and/or supra-valvar);
- an overriding aorta; and.
- right ventricular hypertrophy.

www.medicalcoupons.in

104. A neonate was found to have Grimace on inserting a catheter into nose, some flexion of limbs was noted. Heart rate - 120 / minute, Blue extremities were noted with irregular breathing. Find out the APGAR Score?

a) 4
b) 6
c) 8
d) 10



105. A patient's mother while bathing her child noticed there was no testes in the scrotum and on examination- one right testes was in Inguinal region and left was in perineum. What is the DIAGNOSIS?

a) bilateral undescended testes
b) bilateral ectopic testes
c) right undescended and left ectopic testes
d) left undescended and right ectopic testes

from the examination above we can easily deduct that the right testes are undescended and left testes had become ectopic.

106. A 4-year-old child presented with fever, running nose and following lesions. Probable aetiology?

a) HSV 1
b) HSV 2
c) Coxsackie virus
d) Streptococci pneumonia

www.medicalcoupons.in

Coxsackievirus can produce a wide variety of symptoms. About half of all kids with an infection have no symptoms. Others suddenly get a high fever, headache, and muscle aches, and some also develop a sore throat, abdominal discomfort, or nausea. a type of coxsackievirus syndrome, causes painful red blisters in the throat and on the tongue, gums, hard palate, inside of the cheeks, and the palms of hands and soles of the feet.

107. A patient of 23 years came to the clinic with a known history of myopia with -2 D in left and -3 D in the right eye. He is sent for retinoscopy. What is the movement of retinal reflex in the right eye?

a) It moves in the same direction of the retinoscope


b) It moves in the opposite direction of the retinoscope

c) Vertical movement is having wider reflection and moves in the opposite direction

d) There is no movement of the reflex

it moves in the opposite direction because of the myopia negative i.e. -2D and -3D respectively

108. Identify the clinical condition from the skin finding as given below:



a) Chicken pox

b) HSV

c) HPV

d) Measles

Chickenpox is a very contagious infection caused by the varicella-zoster virus. It mainly affects kids, but adults can get it, too. The tell-tale sign of chickenpox is a super-itchy skin rash with red blisters. Over the course of several days, the blisters pop and start to leak. Then they crust and scab over before finally healing.

www.medicalcoupons.in

109. Which curvature develops after the baby starts to walk?

a) Primary cervical curvature
b) Secondary sacral curvature
c) Primary sacral curvature
d) Secondary lumbar curvature

As babies, children have a C-shaped spine. Secondary curves in the cervical and lumbar spine develop as **infants become able to lift their heads** sit up, crawl, stand, and walk. As children grow, their spine continues to develop natural curves into a normal, mature spine.

110. A mother complains that her baby is not sucking properly from breast and now complaints of pain and swelling of the breast. Probable condition?

a) Breast abscess
b) Mastitis
c) Breast engorgement
d) Sore nipple

Breast engorgement means **your breasts are painfully overfull of milk**. This usually occurs when a mother makes more milk than her baby uses.
Once engorgement occurs **gentle massage**, frequent feeding, correct positioning and warm compresses to the breast have been advocated to relieve symptoms along with analgesia to relieve pain.

Other Causes:
Sore and cracked Nipples
Inverted Nipples
Engorged breasts
Mastitis
Previous surgeries on breast
Neonatal causes
Insufficient/Unsuitable Milk

111. In a paediatric patient with seizure which drug is initially used in management?

a) Diazepam
b) Phenytoin
c) Phenobarbitone
d) Haloperidol

www.medicalcoupons.in

In the absence of intravenous access, **buccal midazolam or rectal diazepam** are therefore acceptable first- line anticonvulsants for the treatment of an acute tonic-clonic convulsion that has lasted at least five minutes.

112. A 54-year-old female patient with the history of Amenorrhea for last 16 months. What should be the hormone

levels identified in this patient?

a) Increased LH, decreased FSH

b) Increased LH and FSH

c) High FSH, low LH

d) LOW LH AND FSH

- Menopause
- Permanent cessation of menses dated 12 months of amenorrhea
- **The avg. age - 51 years (range of 43-57 years)**
- Perimenopause
- 2-8 years preceding + 1 year after last menses - fluctuating hormone levels irregular menstrual cycle & symptoms

When estrogen deficiency occurs in the menopause LH levels increase. Later the FSH is raised and remains high for the rest of life. This raised FSH and low estrogen levels appear to cause the characteristic **hot flushes.**

113. In which of the following condition, MTP is not indicated?

a) Medical: mother's life is in danger

b) Socioeconomic: large family size

c) Eugenic: coming baby is handicap

d) Failure of contraception

MTP Act INDICATIONS:

- Continuation of pregnancy constitutes risk to the life or grave injury to the physical or mental health of woman
- Substantial risk of physical or mental abnormalities in the fetus as to render it seriously handicapped
- Pregnancy caused by rape (presumed grave injury to mental health)
- Contraceptive failure in married couple (presumed grave injury to mental health)
- "In determining whether the continuance of pregnancy would involve such risk of injury to the health (as mentioned above), account may be taken of the pregnant woman's actual or reasonable foreseeable environment"

www.medicalcoupons.in

www.medicalcoupons.in

114. A 40 years old female with severe cardiac compromise, she had to undergo an operation for which the surgeon gave clearance, which of the following surgery must not be done?

a) Total hysterectomy
b) Subtotal hysterectomy
c) Vaginal hysterectomy
d) Laparoscopic hysterectomy

Positive pressure pneumoperitoneum with CO2 has deleterious effects on CVS so Laparoscopic Hysterectomy is not done in this patient

Positive-pressure capnoperitoneum (PPCpn) is **the conventional method of exposure in laparoscopic surgery**. This positive intra-abdominal pressure (10–12 mm Hg) is associated with a variable but significant reduction in cardiac output that may be detrimental in high-risk patients with poor cardiac reserve.
Pneumoperitoneum in excess of 15mmHg has deleterious effects on the cardiovascular system. The pneumoperitoneum compresses the vena cava and thus **decreases venous return to the heart**; this results in blood pooling in the lower half of the body and a decrease in cardiac output.

115. What is the first sign of $MgSO_4$ toxicity?

a) Respiratory depression
b) Loss of patellar reflex
c) Cardiac arrest
d) Absent urine output

Serum Mg2+levels(mEq/l)	Clinical effects
4-7	Normal therapeutic level
8-10	Patellar reflex disappear
10-15	Respiratory depression
12-15	Respiratory paralysis
25 - 30	Cardiac arrest

Calcium gluconate is the antidote for Magnesium Sulfate toxicity.

116. A mother brings her 18-year-old child with complaints of not attained menarche. GPE shows normally developed breast and pubic hair. On

www.medicalcoupons.in

examination vaginal ending is blind and uterus not palpable. What is the next line of investigation?


a) USG
b) T3, TSH
c) Estrogen and Progesterone
d) FSH and LH

USG would be most helpful to know presence or absence of uterus.
Karyotype: To differentiate between MRKH and Androgen insensitivity.
Mayer-Rokitansky-Küster-Hauser (MRKH) syndrome is a **rare disorder that affects women**. It is characterized by the failure of the uterus and the vagina to develop properly in women who have normal ovarian function and normal external genitalia.

117. A Female patient came with c/o pruritis, greenish discharge per vaginum. What is the causative organism?

a) Trichomonas
b) Candida - curdy white discharge + itching
c) Bacterial vaginosis - Foul smelling greyish white + no itching
d) Chlamydia - mucopurulent endocervical discharge

TRICHOMONAS VULVOVAGINITIS :



- M/c vaginitis
- Etiology - **Trichomonas vaginalis** **flagellated** protozoan
- Mode of Spread: Sexual & non Sexual
- Symptoms - Nature of discharge: **Profuse frothy**, Colour of discharge: **Yellowish green, Pruritis: Present**
- Signs - Discharge: **serous & frothy & greenish**, pH: > 5, Vulva: inflamed, Vagina: **Strawberry Vagina**
- Dx: **Wet mount: motile flagellated trichomonas**, Clue cells: absent, Whiff test: negative
- Definitive Dx: culture on **Feinberg Whittington media** or **Diamonds TYM media**
- Rx - **Metronidazole 400mg TID** x 5days 'or' Metronidazole 2gm single dose (Mode : both partners)

www.medicalcoupons.in

118. A 14-year-old girl with well-developed secondary sexual characteristics but presents with amenorrhea. What is the treatment you will prefer next?

a) Wait and Watch
b) Estrogen
c) Give ADH
d) GnRH

- Delayed puberty is said to occur when there is no breast development by the age of 13 and no menarche by age of 15 years, so the best answer to this question is to wait and watch.

119. 33-year-old unmarried female presents with the complaint of amenorrhea and fainting. what is the next step you should perform in this patient?

a) UPT (urine pregnancy test)
b) Endoscopic Biopsy
c) D and C
d) Hysteroscopy

Always rule out Pregnancy first in case of secondary amenorrhea.

Amenorrhea can occur for a variety of reasons. Some are normal, while others may be a side effect of medication or a sign of a medical problem.

Causes of 2^0 Amenorrhea


- Pregnancy
- Breastfeeding
- Menopause
- Contraceptives
- Medications
- Lifestyle factors
- Hormonal imbalance
- Structural problems

120. A female in labor is having occipital posterior position. The cervix gets 8cm dilated, forceps applied in occipitomandibular direction and tried to rotate, but doctor is not able to rotate. In which type of pelvis, the fetal head cannot be rotated?

a) Android pelvis
b) Anthropoid pelvis
c) Platypelloid pelvis
d) Gynecoid pelvis

www.medicalcoupons.in

Contracted pelvis: rotation of the head cannot easily occur in android pelvis due to projection of the ischial spines and convergence of the side walls.
Anthropoid: The elongated shape of the anthropoid pelvis makes it roomier from front to back than the android pelvis. But it's still narrower than the gynecoid pelvis. Some pregnant women with this pelvis type may be able to have a vaginal birth, but their labor might last longer.



IMPORTANT POINTS

- M/c pelvis – Gynecoid*
- Least common pelvis - Platypelloid pelvis*
- The only pelvis with AP > Tr – Anthropoid pelvis*
- Face to pubis is M/c in – Anthropoid pelvis*
- OP position is M/c in – Anthropoid pelvis*
- DTA is M/c in – Android pelvis*
- Flat bowl shaped pelvis is – Platypelloid pelvis*
- Marked posterior asynclitism is seen in – Platypelloid pelvis*

121. Female having 14 weeks of missed abortion. Which is the agent of choice to be used in this condition?

a) Mifepristone
b) Misoprostol
c) Oxytocin
d) Methotrexate

Misoprostol is a non-invasive, effective medical method for completion of abortion in missed abortion.

➢ **Missed Abortion**

- When the fetus **is dead and retained inside the uterus** for a variable period, it is known as *missed abortion*


✓ C/o - **no bleed**, no morning sickness and regression of breast changes

www.medicalcoupons.in

- ✓ O/E - Uterus – **lesser than gestational age** (arrested at the stage when fetus was lost) Cx - Internal **os is closed**
- ✓ USG – **dead fetus**

122. A 35yr old Female presented with 14 week of gestation with bleeding per vaginum, doctor started misoprostol with dose 200 mcg. How much maximum dose can be given?

a) 3
b) 4
c) 5
d) 6



123. Cervical cancer is caused by which type of virus?

a) HPV
b) HIV
c) HBV
d) EBV

Long-lasting infection with certain types of human papillomavirus (HPV) is the main cause of cervical cancer. HPV is a common virus that is passed from one person to another during sexual intercourse.

HUMAN PAPILLOMA VIRUS

- The M/c etiology - epitheliotropic, oncogenic virus
- **9 major genes - 7 (E1 to E7) + 2 (L1, L2)**
- L1(major capsid) & L2(minor capsid protein)
- Viral life cycle – enters genital tract → attaches to the cells → enters basal cells (endocytosis) →viral DNA injected into human genome→Expression
- Oncogenes - E6 (inhibits p53 anti-oncogene) & E7 (inhibits Rb anti-oncogene)

www.medicalcoupons.in

- High risk serotypes **- 16,18,31,33,35, 39,45, 52, 56, 58,59, 68 (strongly associated œ16, 18)**
- **Low risk - genital warts œ6 & 11.**
- Immune handling - 80% women at some point infected - infection clears in **9 œ15 months** - by body's immunity
- PCR / Southern blot – detects HPV DNA – used as screening along with pap smear


124. What is shown in the following image?



a) Polyhydraminos
b) Endometriosis
c) Endometritis
d) Mucinous ovarian adenoma

Ovarian mucinous cystadenoma is **a benign tumour that arises from the surface epithelium of the ovary**. It is a multilocular cyst with smooth outer and inner surfaces. It tends to be huge in size.






www.medicalcoupons.in

125. A Young Female is having inserted IUCD 6 months ago. Now she has presented with the complaints of amenorrhea and fainting since two cycles. What may be the cause for this condition?

a) PID
b) Endometriosis
c) Ectopic pregnancy
d) Failed contraception

The main risk factor for contraceptive-failure ectopic pregnancy is IUD failure. **IUDs do not increase the absolute risk of ectopic pregnancy**, but a pregnancy occurring with an IUD is more likely to be ectopic than intrauterine.

126. Identify the given non spore forming bacteria?



a) Clostridium difficile
b) Clostridium perfringens
c) Clostridium tetani
d) Corynebacterium diphtheria

Corynebacterium diphtheriae is the leading causing agent of diphtheria. It is a non-motile, non-encapsulated, non-sporulating gram-positive rod-shaped bacterium with a high GC-content and occurs in four biovars: gravis, mitis, intermedius, and belfanti, based on colonial morphology and biochemical profiles.

127. A primigravida is in labor from last 12 hours with tender and rigid uterus. Fetal head not palpable, loss of Fetal movements. what is the most appropriate diagnosis?

www.medicalcoupons.in

a) Abruptio placenta

b) placenta previa

c) polyhydramnios

d) Prom

Placental abruption is a complication of pregnancy that happens when the placenta separates from your uterus before your baby is born. Symptoms or signs can also include:

- Abdominal pain.
- Uterine contractions that are longer and more intense than average labor contractions.
- Uterine tenderness.
- Backache or back pain.
- Decreased fetal movement.

128. A female with C-Section delivery the baby was seeming to be normal but the placenta was morbidly adherent, what could be the reason for this condition?

a) absence of nitabuchs membrane

b) absence of decidua parietalis

c) absence of syncytiotrophoblast

d) absence of cytotrophoblast

Morbidly Adherent Placenta

- A morbidly adherent placenta doesn't separate from the uterine wall upon attempted manual removal
- It is a condition wherein the placenta is firmly adherent to the uterine wall and doesn't separate during the third stage
- ✓ Pathological Findings
- Absence of decidua basalis or improper decidua basalis formation
- Absence of Nitabuch's fibrinoid layer
- The trophoblasts completely invade the endometrium and have the potential to invade the myometrium

129. A 37 weeks of pregnant female came with the complaint of bleeding per vaginum associated with pain. the fetal heart rate was absent and uterus is tender. what is the diagnosis and management for this patient?

a) Placental abruption and normal vaginal delivery

b) Placental abruption and LSCS

c) Placenta previa and normal vaginal delivery

d) Placenta previa and LSCS

www.medicalcoupons.in

bleeding per vagina with pain indicates placental abruption and since the fetal heart rate was absent LSCS would be the best management.

130. Hormone responsible for Lactogenesis?

a) Oxytocin

b) GH

c) Estrogen

d) Progesterone

The two primary hormones that are needed for lactation are **prolactin and oxytocin**. Prolactin stimulates milk biosynthesis within the alveolar cells of the breast and oxytocin stimulates contraction of the myoepithelial cells that surround the alveoli, causing the milk to be ejected into the ducts leading to the nipple.

131. A pregnant lady delivered a child and after 24 hrs later she came with complaint of heavy bleeding. What might be the cause for the bleeding in this patient?

a) Battledore placenta

b) Succenturiate lobe

c) Velamentous placenta

d) Circumvallate placenta

A **succenturiate lobe** is a variation in placental morphology and refers to a smaller accessory placental lobe that is separate to the main disc of the placenta. There can be more than one succenturiate lobe.

Complications

- increased incidence of type II vasa previa
- increased incidence of postpartum hemorrhage due to retained placental tissue



www.medicalcoupons.in

132. Most common symptom seen in endometriosis is;

a) Polymenorrhea
b) Dysmenorrhea
c) HMB
d) Amenorrhea

- Endometriosis Symptomatic - Triad of **Dysmenorrhea, Infertility & Dyspareunia**

- **Dysmenorrhea - M/c symptom**, secondary dysmenorrhoeal, progressive dysmenorrhea, **deep-seated pelvic pain**, Associated with premenstrual and postmenstrual spotting, little correlation /w extent of lesion, correlates well with the **depth of the lesion**
- Dyspareunia – **POD** & rectovaginal septum involvement
- **Infertility - 30-40%** patients with endometriosis have infertility, **15-30%** of infertile will have endometriosis

133. A 32-year-old female patient complaints on her 4th day of postpartum delivery of fever, dysuria, foul smelling discharge. what is the probable diagnosis in this patient?

a) Puerperal sepsis
b) Thrombhophlebitis
c) UTI
d) Endometritis

Puerperal sepsis was defined as infection of the genital tract occurring at any time between the onset of rupture of membranes or labour, and the 42nd day postpartum in which two or more of the following are present:

- Fever (oral temperature 38.5°C/101.3°F or higher on any occasion).
- Pelvic pain.
- Abnormal vaginal discharge, e.g. presence of pus.
- Abnormal smell/foul odor of discharge.
- Delay in the rate of reduction of the size of the uterus (involution).

134. Auspitz sign is seen in?

a) Pemphigus vulgaris
b) Psoriasis Vulgaris
c) Pustular psoriasis
d) DLE

www.medicalcoupons.in

The Auspitz sign refers to the **appearance of small bleeding points after successive layers of scale have been removed from the surface of** psoriatic papules or plaques

135. **Ulcer "with overhanging / undermined edges" in the neck region as shown in the image given below.**
Diagnosis?



a) Tubercular lymph node
b) Malignant Cervical lymphadenopathy
c) Syphilis
d) Reactive lymphadenopathy

Scrofula refers to **painless swelling of the cervical and supraclavicular lymph nodes** that is most often caused by M. tuberculosis in adults and nontuberculous mycobacteria in children.

136. Which mineral deficiency can lead to following condition?



www.medicalcoupons.in

a) Zinc
b) Iron
c) Calcium
d) Vitamin A

Zinc deficiency occurs in **infants when its demand exceeds its supply**. It presents with cutaneous signs which, in severe cases, are associated with diarrhea, alopecia, and irritability. Genetic and acquired forms of zinc deficiency have been described and often have overlapping clinical features.

137. First line treatment for acne comedones?




a) Topical steroids
b) Topical antibiotics
c) Benzyl peroxide
d) Topical retinoids

Topical retinoids such as tretinoin or adapalene are effective in many patients with comedonal acne. Patients with inflammatory lesions benefit from treatment with benzoyl peroxide, azelaic acid or topical antibiotics. Frequently, the use of comedonal and antibacterial agents is required

www.medicalcoupons.in

138. Identify the condition shown in the image?



a) Malignant melanoma
b) BCC
c) Squamous cell carcinoma
d) Nevus

Melanoma, also redundantly known as malignant melanoma, is **a type of skin cancer that develops from** the pigment-producing cells known as melanocytes. Melanomas typically occur in the skin, but may rarely occur in the mouth, intestines, or eye (uveal melanoma).

139. Which organism is responsible for causing erysipelas?



a) Streptococcus
b) Staphylococcus
c) Bacteroides
d) Fungus

Erysipelas is an infection of the upper layers of the skin (superficial). The most common cause is **group A streptococcal bacteria**, especially Streptococcus

www.medicalcoupons.in

pyogenes. Erysipelas results in a fiery red rash with raised edges that can easily be distinguished from the skin around it.

140. Cutaneous horn as shown in the image is associated with which of the following skin cancer?



a) Squamous cell carcinoma
b) Melanoma
c) BCC
d) Actinic keratosis

Squamous cell carcinoma (SCC) is **the second most common form of skin cancer**. It's usually found on areas of the body damaged by UV rays from the sun or tanning beds. Sun-exposed skin includes the head, neck, chest, upper back, ears, lips, arms, legs, and hands. SCC is a fairly slow-growing skin cancer.

141. Identify the condition shown in the image?



www.medicalcoupons.in

a) Condyloma acuminata
b) Bowen disease
c) Condyloma lata
d) Hemorrhoids

Condylomata acuminata (CA; singular: condyloma acuminatum), also known as **anogenital warts**, are manifestations of HPV infection that occur in a subset of individuals with anogenital HPV infection

142. Patient suffering from ulcerative colitis presents with ulcer on anterior leg as shown in the image. Diagnosis?



a) Pyoderma gangrenosum
b) Venous ulcer
c) Tubercular ulcer
d) Chancroid

Pyoderma gangrenosum is a rare condition that **causes large, painful sores (ulcers) to develop on your skin**, most often on your legs. The exact causes of pyoderma gangrenosum are unknown, but it appears to be a disorder of the immune system.

143. Common complication associated with the following condition?



www.medicalcoupons.in

a) Post streptococcal glomerulonephritis
b) SSSS
c) Encephalitis
d) Pneumonitis

PSGN is a kidney disease that can develop after infections caused by bacteria called group A Streptococcus (group A strep). These infections include throat and skin infections like strep throat, scarlet fever, and impetigo. PSGN is not a group A strep infection of the kidneys.

144. A tourist guide in Mt. Everest, after one of his trips developed blisters on the feet.
Which is not useful for his treatment?



a) Hyperbaric Oxygen
b) Aspirin
c) Phenylephrine
d) Pentoxifylline

Phenylephrine is **used to relieve nasal discomfort caused by colds, allergies**, and hay fever. It is also used to relieve sinus congestion and pressure. Phenylephrine will relieve symptoms but will not treat the cause of the symptoms or speed recovery. So it cannot be helpful for the patient with the above condition.

145. Patient presents with history of severe sunburn after only a few minutes in the sun (photo sensitivity), freckling in sun exposed areas, dry skin and changes in skin pigmentation. Diagnosis?



www.medicalcoupons.in

a) Xeroderma pigmentosum
b) Bloom syndrome
c) Warts
d) Melanocytes nevus

Xeroderma pigmentosum (XP) is a **very rare skin disorder** where a person is highly sensitive to sunlight, has premature skin ageing and is prone to developing skin cancers. Xeroderma pigmentosum is caused by cellular hypersensitivity to ultraviolet (UV) radiation, as a result of a defect in the DNA repair system.

146. Child presents with sores in the mouth and a rash on the hands and feet as shown in the image below. What is the causative organism for it?



a) Coxsackievirus A16
b) Pox virus
c) Herpes virus
d) Human papilloma virus

Coxsackievirus can produce a wide variety of symptoms. About half of all kids with an infection have no symptoms. Others suddenly get a high fever, headache, and muscle aches, and some also develop a sore throat, abdominal discomfort, or

www.medicalcoupons.in

nausea. a type of coxsackievirus syndrome, causes painful red blisters in the throat and on the tongue, gums, hard palate, inside of the cheeks, and the palms of hands and soles of the feet.

147. Child presents with fever and pleomorphic rash as shown in the image. Diagnosis?



a) Chicken pox
b) Small pox
c) Measles
d) Rubella

Chickenpox is a very contagious infection caused by the varicella-zoster virus. It mainly affects kids, but adults can get it, too. The tell-tale sign of chickenpox is a super-itchy skin rash with red blisters. Over the course of several days, the blisters pop and start to leak. Then they crust and scab over before finally healing.

148. A Normal individual underwent an evaluation of his cardiac tissue. Which isoform of lactate dehydrogenase is predominant in the cardiac tissue?
a) LDH1
b) LDH2
c) LDH3
d) LDH4

The LDH-1 isoenzyme is found predominately in **cardiac muscle**, LDH-2 is found primarily in the reticuloendothelial system, LDH-3 predominates in the lungs, LDH-4 in the kidneys, and LDH-5 in the liver and skeletal muscle.

149. A patient has been diagnosed with a solid cancer of the bowel. He also experienced massive proteinuria after a few months of initial diagnosis of cancer. which of the following is a likely cause for the development of the urinary finding?

www.medicalcoupons.in

a) Focal segmental glomerulosclerosis
b) Minimal change disease
c) Membranous Glomerulopathy
d) MPGN

Membranous Nephropathy (MN) is a **kidney disease** that affects the filters (glomeruli) of the kidney and can cause protein in the urine, as well as decreased kidney function and swelling. It can sometimes be called membranous glomerulopathy as well.

150. Which of the following contributes to the immortality of cancer cells?
a) Topoisomerase
b) Telomerase
c) RNA polymerase
d) Helicase

With each cell division, telomeres shorten until eventually they become too short to protect the chromosomes and the cell dies. Cancers become immortal **by reversing the normal telomere shortening process** and instead lengthen their telomeres

151. A patient died of myocardial infarction and his heart was taken out and immersed in a solution. The normal area of the heart turned brick red whereas infarcted portion turned white. Which of the following can be likely component of the solution?
a) Formalin
b) Triphenyl tetrazolium chloride
c) Ethanol
d) Glutaraldehyde

Triphenyl tetrazolium chloride (TTC) is one of the most conventional stains to detect **infarcted** area of the heart in animal experiments. ... Samples with old myocardial infarction produced clear TTC contrast; normal tissue is vivid red, and fibrotic myocardium is white discoloration.

152. A patient presents with the presence of a thyroid swelling and weight loss the physician found presence of cervical lymphadenopathy in the same patient and decided to go in for a Thyroid biopsy. Histopathology report of the patient suggested the presence of cells with orphan Annie eye nuclei. Which of the following is a likely diagnosis for this patient?
a) Follicular thyroid cancer
b) Medullary thyroid cancer

www.medicalcoupons.in

c) Papillary thyroid cancer

d) Anaplastic thyroid cancer

The Orphan Annie-eyed clear nucleus, defined as a large, optically clear nucleus, devoid of chromatin strands, with sharp chromatin rim, is a more specific feature than are nuclear grooves or intranuclear cytoplasmic inclusions in papillary thyroid carcinoma.

153. A patient presented with development of seizures for which the CT guided biopsy was done. The histopathology reported presence of psammoma bodies. Which of the following is the most likely diagnosis of this patient?

a) Meningioma

b) Astrocytoma

c) Ependymoma

d) Medulloblastoma

Psammoma bodies (PBs) are **concentric laminated calcified structures**, observed most commonly in papillary thyroid carcinoma (PTC), meningioma, and papillary serous cystadenocarcinoma of ovary but have rarely been reported in other neoplasms and non-neoplastic lesions

154. Which of the following is implicated in the development of angiosarcoma of the liver?

a) Copper


b) Zinc

c) Arsenic

d) Iron

Hepatic angiosarcoma (HAS) is a **rare type of liver cancer** that is often fatal, and arsenic and vinyl chloride monomer (VCM) are two major causal agents.

155. An old man underwent a bone marrow examination which revealed the presence of the finding shown in the image. What is the likely diagnosis of the affected individual?



www.medicalcoupons.in

a) Multiple myeloma
b) ALL
c) Tuberculosis
d) Sickle cell disease

People with smoldering myeloma have some signs of multiple myeloma, such as any of the following: **Plasma cells in the bone marrow between 10% and 60% High level of monoclonal immunoglobulin** (M protein) in the blood. High level of light chains in the urine (also called Bence Jones protein)

156. A 4-year-old child presents with development of fever, petechial spots and complaint of fatigue. He is also having presence of pallor, hepato-splenomegaly as well as sternal tenderness. The clinical situation described above is most correctly associated with which of the following?

a) AML
b) ALL
c) CLL
d) CML

Acute lymphoblastic leukaemia is the most common childhood cancer. It occurs when a bone marrow cell develops errors in its DNA.

Symptoms may include enlarged lymph nodes, bruising, fever, bone pain, bleeding from the gums and frequent infections.

157. A boy presents in the emergency because of development of allergy due to pollen inhalation. Which of the following cells is important in the pathogenesis of this condition?

a) NK cell
b) Neutrophils
c) Helper T cell
d) Cytotoxic T cell

Allergic reactions arise from an inappropriate immune response to otherwise innocuous substances such as pollen, mold, or peanuts, initiated by a group of immune cells known as the TH2 subset, or type 2 "helper" T cells.

158. **Which of the following is the molecule required for presentation of the antigens by the antigen presenting cell to the TH2 cell?**

a) Toll like receptor
b) MHC
c) G protein coupled receptor

www.medicalcoupons.in

d) Lectin receptor

An antigen-presenting cell (APC) or accessory cell is **a cell that displays antigen bound by major histocompatibility complex (MHC) proteins on its surface**; this process is known as antigen presentation. ... APCs process antigens and present them to T-cells.

159. Which of the following is the mode of inheritance in the patient suffering from color blindness?

a) Autosomal dominant
b) Autosomal recessive
c) X linked dominant
d) X linked recessive

Examples of X-linked recessive conditions include **red-green color blindness** and hemophilia A: Red-green color blindness. Red-green color blindness means that a person cannot distinguish shades of red and green (usually blue-green), but their ability to see is normal.


160. A patient has been diagnosed with a condition due to a DNA repair gene defect. He is having hyper pigmentation on the skin and high risk for development of skin cancers as per the physician. which of the following is a likely enzyme defect in the patient?

a) Helicase
b) Nucleotide excision repair defect
c) ATM gene
d) MSH gene

Nucleotide excision repair (NER) is the main pathway used by mammals to **remove bulky DNA lesions** such as those formed by UV light, environmental mutagens, and some cancer chemotherapeutic adducts from DNA. Deficiencies in NER are associated with the extremely skin cancer-prone inherited disorder xeroderma pigmentosum.

www.medicalcoupons.in

161. Which of the following is a likely diagnosis of the patient looking at the karyotype given in the image as Shown



a) Down syndrome
b) Turner syndrome
c) Klinefelter syndrome
d) Edward syndrome

Down syndrome is caused by **trisomy 21** — the person has three copies of chromosome 21, instead of the usual two copies, in all cells. This is caused by abnormal cell division during the development of the sperm cell or the egg cell.

162. Which of the following amino acids is required for the formation of nitric oxide in the blood vessels?

a) Citrulline
b) Arginine
c) Histidine
d) Tryptophan

In the body, the amino acid arginine changes into nitric oxide (NO). Nitric oxide is a powerful neurotransmitter that helps blood vessels relax and also **improves circulation**. Some evidence shows that arginine may help improve blood flow in the arteries of the heart.

163. Macrophages play an important role in phagocytosis and chronic inflammation. Which of the following cells perform the same function in the liver?

a) Merkel cell
b) Sinusoidal cell

www.medicalcoupons.in

c) Kupffer cell

d) Hepatocytes

Hepatic macrophages play a central role in maintaining homeostasis in the liver, as well as in the initiation and progression of liver diseases. Hepatic macrophages are mainly derived from resident hepatic macrophages called **Kupffer** cells or circulating bone marrow-derived monocytes.

164. Which of the following is a characteristic feature associated with Barrett's oesophagus?

a) Squamous metaplasia


b) Intestinal metaplasia

c) Columnar metaplasia

d) Severe dysplasia

Barrett's esophagus is a condition in which there is an abnormal (metaplastic) change in the mucosal cells lining the lower portion of the esophagus, from normal stratified squamous epithelium to simple columnar epithelium with interspersed goblet cells that are normally present only in the small intestine, and large intestine. This change is considered to be a premalignant condition because it is associated with a high incidence of further transition to esophageal adenocarcinoma, an often-deadly cancer.

165. What is the suggestive diagnosis as per given image?



a) Preseptal Cellulitis

b) Impetigo

c) Orbital cellulitis

d) Conjunctivitis


Orbital cellulitis is an infection of the soft tissues of the eye socket behind the orbital septum, a thin tissue which divides the eyelid from the eye socket. Infection isolated

www.medicalcoupons.in

www.medicalcoupons.in

anterior to the orbital septum is considered to be Preseptal cellulitis. Orbital cellulitis most commonly refers to an acute spread of infection into the eye socket from either extension from periorbital structures (most commonly the adjacent ethmoid or frontal sinuses (90%), skin, dacryocystitis, dental infection, intracranial infection), exogenous causes (trauma, foreign bodies, post-surgical), intraorbital infection (endopthalmitis, dacryoadenitis), or from spread through the blood (bacteremia with septic emboli).

166. The image indicates?



a) Simple Myopia
b) Irregular astigmatism
c) Regular astigmatism
d) Simple hyperastigmatism

Each meridian in the regular astigmatic eye has a **uniform curvature** at every point across the entrance of the pupil. This is the most common type of astigmatism in which the symptoms included are blurry vision, headaches, and light sensitivity

167. A 3-year-old child was brought to the ER with unilateral diagnosed as dacryocystitis. What is the first line of management in the given child
a) Massage
b) Syringing and probing
c) Dacryocystorhinostomy
d) Dacryocystoplasty

www.medicalcoupons.in

The simple method of hydrostatic sac massage has been found to be effective in the treatment of dacryocystitis in children. The massage is begun by pressure at the medial canthus, which is effective in blocking the common canaliculus thus preventing reflux backward through the canaliculi.

168. A 60-year-old patient presented with right sided homonymous hemianopia with Macular sparing. Localise the neurological site of the lesion in this condition.

a) Medial geniculate body
b) Optic Tract
c) Optic Chiasma
d) Occipital cortex

Occipital cortex is the neurological site of lesion which causes homonymous hemianopia with macular sparing.

169. In which chamber of the eye, given keratome is used in Phaco surgery?




a) Anterior chamber
b) Posterior chamber
c) Sclera
d) Lens

- Keratome blade: **a surgical instrument used for making an incision in the cornea in cataract operations**.
- Clear corneal incision and it is used to enter the anterior chamber

170. A 60-year-old women comes to the emergency department due to a sudden onset of severe pain in her left eye with blurred vision, nausea, and vomiting. the symptoms began a few minutes ago, while was watching a movie in a theatre. Her temperature is 36.8c (98.4F), blood pressure is 140/90mm.hg, Pulse is 82/min, and respirations are 14/min, on examination reveals

www.medicalcoupons.in

decreased visual activity. Her left eye appears red, with a hazy cornea, shallow anterior chamber and dilated, fixed pupil. Her left eye is stony hard to touch, what is most probable diagnosis?



a) Primary open angle glaucoma
b) Conjunctivitis
c) Acute angle closure glaucoma
d) Anterior uveitis

Acute angle closure is an urgent but uncommon dramatic symptomatic event with blurring of vision, painful red eye, headache, nausea, and vomiting. Diagnosis is made by noting high intraocular pressure (IOP), corneal edema, shallow anterior chamber, and a closed angle on gonioscopy. Medical or surgical therapy is directed at widening the angle and preventing further angle closure. If glaucoma has developed, it is treated with therapies to lower IOP.

171. The patient was brought to the emergency. The patient was hit by a tennis ball…. And noted to have hyphema. What is the source of bleed in this patient?

a) Anterior ciliary artery
b) Posterior ciliary artery
c) Major arterial circle of iris
d) Minor arterial circle of iris

Hyphema is the collection of blood in the anterior chamber of the eye. The most common cause of hyphema is **blunt trauma**, though spontaneous hyphemas can occur in the setting of sickle cell disease or other increased bleeding states. Hyphemas are graded based on the degree of blood obscuring the cornea. major arterial circle of iris is the source of bleed.

www.medicalcoupons.in

172. A 2-year-old Child presented with leucocoria in the right eye since 2 months. On examination a total retinal detachment was the present in the same eye. Ultrasound B scan revealed a heterogonous sub- retinal mass with calcification. The most probable clinical diagnosis is:



a) Coats – Disease
b) Retinoblastoma
c) Toxocariasis
d) Retinal tuberculoma

Retinoblastoma is **an eye cancer that begins in the retina** — the sensitive lining on the inside of your eye. Retinoblastoma most commonly affects young children, but can rarely occur in adults. Your retina is made up of nerve tissue that senses light as it comes through the front of your eye

173. Which of the following is Contra Indicated in Acute Intermittent Porphyria?
a) Thiopentone
b) Propofol
c) Ethomidate
d) Ketamine

hypersensitivity to barbiturates. acute intermittent porphyria $^{\Delta}$ (barbiturates enhance porphyrin synthesis, and therefore are absolutely contraindicated in patients with a history of acute intermittent porphyria)

www.medicalcoupons.in

174. Identify the Following Instrument.



a) LMA PROSEAL
b) Tight Seal
c) High Seal
d) Classical LMA

ProSeal LMA has **a gastric drainage tube**, placed lateral to the main airway tube. The gastric drainage tube forms a channel for regurgitated gastric contents and prevents gastric insufflation and pulmonary aspiration.

175. Identify the Following Tube.



a) Univalent tube
b) Single Endo tracheal Lumen
c) Double Lumen Endo tracheal Tube
d) Ryles tube

www.medicalcoupons.in

Double lumen tubes play a crucial role in:

1. Airway management during thoracic surgery.
2. Anatomical lung separation or isolating normal from the diseased lung in situations such as: Massive hemorrhage from one lung. ...
3. Differential lung protective ventilation in case of lung transplantation patients.

176. A lady was taken for C-section operation. the duration of surgery was more than anticipated and the patient lands into post op hyperventilation. which of the following could be the reason for that?

a) High Spinal Anaesthesia

b) Malingering

c) Hypoxic Encephalopathy

d) Cranial nerve palsy

High spinal/regional block: **spread of local anaesthetic affecting the spinal nerves**. **above T4**. The effects are of variable severity depending upon the maximum level. that is involved, but can include cardiovascular and/or respiratory compromise.

177. A person was admitted into hospital for tibial fracture and a cast was applied. After 3 weeks the cast was removed and another cast was applied. The patient was given crutches and asked to walk. What is the stage of healing of the bone?

a) Inflammation Phase

b) Remodeling Phase


c) Proliferative Phase

d) Infiltration Phase

Remodeling or also known as maturation phase is **the fourth and final phase in wound healing** and lasts from 21 days up to 2 years. In this final and longest phase, collagen synthesis is ongoing in order to strengthen the tissue. Remodeling occurs as wound continues to contract and fibers are being reorganized.

www.medicalcoupons.in

178. Most common Associated Site of Tb Infection?



a) 1
b) 2
c) 3
d) 4

Bone tuberculosis is simply a form of TB that affects **the spine**, the long bones, and the joints

179. Identify the condition?



www.medicalcoupons.in

a) DDH (Developmental Dysplasia of Hip)
b) Ankle Foot Arthrosis Brace
c) Perthes Disease
d) CTEV (Congential Talipes Equino Varus)

In all cases of DDH, the socket (acetabulum) is shallow, meaning that the ball of the thighbone (femur) cannot firmly fit into the socket. Sometimes, the ligaments that help to hold the joint in place are stretched. The degree of hip looseness, or instability, varies among children with DDH.

180. A patient had trauma and was applied cast. On removal of the cast the doctor pressed the fracture site and described as movable or soft. True among the following?



a) Patellar Cast with Soft Callus Formed
b) Patellar Cast with Hard Callus Formed
c) Patellar Cast with Bone Remodelling
d) Patellar cast with hematoma.

The callus holds the bone together, but isn't strong enough for the body part to be used. Over the next few weeks, the soft callus becomes harder. By **about 2–6 weeks**, this hard callus is strong enough for the body part to be used.

www.medicalcoupons.in

181. All are true except



a) Most Common Infection of Finger

b) Can Progress into Osteomyelitis

c) Treatment Include Incision& Drainage and Antibiotics

d) A Part of Nail Is Removed as Treatment Option

Paronychia is the inflammation of the tissue that immediately surrounds the nail. It is the most common infection of the hand and it occurs after disruption of the seal between the nail fold and the nail plate due to penetrating trauma, nail biting, manicures or a hangnail

182. A parents brought child with the following image of thigh (image has protruded bone with discharge). Parents confirmed there is history of bone pain and discharge from the site prior to the protrusion. what could be the diagnosis?



a) Acute Osteomyelitis

b) Chronic Osteomyelitis

c) Involucrum with Cloaca

d) Osteosclerosis

www.medicalcoupons.in

Chronic osteomyelitis, defined as continuous infection of a low-grade type or of a recurrent type, is characterized predominantly by bony sclerosis, periosteal new bone formation, and the presence of sequestra and/or draining sinuses.

183. A 45-year-old female came to outpatient department with complaints of pain fatigue. Diagnosed tumor of epiphysis



a) GCT
b) Chondrosarcoma
c) Osteosarcoma
d) Ewings Sarcoma

A giant cell tumor is a **rare, aggressive non-cancerous tumor**. It usually develops near a joint at the end of the bone. Most occur in the long bones of the legs and arms. Giant cell tumors most often occur in young adults when skeletal bone growth is complete.

184. What is the Diagnosis for the given?




www.medicalcoupons.in

a) Gall Stones
b) Carcinoma of Gall bladder
c) Mixed gall stones
d) Porcelain gallbladder

Gallbladder cancer is a relatively uncommon cancer. Most often it is found after symptoms such as abdominal pain, jaundice and vomiting occur, and it has spread to other organs such as the liver. It is a rare cancer that is thought to be related to gallstones building up, which also can lead to calcification of the gallbladder, a condition known as porcelain gallbladder. Porcelain gallbladder is also rare. Some studies indicate that people with porcelain gallbladder have a high risk of developing gallbladder cancer

185. Identify the given image




a) P. Vivax
b) P.falciparum
c) P.malariae
d) P.ovale

Plasmodium falciparum is a **unicellular protozoan parasite of humans**, and the deadliest species of Plasmodium that causes malaria in humans. The parasite is transmitted through the bite of a female Anopheles mosquito and causes the disease's most dangerous form, falciparum malaria.

www.medicalcoupons.in

186. Identify the give Image



a) Teniasolium
b) Trichuris spiralis
c) Trichuris trichiura eggs
d) Wuchereria bancrofti

Trichuris trichiura eggs are **50-55 micrometers by 20-25 micrometers**. They are barrel-shaped, thick-shelled and possess a pair of polar "plugs" at each end. The eggs are unembryonated when passed in stool. Trichuris trichiura eggs are **50-55 micrometers by 20-25 micrometers**. They are barrel-shaped, thick-shelled and possess a pair of polar "plugs" at each end. The eggs are unembryonated when passed in stool.


187. A 30-year-old patient of valvular heart disease is having fever for last 2 weeks. On physical examination the following features are noticed in this patient. Which of these will be the abdomen examination finding of this patient?

a) Ascites
b) Portal Hypertension
c) Splenomegaly
d) Renal artery Bruit

An enlarged spleen is the result of damage or trauma to the spleen from any of several different medical conditions, diseases, or types of physical trauma. **Infections, liver problems, blood cancers, and metabolic disorders** can cause your spleen to become enlarged, a condition called splenomegaly.

188. A 70-year-old man collapsed in his house and was rushed to the hospital. On arrival his ECG tracing was as shown below. Which of the following is the best intervention for this patient?

www.medicalcoupons.in



a) Use of Automated external defibrillator
b) Injection of Vasopressin 40 IU
c) Injection of adrenaline 1 mg iv
d) Injection of atropine 0.5mg iv

Adrenaline 1:1000 (1mg/mL) Solution for injection is used in **life threatening emergencies such as severe allergic reactions or cardiac arrest**. 2. you have atherosclerosis which is a narrowing and hardening of the body's blood vessels

189. A 60-year-old woman presents with weakness in right arm for four hours' duration. The weakness gradually became lesser and resolves. She is a known case of Hypertension and BMI is elevated. Which of the following is the probable diagnosis of this case?

a) Transient ischemic attack
b) Compressive Neuropathy
c) Transverse Myelitis
d) Stroke

weakness in the right arm for 4 hours, which self resolves for a patient with elevated BMI and Hypertension is suggestive of stroke

190. A 35-year-old recently divorced male is suffering from palpitations, headache and diaphoretic episodes on recurrent basis. He has visited the psychiatrist multiple times. What should be the next best step for management of this case?

a) He should visit psychologist instead of psychiatrist
b) Check for catecholamines levels to rule out pheochromocytoma
c) Check for hyperthyroidism
d) Increase dose of anti-anxiety medication

High levels of catecholamines, vanillylmandelic acid (VMA), or metanephrine can mean that **an adrenal gland tumor (pheochromocytoma) or another type of tumor that makes catecholamines is present**.

www.medicalcoupons.in

191. A patient has presented with gross abdominal distention and the following test is being performed. Comment on the finding being elicited?



a) Fluid thrill
b) Shifting dullness
c) Puddle sign
d) Hingorani sign

In medicine, the fluid wave test or fluid thrill test is **a test for ascites** (free fluid in the peritoneal cavity). It is performed by having the patient (or a colleague) push their hands down on the midline of the abdomen. The examiner then taps one flank, while feeling on the other flank for the tap.

192. An elderly patient with Hypertension fell down in his bathroom. His BP was 220/110 on admission. Which of the following statements is correct about this patient?

a) Start anti platelet drugs
b) Platelets are transfused even if the platelet count is not low
c) Reduce the BP to <120:
d) Putamen is the most common site

The putamen is **the most common site for hypertensive ICH**. Hemorrhages may remain localized to the putamen; enlarge to involve the internal capsule, corona radiata, centrum semiovale, or temporal lobe; or rupture into the ventricular system

www.medicalcoupons.in

193. A 70-year-old man has recurrent episodes of exertional syncope and recurrent episodes of chest pain on climbing stairs. He was diagnosed as a case of valvular aortic stenosis. Which of the following is the radiological finding of XRC of this patient?

a) Dilated aortic Root:
b) Widening of aortic knob
c) Widening of vascular pedicle
d) Post stenotic dilatation of aorta

Variable appearance on chest radiographs depending on stage and severity of the disease.

In early disease, the chest radiograph can be entirely normal or it may show dilatation of the ascending aorta with a normal heart size. Differentiation with hypertension can usually be made as in hypertension the entire descending aorta is enlarged

194. Elderly Chronic alcoholic patient has presented to OPD today. The following findings were noted in the patient. What is the best investigation to be done in this patient?





a) Plasma ascorbate levels
b) Factor8 assay
c) RBC transketolase activity
d) Thrombin time

ascorbate is the **most effective aqueous-phase antioxidant in human** blood plasma and suggest that in humans ascorbate is a physiological antioxidant of major

www.medicalcoupons.in

importance for protection against diseases and degenerative processes caused by oxidant stress.

195. A patient presents with complaints of palpitations. On examination her heart rate is 110 / min with Respiratory rate of 10 / min. She tells you that her mother has recently expired due to renal cell cancer. What investigation would you perform to manage this case?

a) Serum Hydroxy Indole Acetic Acid levels (H.I.A.A)
b) Serum Catecholamines
c) 24-hour Urine VMA (Vanillylmandelic acid) for 24 hours
d) 24-hour fractionated Metanephrine levels

- If your 24-hour metanephrine level is between 1 and 2 times the normal amount, there is about a 30% chance you have a pheochromocytoma or paraganglioma.
- If your 24-hour metanephrine level is twice the normal level or higher, it's likely that you have a pheochromocytoma or paraganglioma.

196. Comment on serological report of this hepatitis B patient

IgG Anti HBc	Positive
HBsAg	Negative
HBeAg	Negative
Anti HBs	positive

a) Acute hepatitis B, High infectivity
b) Chronic hepatitis B, High infectivity
c) Chronic hepatitis B, low infectivity
d) Previous infection with hepatitis B

HBeAg negative shows that the virus is not actively replicating and the patient is not infectious

HbsAg negative shows the patient is not currently infected with the virus

Anti HBs, IgG Anti HBc both are positive showing that the antibodies are formed for the previous infection

197. A 30-year-old female patient presented with pain in joints of hand with rash on dorsum of hand with extensive skin induration and skin tightening. Comment on the diagnosis.

a) Rheumatoid arthritis
b) Psoriatic arthritis

www.medicalcoupons.in

c) Osteoarthritis
d) Scleroderma

Psoriatic arthritis is **a form of arthritis that affects some people who have psoriasis** — a disease that causes red patches of skin topped with silvery scales. Most people develop psoriasis years before being diagnosed with psoriatic arthritis.

198. A 40-year-old patient came to OPD with symptoms of excessive sleepiness and lethargy all the time. On physical examination the clinical features depicted in the images were seen. Per abdominal examination shows no free fluid in abdomen. Liver and spleen are not palpable. He tells of regular Alcohol consumption. Which is the most probable diagnosis?





a) Hepatolenticular degeneration
b) Alcoholic Cirrhosis
c) Portal Hypertension
d) Budd Chiari Syndrome

Alcoholic hepatitis is a condition caused by continued alcohol use that results in long-term (chronic) inflammation in your liver. Alcoholic cirrhosis is an advanced stage of alcoholic liver disease that **causes your liver to become stiff, swollen, and barely able to do its job**.

199. A 35-year-old woman presented with complaints of palpitations and weight loss. On examination sinus tachycardia is present, and the following lesion was noticed on the shin of the patient. What is the clinical diagnosis?



www.medicalcoupons.in

a) Hypothyroidism
b) Hyperthyroidism
c) Filariasis
d) Milroy disease

Hyperthyroidism, also called overactive thyroid, is a **condition where the thyroid releases high levels of thyroid hormone into the body**. This condition can make your metabolism speed up. Symptoms of hyperthyroidism include a rapid heartbeat, weight loss, increased appetite and anxiety.

200. On Post-operative day 2, routine X-Ray of patient was performed. The patient is not complaining of any respiratory distress and is comfortable in hospital bed. On bilateral auscultation equal air entry is heard. What is the next step in management of patient?

a) Wait and watch and give supplemental oxygen if required
b) Perform needle thoracotomy urgently
c) Do a thoracostomy with wide bore tube and under water seal
d) Perform elective intubation and positive pressure ventilation

The patient condition is getting better and since there is no complaints of any respiratory distress and equal air entry is heard in bilateral auscultation, we just continue the observation of the patient and provide oxygen if needed.

201. A 55-year-old man has chest pain on rest for last 6 hours. ECG was performed on admission and showed findings of myocardial ischemia. His biomarkers report was sent to lab for diagnostic workup. Which of the following will help in diagnosis?

a) LDH1
b) LDH2
c) LDH3
d) LDH4

The LDH-1 isoenzyme is found predominately in **cardiac muscle**, LDH-2 is found primarily in the reticuloendothelial system, LDH-3 predominates in the lungs, LDH-4 in the kidneys, and LDH-5 in the liver and skeletal muscle.

202. A 50-year-old man presents with recurrent episodes of chest pain on exertion. He is diagnosed as a case of chronic stable angina. On routine monitoring BP is found to be consistently elevated. Which of the following is the best drug for management of this patient?

a) Enalapril
b) Amlodipine

www.medicalcoupons.in

c) Atenolol

d) Thiazides

Atenolol belongs to a group of medicines called beta blockers. It's **used to treat high blood pressure and irregular heartbeats** (arrhythmia). It can also be used to prevent chest pain caused by angina. If you have high blood pressure, taking atenolol helps prevent future heart disease, heart attacks and strokes.

203. A 30-year-old man has presented with complaints of fever for 3 days with SOB. On examination on right mammary area bronchial breathing is heard with normal air entry bilaterally. Comment on the diagnosis.

a) Right lower lobe pneumonia


b) Right middle lobe pneumonia

c) Right sided hydro-pneumothorax

d) Right sided Hydrothorax

Bronchial sounds are high pitched & usually heard over the trachea. Timing includes an inspiratory phase that is less than the expiratory phase. If bronchial sounds are heard in the actual lung fields, this may indicate **consolidation**.

204. What is the impression from the following CXR?



a) Chemo-port device

b) Pacemaker

c) Artifact

d) Travel tract for bypassing accessory pathway

www.medicalcoupons.in

A pacemaker is a **device that sends small electrical impulses to the heart muscle** to maintain a suitable heart rate or to stimulate the lower chambers of the heart (ventricles). A pacemaker may also be used to treat fainting spells (syncope), congestive heart failure and hypertrophic cardiomyopathy.

205. Prophylactic CNS radiation is given for which of the following cancers?

a) Liver cancer

b) Small cell cancer of lung

c) Prostate cancer

d) Seminoma

Prophylactic cranial irradiation is now known to **improve survival** to a significant degree in small-cell lung cancer (SCLC) patients; this is in addition to its established role in preventing the disabling symptoms of brain metastases.

206. An elderly patient fell in the bathroom in early morning hours and was rushed to the casualty. On arrival GCS is E4V4M6 and urgent NCCT head was performed. Comment on the diagnosis.

a) Extradural hemorrhage

b) Subdural hemorrhage

c) Diffuse axonal injury

d) Intracerebral haemorrhage

A subdural hematoma is **a type of bleed inside head**. More precisely, it is a type of bleed that occurs within the skull of head but outside the actual brain tissue. The brain has three membranes layers or coverings (called meninges) that lay between the bony skull and the actual brain tissue

207. A young woman presents as shown below, which is triggered by cold temperatures or emotional stress. Episodes usually affect the fingers and toes along with numbness and pain in the affected areas. There is no underlying illness or comorbidity in the patient. Comment on the diagnosis.




www.medicalcoupons.in

a) Primary Raynaud phenomenon
b) Secondary Raynaud phenomenon
c) Harlequin sign
d) Ainhum

Secondary Raynaud's is caused by **an underlying disease or condition**. It is especially common in people with connective tissue diseases. Some of these diseases reduce blood flow to the fingers and toes by causing the blood vessel walls to thicken and the vessels to constrict too easily

208. Comment on the diagnosis of the Flow volume curve shown below



a) COPD
b) Interstitial lung disease
c) Retrosternal Goiter
d) Tracheal stenosis

The F/V curve is **a representation of air stream volume pattern during inspiration and expiration**. It exhibits the obstructive, restrictive and mixed pattern pathological conditions. ... The classic flow-volume loop shapes describe the obstructive and restrictive lung pathology as well as anatomical abnormalities. Patients with chronic obstructive pulmonary disease (COPD) exhibit **increases in lung volume due to expiratory airflow limitation**. Increases in lung volumes may affect upper airway patency and compensatory responses to inspiratory flow limitation (IFL) during sleep.

209. A 28-year-old pregnant lady presents for routine antenatal check-up. Which Arterial Blood gas abnormality is expected to be present in this case?
a) Respiratory acidosis
b) Respiratory Alkalosis
c) Metabolic Acidosis
d) Metabolic Alkalosis

www.medicalcoupons.in

Progesterone levels are increased during pregnancy. Progesterone causes stimulation of the respiratory center, which can lead to respiratory alkalosis. Chronic respiratory alkalosis is a common finding in pregnant women.

210. Rabies virus will use which kind of molecular motor to cause involvement of Central nervous system from the site of inoculation into the host?

a) Dynein
b) Kinesin
c) Kinin
d) Actin

Dynein is a family of cytoskeletal motor proteins that move along microtubules in cells. They convert the chemical energy stored in ATP to mechanical work. Dynein **transports various cellular cargos**, provides forces and displacements important in mitosis, and drives the beating of eukaryotic cilia and flagella. Rabies virus uses this to cause involvement of central nervous system from the site of inoculation into the host.

211. A 31-year-old car driver was involved in an accident. He is brought to ER with complaints of breathlessness. On physical examination vitals are PR-110 / min, BP 90/60 mm Hg and temp of 37 degrees. Extensive bruising is noticed on right side of chest. An urgent bedside CXR was performed. What is the best management of this case?

a) Thoracotomy
b) Thoracostomy
c) Elective intubation and PPV
d) Fluid resuscitation

The patient is suspected to have hemothorax due to the contusion and so first we go for elective intubation and PPV

212. A 60-year-old woman a known case of lung cancer is having shortness of breath. On examination her neck veins are engorged showing gross elevation of JVP, Heart sounds are distant and breath sounds appear reduced on left side infra-scapular area. Requisition for CXR is done and is shown below. Comment on the diagnosis of the patient.

a) Cardiac Tamponade
b) Congestive heart failure
c) Bilateral pleural effusion
d) SVC Syndrome


Cardiac tamponade is a **serious medical condition in which blood or fluids fill the space between the sac** that encases the heart and the heart muscle. This places extreme pressure on your heart. The pressure prevents the heart's ventricles from expanding fully and keeps your heart from functioning properly.

213. An elderly patient with malignancy develops haematuria with dipstick performed on urine sample showing 4+ proteinuria. Which of these is the possible diagnosis of this patient?

a) MGN
b) MPGN
c) FSGS
d) Diffuse Glomerulosclerosis

Membranous glomerulonephritis (MGN) is a specific type of GN. MGN develops **when inflammation of your kidney structures causes problems with the functioning of your kidney**. MGN is known by other names, including extramembranous glomerulonephritis, membranous nephropathy, and nephritis.

214. In the given tracing shown below, HV interval denotes?



a) AV intranodal conduction
b) Septal Activation of bundle of His
c) Antegrade conduction from Bundle of His to bundle branches
d) Antegrade conduction from Bundle to His to Purkinje fibers

HV interval measures **the conduction time through the distal His- Purkinje tissue** and is measured from the onset of the His-bundle deflection to the earliest ventricular activation.

www.medicalcoupons.in

215. A 60-year-old patient has right sided homonymous hemianopia with macular sparing. Locate the neurological site of lesion in this patient.

a) Optic chiasma
b) Optic tract
c) Medial geniculate body
d) Occipital cortex

Homonymous hemianopsia can be congenital, but is usually caused by brain injury such as from stroke, trauma, tumors, infection, or following surgery.
Vascular and neoplastic (malignant or benign tumours) lesions from the optic tract, to visual cortex can cause a contralateral homonymous hemianopsia. Injury to the right side of the brain will affect the left visual fields of each eye. The more posterior the cerebral lesion, the more symmetric (congruous) the homonymous hemianopsia will be.

216. A 60-year-old man has presented with chronic projectile vomiting and significant weight loss. Which of the following electrolyte abnormality is seen in this patient?

a) Hypokalemic hypochloremic Metabolic alkalosis with Hyponatremia
b) Hypokalemic hypochloremic Metabolic alkalosis with Hypernatremia
c) Hypokalemic hypochloremic Metabolic alkalosis with Hypercalcemia
d) Hypokalemic hypochloremic Metabolic alkalosis with Hypomagnesemia

Hypochloremic alkalosis results from **either low chloride intake or excessive chloride wasting**. Whereas low chloride intake is very uncommon, excessive chloride wasting often occurs in hospitalized children, usually as a result of diuretic therapy or nasogastric tube suctioning.

217. A 35-year-old patient with ulcerative colitis develops ulcer on the anterior compartment of the leg. Comment on the diagnosis of patient.

a) Venous ulcer
b) Erythema migrans
c) Erythema Nodosum
d) Pyoderma gangrenosum

Pyoderma gangrenosum is a rare condition that causes large, painful sores (ulcers) to develop on your skin, most often on your legs. The exact causes of pyoderma gangrenosum are unknown, but it appears to be a disorder of the immune system

www.medicalcoupons.in

218. A patient presented with back pain. Labs show a grossly elevated ESR, X- Ray skull shows multiple punched out lesions. Which is the most important investigation to be done for this case?

a) Serum acid phosphatase
b) CT head with Contrast
c) PET scan
d) Serum Electrophoresis

back pain, elevated ESR X-ray showing multiple punched out lesions leading to multiple myeloma. Serum protein electrophoresis is used to identify patients with multiple myeloma and other serum protein disorders. Electrophoresis separates proteins based on their physical properties, and the subsets of these proteins are used in interpreting the results. Plasma protein levels display reasonably predictable changes in response to acute inflammation, malignancy, trauma, necrosis, infarction, burns, and chemical injury.

219. **An AIDS Positive 30-year-old male patient has developed fever, vomiting and meningismus. Which of the following test: will have help in rapid diagnosis of Cryptococcal meningitis**

a) India Ink preparation of CSF
b) Blood Lateral flow assay for Cryptococcal antigen
c) Blood culture
d) CSF Cultures

he recent development of the **Cryptococcal antigen lateral flow assay (LFA)**, a commercially available rapid diagnostic test that detects capsular polysaccharide antigens of the four major cryptococcal serotypes (A and D for C. neoformans and B and C for C.

220. A 30-year-old man who works in a glass factory presents with complaints of progressive shortness of breath on exertion. He is predisposed to development of which of the following?

a) Histoplasmosis
b) Tuberculosis
c) Mesothelioma
d) Pleural calcification

The common Health hazards associated with glass manufacturing processes include **noise, exposures to respirable airborne particulates [especially silica (SiO2) that leads to chronic effects such as decreased pulmonary function, lung disease, complicated pneumoconiosis**

www.medicalcoupons.in

221. **A 12-year-old girl has developed Acute Immune thrombocytopenic purpura (ITP) and is managed with steroids. There is recurrence of low platelet count once steroids were stopped. On current physical examination there is no petechiae or any physical evidence of bleeding. Which is the next line of treatment?**

a) Continue steroids as long they are effective

b) Splenectomy is greater than 2 relapses

c) Wait and watch as self-resolution will occur in most cases of acute ITP in children

d) Rho Immunoglobulin IV

since the patient was under steroid therapy for Immune thrombocytopenic purpura (ITP). Since there is no petechiae or any bleeding other than less platelet count , most probable next step should be to wait and observe as acute ITP is self-resolved in most cases of children.

222. What is the primary defect in vitamin D resistant rickets?

a) Defective phosphate absorption from gut

b) Phosphate wasting from proximal renal tubule

c) Defective Calcium absorption from gut

d) Calcium wasting from proximal renal tubules

Hypophosphatemic rickets is a disorder characterized by hypophosphatemia, defective intestinal absorption of calcium, and rickets or osteomalacia unresponsive to vitamin D. It is usually hereditary.

223. A 31-year-old alcoholic had his last drink 48 hrs ago. He develops anger and rage attacks and appears unable to identify faces of family members and can see inanimate objects. He has been brought to the ER. What is your diagnosis?

a) Delirium tremens

b) Alcoholic hallucinosis

c) Korsakoff psychosis

d) Wernicke's encephalopathy

Delirium tremens (DTs) is the **most severe form of ethanol withdrawal**, manifested by altered mental status (global confusion) and sympathetic overdrive (autonomic hyperactivity), which can progress to cardiovascular collapse. Minor alcohol withdrawal is characterized by tremor, anxiety, nausea, vomiting, and insomnia


www.medicalcoupons.in

224. A 25-year-old male patient is having loose stools 18 times per day and his urine frequency is reduced. What will be the correct finding of this patient?

a) Urine sodium >40
b) Kidney biopsy showing Glomerular damage
c) Hyperosmolar urine
d) Post renal azotemia

Both high and low urine osmolality have several causes. Some of these, such as dehydration, are relatively easy to treat. Others can be more serious or ongoing. Here due to severe dehydration, hyperosmolar urine can be found.

225. A 50-year-old lady with Hypertension was admitted with breathlessness in supine position and palpitations for last 4 days. On physical examination her heart rate is 120 / min with congested neck veins. Per abdomen shows liver palpable 3 cm below costal margin. The finding shown in the image below is?



a) Deep vein thrombosis
b) Superficial venous thrombophlebitis
c) Pitting Pedal edema
d) Lymphedema

Pitting edema is a specific type of edema that is associated with pitting or indentation in the affected areas, and is often caused by conditions that lead to the pooling of blood in the feet or legs. The excess fluid that builds up in pitting edema is mainly composed of water.

226. A 6-year-old boy has presented with multiple episodes of hematemesis. On admission his vitals are stable and splenomegaly is noted. The mother tells that at birth the boy had developed jaundice and an exchange transfusion procedure had been performed in the same hospital. Diagnosis is

a) Budd Chiari Syndrome
b) Sinusoidal fibrosis
c) Portal vein thrombosis
d) Non cirrhotic portal fibrosis

www.medicalcoupons.in

Portal vein thrombosis is **blockage or narrowing of the portal vein** (the blood vessel that brings blood to the liver from the intestines) by a blood clot. Most people have no symptoms, but in some people, fluid accumulates in the abdomen, the spleen enlarges, and/or severe bleeding occurs in the esophagus.

227. A 6-year-old child weighs 12 kg. The urine output of the child is only 0.5ml / kg / hour for last 12 hours and serum creatinine has shown a doubling over the baseline. What is stage of AKI by KDIGO criteria?

a) 1

b) 2

c) 3

d) 4

AKI stage	Serum creatinine (SCr)	Urine output
1	1.5-1.9 times baseline OR ≥ 0.3 mg/dL increase	< 0.5 mL/kg/h for 6-12 hours
2	2.0–2.9 times baseline	< 0.5 mL/kg/h for ≥ 12 hours
3	3.0 Times baseline OR Increase in SCr to ≥ 4.0 mg/dL OR Initiation of RRT OR Decrease in eGFR to < 35 mL/min/1.73 m² in patients < 18 years	< 0.3 mL/kg/h for ≥24 hours OR Anuria for ≥ 12 hours

228. A girl is having honey crust injuries on face which was diagnosed by her paediatrician as impetigo. She has now been brought your hospital with facial puffiness and not eating well. When you work up the patient, which of the following will be the most common finding encountered in this case:

a) RBC in urine

b) Nephrotic range proteinuria

c) Derangement of liver enzymes

d) Focal neurological deficit

Transient proteinuria and hematuria **may occur during impetigo** and resolve before renal involvement develops. Antibiotic treatment does not prevent the development of glomerulonephritis, but it limits the spread of the disease to other individuals. Other complications may include the following: Scarlet fever.


229. A 61-year-old known case of Diabetes mellitus has been noticed on laboratory work up to have microalbuminuria. What is the site of reabsorption of proteins in the ultra-filtrate?

www.medicalcoupons.in

a) Podocytes
b) Proximal convoluted tubule
c) Distal convoluted tubule
d) Collecting duct

The proximal convoluted tubule is where a majority of reabsorption occurs. About **67 percent** of the water, Na^+, and K^+ entering the nephron is reabsorbed in the proximal convoluted tubule and returned to the circulation.

230. A 35-year-old patient presents with high grade fever, chills and cough with copious production of purulent sputum. An x-ray examination was made as shown in the image, what is the most likely diagnosis?




a) Rt middle lobe consolidation
b) Rt lower lobe consolidation
c) Atelectasis
d) Pneumothorax

Consolidation refers to the alveolar airspaces being filled with fluid (exudate/transudate/blood), cells (inflammatory), tissue, or other material. The list of causes of consolidation is broad and includes: **pneumonia**. adult respiratory distress syndrome (ARDS)

231. An 65 yrs old hypertensive patient on irregular medication has headache and a BP-220/110, CT head is shown below

www.medicalcoupons.in



a) Intraventricular bleed
b) Putaminal bleed
c) Sub arachnoid hemorrhage
d) Pontine hemorrhage

The putamen is a common site of **hypertensive cerebral hemorrhage**. Such hemorrhages show a large range of possible presentations – from enormous hemorrhages involving the white matter of the hemispheres and the ventricular system, to cases occurring without causing any symptoms or neurological signs.

232. Pt with History of Recurrent UTI, the IVP image is given, Identify the location of insertion of ureter arising from left superior pelvis?




a) Below and medial to trigone of bladder
b) Above the dome of the bladder
c) Prostatic urethra
d) Perineum

www.medicalcoupons.in

The **trigone** (a.k.a. vesical trigone)is a smooth triangular region of the internal urinary bladder formed by the two ureteric orifices and the internal urethral orifice.

233. Identify the marked structure on CT head



a) Optic Nerve
b) Optic Chiasma
c) Optic tract
d) P Superior Ophthalmic Artery

The marked part in the CT scan is the optic nerve

234. 5yr old boy is unable to pronate and supinate since childhood?




www.medicalcoupons.in

a) Radioulnar synostosis
b) Radial head dislocation
c) Monteggia fracture
d) Myositis ossificans

Radioulnar synostosis is a **rare condition in which the two bones of the forearm** — the radius and the ulna — are abnormally connected. This limits rotation of the arm. Radioulnar synostosis is usually congenital). It can also occur as the result of a forearm fracture or trauma.


235. Identify the marked structure shown by the arrow:



a) Pons
b) Medulla
c) Midbrain
d) Spinal cord

The marked structure in the above radiological finding is the mid-brain

236. 45 year old Male with Heart Disease undergoes a chest X-ray, what is the diagnosis?




www.medicalcoupons.in

a) Prosthetic aortic valve
b) Prosthetic mitral valve
c) Pericardial effusion
d) Implantable pacemaker

An artificial heart valve is a one-way valve implanted into a **person's heart to replace a valve that is not functioning properly (valvular heart disease)**. The mechanical prosthetic heart valves are **very durable and guarantee 20-30 years of functioning in ideal conditions**. The coating with pyrolytic carbon in mechanical valves also decreases the chance of blood clotting on the valve surface.

237. The Given Skull X-ray finding is seen in?



a) Hemolytic anemia
b) Multiple myeloma
c) Increased ICT
d) Osteopetrosis

Hemolytic anemia is **a disorder in which red blood cells are destroyed faster than they can be made**. The destruction of red blood cells is called hemolysis. Red blood cells carry oxygen to all parts of your body. If you have a lower than normal amount of red blood cells, you have anemia.

www.medicalcoupons.in


238. X-ray of a 2-year-old child is given, what is the possible diagnosis?



a) Pneumastosis intestinalis
b) Pneumoperitoneum
c) Subcutaneous emphysema
d) Duodenal atresia

Pneumoperitoneum is **the presence of air or gas in the abdominal (peritoneal) cavity**. It is usually detected on x-ray, but small amounts of free peritoneal air may be missed and are often detected on computerized tomography (CT).

239. Identify the pathology on given Barium enema image?



www.medicalcoupons.in

a) Intussusception
b) Ulcerative colitis
c) Cecal cancer
d) Ileocecal TB

Intussusception is a **condition in which one segment of intestine "telescopes" inside of another**, causing an intestinal obstruction (blockage). Although intussusception can occur anywhere in the gastrointestinal tract, it usually occurs at the junction of the small and large intestines


240. A 45-year-old female came for post-surgery follow up, A CXR was taken what is the most likely diagnosis?



a) Malignancy
b) Fibroadenoma
c) Emphysema
d) Left sided mastectomy

In the picture we can clearly absence of left breast.

241. Identify the Given HSG image?



www.medicalcoupons.in

a) Bicornuate
b) Unicornuate
c) Septate
d) Didelphys

Unicornuate uterus is **a rare genetic condition in which only one half of a girl's uterus forms**. A unicornuate uterus is smaller than a typical uterus and has only one fallopian tube. This results in a shape often referred to as "a uterus with one horn" or a "single-horned uterus.

242. A female patient with dysphagia, Barium swallow image is shown, what is the likely diagnosis?



a) Diffuse esophageal spasm
b) Esophageal cancer
c) Achalasia cardia
d) Esophagitis

Achalasia Cardia is a **rare disorder** that makes it difficult for food and liquid to pass from the swallowing tube connecting your mouth and stomach (esophagus) into your stomach. Achalasia Cardia occurs when nerves in the esophagus become damaged. The hallmark of the achalasia pattern is a **gradually tapering, smooth, conical narrowing (bird's beak or rat-tail appearance) of** the distal esophageal segment which extends about 1-3 cm in length

243. A 65-Year-old female fell down in the bathroom &sustained head injury, Ct scan was performed, what is likely diagnosis?

www.medicalcoupons.in



a) Extradural hematoma
b) Intracranial bleed
c) Thrombosis
d) Subdural hematoma

A subdural hematoma is **a type of bleed inside head**. More precisely, it is a type of bleed that occurs within the skull of head but outside the actual brain tissue. The brain has three membranes layers or coverings (called meninges) that lay between the bony skull and the actual brain tissue

244. Identify the given MRI image




www.medicalcoupons.in

a) B/L Ureterocele
b) Hydronephrosis
c) Staghorn calculi
d) Bladder calculi

A ureterocele is **a swelling at the bottom of one of the ureters**. Ureters are the tubes that carry urine from the kidney to the bladder. The swollen area can block urine flow. A ureterocele is a birth defect


245. What structure is the arrow pointing to



a) Medial epicondyle
b) Trochlea
c) Capitulum
d) Olecrenon

In human anatomy of the arm, the capitulum of **the humerus** is a smooth, rounded eminence on the lateral portion of the distal articular surface of the humerus.

246. Identify the Given Chest Radiograph?




www.medicalcoupons.in

a) Pneumothorax
b) Hydropneumothorax
c) Pericardial effusion
d) Emphysema

A pneumothorax is **a collapsed lung**. A pneumothorax occurs when air leaks into the space between your lung and chest wall. This air pushes on the outside of your lung and makes it collapse. A pneumothorax can be a complete lung collapse or a collapse of only a portion of the lung.

247. 33 yr. old female with fever, weight loss, elevated levels of ACE levels with hilar lymphadenopathy on x ray, Probable diagnosis?



a) TB
b) Sarcoidosis
c) Silicosis
d) Small cell cancer

Sarcoidosis is **an inflammatory disease that affects one or more organs** but most commonly affects the lungs and lymph glands. As a result of the inflammation, abnormal lumps or nodules (called granulomas) form in one or more organs of the body.

www.medicalcoupons.in


248. Identify the structure being pointed by the Arrow?



a) Hepatic artery
b) Portal vein
c) Hepatic vein
d) Inferior venacava

The portal vein (PV) is the main vessel of the portal venous system (PVS), which **drains the blood from the gastrointestinal tract, gallbladder, pancreas, and spleen to the liver**.

249. A patient met with a RTA, is unconscious and the CT scan does not show any findings except for few petechial hemorrhages, what is the probable diagnosis?



www.medicalcoupons.in

a) Diffuse Axonal injury
b) Concussion
c) Contusion
d) Subarachnoid hemorrhage

250. A Patient with prostate cancer receives sequential radiotherapy at a dose of 1.8-2.4 Gy for 5 days every week, for 5-7 weeks this is called as?

a) Regular fractionated radiotherapy
b) Hypofractionated RT
c) Hyperfractionated RT
d) Accelerated RT

Regular fractionated radiotherapy -delivers **1.8-2 Gy single fraction per day, 5 times weekly**. • Accelerated fractionation – aims to shorten the treatment time by delivering larger than standard size fractions five times weekly or more than five fractions per week of 2 Gy. Multiple fractions may also be given daily.

251. A 35 yrs old patient from Bihar comes with generalized bone pain and was diagnosed to have Fluorosis. Fluorosis of bone is commonly associated with?



a) Osteosclerosis
b) Osteomalacia
c) Osteopetrosis
d) Osteochondrosis

Exposure to very high fluoride over a prolonged period results in acute to chronic skeletal fluorosis Vague, diffuse aches, muscle weakness, chronic fatigue and stiffness of joints with decreased range of motion are common initial symptoms. These symptoms may be dismissed as functional but may, in fact, be early signs of fluoride damage to tendinous insertions and ligaments as well as joint capsules. During later stages, calcification of the bones takes place, osteoporosis in long bones and symptoms of osteosclerosis where the bones become denser and develop abnormal crystalline structure develop.

www.medicalcoupons.in

252. Patient with Right Hypochondrial pain, Water Lily Sign is seen on Ultrasound, what is the diagnosis?

a) Entamoeba
b) Schistosoma
c) Ascariasis
d) Hydatid cyst

Cystic echinococcosis (CE), also known as hydatid disease, is **caused by infection with the larval stage of Echinococcus granulosus**, a ~2–7 millimetre-long tapeworm found in dogs (definitive host) and sheep, cattle, goats, and pigs. The water-lily sign, also known as the camalote sign, is seen in hydatid infections when **there is detachment of the endocyst membrane** which results in floating membranes within the pericyst that mimic the appearance of a water lily

253. A 2-year-old child with sudden onset breathlessness, Wheezing and Decreased breath sounds in right lung, Chest Radiograph shows diffuse opacity in right hemithorax, what is probable diagnosis?

a) Bronchial asthma
b) Pneumonia
c) Foreign body aspiration
d) Pneumothorax

Aspirated foreign bodies most commonly are lodged in **the right main stem and lower lobe**. Aspiration has been documented in all lobes, including the upper lobes, though with less frequency

254. Prophylactic craniospinal irradiation is done for which cancer?

a) Seminoma
b) Small cell lung ca
c) Prostate cancer
d) Hodgkin's lymphoma

Prophylactic cranial irradiation is now known to **improve survival** to a significant degree in small-cell lung cancer (SCLC) patients; this is in addition to its established role in preventing the disabling symptoms of brain metastases.

255. A Patient with Brain Metastasis had to undergo Head and Neck Irradiation, what is the most common delayed symptom with Head and Neck Irradiation?


a) Xerostomia
b) Dysphagia
c) Dysgeusia

www.medicalcoupons.in

d) Anhidrosis

The most common long-term complication of radiation therapy (RT) and chemoradiotherapy for head and neck cancer is **xerostomia**, which is the result of damage to the salivary glands. The magnitude of this damage is dose dependent

256. Identify the given X-RAY?




a) Hydropneumothorax
b) Pneumothorax
c) Pneumo-mediastinum
d) Lower lobe pneumonia

Hydropneumothorax is **an abnormal presence of air and fluid in the pleural space**. Even though the knowledge of hydro-pneumothorax dates back to the days of ancient Greece, not many national or international literatures are documented.

257. A Patient with tingling, numbness and paresthesias in his lower extremities, suggestive of nerve root Compression. From the given radiological image which disc is likely to be herniated?

www.medicalcoupons.in



a) L1
b) L2
c) L3
d) L4

258. A 35 years old female known case of HIV, with CD4<300, presents with fever, dyspnea, cough, and x-ray has been taken shows Causative Organism is?




a) Staphylococcus
b) Streptococcus
c) Histoplasmosis
d) Miliary TB

Miliary TB is **a potentially fatal form of TB** that results from massive lymphohematogenous dissemination of Mycobacterium tuberculosis bacilli. The epidemiology of miliary TB has been altered by the emergence of the human immunodeficiency virus (HIV) infection and widespread use of immunosuppressive drugs

www.medicalcoupons.in


259. Identify the organism in the diagram?



a) Microsporium gypseum.
b) Microsporium canis
c) Trichophyton Epidermidis.
d) Trichophyton tonsurans

Microsporium canis is a pathogenic, asexual fungus in the phylum Ascomycota that infects the upper, dead layers of skin of domesticated cats, and occasionally dogs and humans. The species has a worldwide distribution

260. A 60-year-old woman, a known case of lung cancer complains of shortness of breath. On examination her neck veins are engorged which showed a gross elevation of JVP, heart sounds are distant and breath sounds appear reduced on the left side infra scapular area. Requisition for CXR is done and is shown below. Comment on the diagnosis of the patient.



a) Congestive heart failure
b) Bilateral pleural effusion
c) Cardiac tamponade
d) Superior venacava syndrome

www.medicalcoupons.in

Cardiac tamponade is a **serious medical condition in which blood or fluids fill the space between the sac** that encases the heart and the heart muscle. This places extreme pressure on your heart. The pressure prevents the heart's ventricles from expanding fully and keeps your heart from functioning properly. Conventional radiographs of the chest in patients with cardiac tamponade may demonstrate an **enlarged cardiac silhouette** with or without an epicardial fat pad sign suggesting a pericardial effusion

261. A boy fell down while playing and the nose is deviated but septum was found normal, what is the next step done?




a) Lower the edema then closed reduction after 7 days
b) Closed reduction and automatically swelling decreases
c) Septorhinoplasty
d) Open reduction

Initially medications given to treat the edema and close reduction done after 7 days.

www.medicalcoupons.in

262. Where is the marked structure placed in the heart to activate the device present here?



a) AVN
b) SAN
c) Bundle of His
d) Purkinje fibres

263. A 15-year-old boy playing football has an injury to the leg but now complains of deep bone pain and high grade fever. A x ray was taken. What is the diagnosis?



www.medicalcoupons.in

a) Osteochondroma
b) Ewings sarcoma
c) Osteosarcoma
d) Osteochondroma

Typically, radiographic appearance of Ewing's sarcoma is that of a permeative destruction of bone. It is associated with a soft tissue mass. Lesions can **cause a lamination type of periosteal reaction** (onion skinning) or vertical spiculation

264. A 43-year-old male returned from abroad and after a week develops breathing difficulty with loss of smell and taste. Which is the next best line of investigation?

a) Sputum examination
b) Nasopharyngeal swab
c) CT chest
d) Bone marrow biopsy

The patient shows the symptoms of SARS COVID-19, the next appropriate step for the investigation would be nasopharyngeal swab

265. All the following measures are useful against COVID 19 except?

a) 2% Glutaraldehyde
b) 70% ethanol
c) Face mask
d) Hand wash

Hand wash, Face mask, 70% ethanol are all effective against the spread of COVID-19.
2% glutaraldehyde is the **disinfectant used for hospital instruments**

www.medicalcoupons.in


266. Identify the image given below?



a) Meter Dose inhaler
b) Capsule inhaler
c) Rotahaler
d) Dry powder inhaler

Rotahaler is a **plastic inhalation device which is breath activated**. This means when you inhale, the Rotahaler releases medication from the Rotacap. When inhaled correctly, the medication has a better chance to reach the small airways. This increases the medication's effectiveness.

267. In FAST which of the following needs immediate laparo-exploration?



a) 1 and 2
b) 2 and 3
c) 1 and 3
d) All

1 and 3 are the regions which immediate exploration is needed in FAST

www.medicalcoupons.in

DEC 2020
ANATOMY

1. What is the arrow marked structure in the following image?




A. Thalamus
B. Caudate nucleus
C. Lentiform nucleus
D. Claustrum

Answer: C

Explanation:

This is the transverse section of cerebrum




www.medicalcoupons.in

2. Which of the following nerve is involved in the given condition?



A. Dorsal Scapular Nerve
B. Long thoracic Nerve
C. Suprascapular Nerve
Subscapular Nerve Answer: B
Explanation: Winging of Scapula

- Medial border of scapula becomes more prominent
- Injury to long thoracic nerve from nerve root C5, C6, C7 supplying to serratus anterior muscle,
- Injury causes the condition called winging of scapula




www.medicalcoupons.in


3. Nerve involved in injury to the medial Epicondyle of Humerus?

A. Axillary nerve
B. Ulnar nerve
C. Median nerve
D. Radial nerve

Answer: B

Explanation: from the following images of nerves relate[illegible] humerus, we can clearly see that the ulnar nerve passes through the medial epicondyle.





4. Muscle attached to the arrow marked structure?




A. Gluteus maximus
B. Gluteus medius
C. Gluteus minimus
D. Iliopsoas

Answer: D

Explanation: the muscle attached to lesser trochanter is Iliopsoas whereas in the greater trochanter , Gluteus medius

www.medicalcoupons.in



5. Identify the type of joint that is marked by the arrow?




www.medicalcoupons.in

A. Syndesmosis
B. Primary Cartilaginous joint
C. Secondary Cartilaginous joint
D. Synchondrosis

Answer: C

Explanation: Secondary Cartilaginous joint are present in the midline of the body except Symphysis Menti




6. Which of the tarsal bone is marked by arrow in the image?



A. Cuboid
B. Talus
C. Navicular
D. Cuneiform


www.medicalcoupons.in

Answer: C
Explanation:




7. Which of the following nerve is involved in eversion of foot and loss of sensation in dorsal aspect of foot?

A. Deep peroneal nerve
B. Superficial peroneal nerve
C. Sural nerve
D. Saphenous nerve
Answer: B
Explanation:



www.medicalcoupons.in


8. Where does the structure marked in the following image ends?



A. Junction of left jugular vein and subclavian vein
B. Brachiocephalic vein
C. Superior vena cava
D. Right subclavian vein


Answer: A

Explanation: the structure marked is the thoracic duct. And it opens at the junction of left jugular vein and left subclavian vein.



www.medicalcoupons.in


9. ‘X’ marked tendon in the given image is?



A. Flexor digitorum superficialis
B. Flexor digitorum profundus
C. Flexor carpi radialis
D. Flexor carpi ulnaris


Answer: C

Explanation: the tendon marked is in lateral side or radial side and it is Flexor carpi ulnaris



www.medicalcoupons.in

10. Nerve passing behind the marked structure…





A. Radial nerve
B. Ulnar nerve
C. Median nerve
D. Musculocutaneous nerve

Answer: B

Explanation:


from the following images of nerves related to humerus, we can clearly see that the ulnar nerve passes through the medial epicondyle.





www.medicalcoupons.in

11. Tingling sensation and paresthesia on medial side of hand. Which structure most likely to be affected?




A. A
B. B
C. C
D. D

Answer: A

Explanation:

- A is Brachial plexus that is marked here
- B is Subclavian artery and vein
- C is Trachea
- D is Clavicle



www.medicalcoupons.in

BIO CHEM

1. Enzyme deficiency in von gierke's disease?

   (a) Glucose-6-phosphatase
   (b) Branching enzyme
   (c) Acid maltase
   (d) Glycogen Phosphorylase

   Ans. (a) Glucose-6-phosphatase

SIGNS AND SYMPTOMS of Glycogen storage disease type 1-
Glycogen accumulation in liver and renal tubule cells(hepatomegaly & renomegaly)
Hypoglycemia
Lactic acidemia
Ketosis
Hyperlipemia

Disease	Enzyme Deficient
Carbohydrates	
Type I/Von Gierke's Disease	Glucose – 6-phosphatase
Type II/Pompe's Disease	Acid Maltase
Type III/Cori's Disease	Debranching Enzyme
Type IV / Anderson's Disease	Branching Enzyme
Type V/Mc Arlde's Disease	Muscle Glycogen Phosphorylase
Type VI/ Her's Disease	Hepatic Glycogen Phosphorylase
Essential Fructosuria	Fructokinase
Fructose Intolerance	Aldolase B
Galactossemia	Galactose-1-phosphatase uridyl transferase

2. Patient presents with crying of skin and appears as shown below. Identify the vitamin deficiency?

   (a) Biotin
   (b) Niacin
   (c) Riboflavin
   (d) Ascorbic acid



www.medicalcoupons.in

Ans. (b) Niacin deficiency

Niacin (vitamin B3) deficiency results in a condition known as **pellagra**. Pellagra includes the triad of dermatitis, dementia, and diarrhea and can result in death. Niacin deficiency can occur through genetic disorders, malabsorptive conditions, and interaction with certain medications.





3. Complex IV of ETC is inhibited by?

(a) Oligomycin
(b) Malonate
(c) Cyanide
(d) Rotenone
Ans. (c) Cyanide

Complex IV, ***Cytochrome Oxidase***:
Components: Cu
Inhibitors: Cyanide, Carbon mono-oxide, H_2S

Complex I, ***NADH-Q Reductase***:
Components: NADH DH, FMN (flavin mononucleotide)
Inhibitor: Rotenone (Rat Poison and Insecticide)

www.medicalcoupons.in

<u>Complex II</u>, ***Succinate-Coenzyme Q Reductase***:
Components: succinate dehydrogenase, FAD
Inhibitors: Malonate

<u>Complex III</u>, ***Cytochrome Reductase***:
Components: b cytochromes, Fe-S-protein center, cytochrome c_1
Inhibitor: Antimycin A (Antibiotic)

4. A 40-year-old man presented with black pigmentation of pinna and sclera of eyes. The color of urine changes to black on long standing. What is the diagnosis?

   (a) Alkaptonuria
   (b) Phenylketonuria
   (c) Maple syrup urine disease
   (d) Hartnup's disease

   Ans. (a) Alkaptonuria

Alkaptonuria is a **rare genetic metabolic disorder characterized by the accumulation of homogentisic acid in the body**. Affected individuals lack enough functional levels of an enzyme required to breakdown homogentisic acid. Affected individuals may have dark urine or urine that turns black when exposed to air.



www.medicalcoupons.in

5. Which of the following is not a micromineral?

(a) Zinc
(b) Copper
(c) Calcium
(d) Iodine

Ans. (c) Calcium

Table: 10 Classification of essential minerals

Macro minerals (> 100 mg/day)*	Micro minerals (<100 mg/day)*
Calcium	Iron
Phosphorus	Zinc
Magnesium	Copper
Sulfur	Iodine
Sodium*	Fluoride
Potassium*	Manganese
Chloride*	Selenium
	Chromium
	Molybdenum

6. Nitrogenous waste is excreted from the body in the form of?

(a) Glutamine
(b) Urea
(c) Uric acid
(d) Nitric oxide

Ans. (b) Urea

Nitrogenous waste is excreted from the body in the form of **uric acid**. The nitrogenous waste that is generated in the body tends to produce toxic ammonia if remained in the body it must be excreted.
Urea and uric acid are the most common nitrogenous waste products in terrestrial animals; freshwater fish excrete ammonia and marine fish excrete both urea and trimethylamine oxide.

www.medicalcoupons.in

FORENSIC

1. Patient's relatives are complaining about patient's death and filed a case against the doctor. Which IPC section do you think will be appropriate to punish the medical practitioner?

   (a) IPC 304 A
   (b) IPC 304 B
   (c) IPC 300
   (d) IPC 302

   Ans. (a) IPC 304 A

Causing death by negligence. --Whoever causes the death of any person by doing any rash or negligent act not amounting to culpable homicide, shall be punished with imprisonment of either description for a term which may extend to two years, or with fine, or with both.

2. Active principle of the Image shown below ?

   (a) Datura
   (b) Hyoscine
   (c) Thebaine
   (d) Tetrahydro-cannabinol



Ans. (b) Hyoscine (Image → Datura)

Datura stramonium, known by the common names **thorn apple**, **jimsonweed** (**jimson weed**), **devil's snare**, or **devil's trumpet**, is a species of flowering plant in the nightshade family Solanaceae.
The active principles are alkaloids such as **hyoscine (scopolamine), hyoscyamine, and atropine**.

www.medicalcoupons.in

3. A female who died within 3 years of marriage &this was the case of suspected dowry. Who should perform the inquest in this case?

(a) Magistrate
(b) Police
(c) Village headman
(d) Judge

Ans. (a) Magistrate

Under section 176 of Criminal Procedure Code, 1973, Magistrate inquest is done in the following cases: **Dowry Death**. **Rape under police custody or government custody**, Death in Jail. Death as a result of Public Shooting.

4. A dead fetus was found in water. The crown heel length was found to be 25 cm. Light hairs were present on the skull with ossification center. Approximate age of fetus is –

(a) 5 months
(b) 7 months
(c) 9 months
(e) 3 months

Ans. (b) 5 months

RULE OF HASSE

- ✓ Rule of hasse is used to determine the age of the foetus from crown to heel length in cm
- ✓ During first five months , length in cm = (age in months) ^{2}
- ✓ After age of 5 months, Age in months = (length in cm)/5

Crown-rump length (CRL) is **the measurement of the length of human embryos and fetuses from the top of the head (crown) to the bottom of the buttocks (rump)**. It is typically determined from ultrasound imagery and can be used to estimate gestational age

5. A man was found to be agitated and shouting at the airport. Later on he fell down. On further checking, his bag was found to have enema apparatus and laxatives. All of the following are true regarding this case except?



(a) Abdominal scan is Indicated
(b) The person is malingering
(c) He is at a risk of poisoning
(d) It's a case of body packer's syndrome

www.medicalcoupons.in

Ans. (b) The person is malingering

A body packer is someone who carries drugs such as heroine or cocaine, packed in rubber or plastic, in his/her body in order to smuggle them. These people can present with symptoms that vary from mild abdominal complaints to respiratory insufficiency and even death. Physical examination and additional radiology tests are helpful for the diagnosis. Any packages can usually be seen on a plain abdominal X-ray.

6. A person was found dead in bushes with tied arms and legs, cyanosis of the nails, scratches over face, neck and chin, laceration near lips was noticed. Which of the following is not the probable mode of death in this case?

   (a) Throttling
   (b) It's a Homicide
   (c) Asphyxia
   (d) Case of a suicide

Ans. (d) Case of a suicide

Suicide is death caused by injuring oneself with the intent to die. A suicide attempt is when someone harms themselves with any intent to end their life

Homicide is an act of a human killing another person.[1] A homicide requires only a volitional act that causes the death of another, and thus a homicide may result from accidental, reckless, or negligent acts even if there is no intent to cause harm

7. A gunshot case brought for autopsy shows an entry wound with inverted margins & surrounded by blackening and tattooing. Burning & singeing of hairs is noted. What is the range of firearm weapon?

   (a) Near
   (b) About 2 meters
   (c) Contact
   (d) Distant

Ans. (a) Near

Depending on the distance of fire, the wound can also classify as contact wounds – again divided into firm-contact and loose contact – near-contact or close-range


wounds, mid-range or intermediate-range wounds, distant or far-range wounds, and indeterminate wounds.

Contact wounds are self-explanatory and are diagnosable by the presence of muzzle-imprint on the skin. **Near-contact or close-range wounds are identifiable by the presence of flame burns and singeing of hair**. Mid-range or intermediate-range wounds do not have flame burns or singeing of hair but will show the presence of smoke, gas, and unburnt particles. Distant or far-range wounds will be embedded with any accompanying components other than the lubricant forming a grease collar. Indiscriminate wounds are atypical wounds beyond the range of the weapon with an irregular shape due to yawing (changing in the axis and direction of flight of the projectile).

8. A piece of cloth was obtained from a crime scene with some spots. On examination, yellow colored needle shaped crystals were found. Which body secretions it can be?

   (a) Serum
   (b) Semen
   (c) Saliva
   (d) CSF

Ans. (b) Semen



Barberio test was invented by Barberio in the year 1905. When the **questioned stain is allowed to react with picric acid it leads to the formation of yellow needle shaped spermine picrate crystals**, including the presence of seminal stain.

9. Which of the following produces catecholamine storm as seen with scorpion sting?
   (a) Spider
   (b) Tick
   (c) Sea Urchin
   (d) Ciguatera

www.medicalcoupons.in

Ans. (a) Spider

Severe scorpion envenoming causes an autonomic storm producing multi-system organ-failure (MSOF) and death
Most of the offensive and defensive zootoxins that organisms have evolved for shutting down the nervous systems of predators and prey e.g., the venoms produced by spiders, scorpions, snakes, fish, bees, sea snails and others work by plugging ion channel pores.

Scorpion sting

- More than > 10 Species are present
- Indian red scorpion is highly venomous
- Scientific name → Mesobuthus tumulus
- Nature of venom → resembles snake venom

But fatality is less as the amount is less.

Action:

- Scorpion venom acts on Na^+ & K Channels
  ↓
  It Stimulates Sympathetic & Parasympathetic system
  ↓
  Uncontrolled release of Catecholemines into the circulations

(Autonomic Storm)

1. Pain → the tag sign
2. Paresthesia
3. Systemic Symptoms → Vomiting, Sweating, Salivation, chest pain, anxiety, cardiac arrhythmias

Rx:-

1. Immobilise
2. Pain relief
3. Prazosin
4. Scorpion antivenom (If available)

10. Dead body of a 12yr old child was found. Which bone would be best for gender assessment?

(a) Skull
(b) Femur
(c) Pelvis
(d) Mandible

www.medicalcoupons.in

Ans. (c) Pelvis

Krogman's Table

%age of accuracy in determination of sex from bone.

Pelvis	95%
Skull	90-92%
Long Bones	80%
Pelvis + Skull	98%
All 206 bones	**100%**

- **Best bone for determination of sex is *Pelvis***

11.A case of RTA was brought for autopsy with stiffness of neck, hands, both upper and lower limbs. What is this condition called?

(a) Rigor mortis
(b) Cadaveric spasm
(c) Heat stiffness
(d) Cold stiffness

Ans. (a) Rigor mortis

Rigor mortis: Literally, **the stiffness of death**. The rigidity of a body after death. ... Rigor mortis is due to a biochemical change in the muscles that occurs several hours after death, though the time of its onset after death depends on the ambient temperature

12. A woman tried to fake the pregnancy & delivery. After sometime she brought a baby claiming it to be her neighbor's child and trying to blackmail him for getting benefit in property. Such kind of child is called as?

(a) Suppositious child
(b) Spurious
(c) Superfecundation
(d) Superfetation

Ans. (a) Suppositious child

Suppositious children are **fraudulent offspring**. These arose when an heir was required and so a suitable baby might be procured and passed off as genuine.

www.medicalcoupons.in

PATHO

1. A 55-year-old diabetic patient has chronic renal failure and is undergoing hemodialysis. What is the associated amyloidosis?

   (a) AL
   (b) AA
   (c) Beta 2 microglobulin
   (d) ATTR

   Ans. Beta 2 micro globulin

Beta-2–microglobulin (beta-2m) amyloidosis is a disabling condition that affects patients undergoing long-term hemodialysis (HD) or continuous ambulatory peritoneal dialysis (CAPD).

2. A person develops pruritic rashes every time on eating sea food within one hour. Which type of hypersensitivity is this?

   (a) I
   (b) II
   (c) III
   (d) IV

   Ans. (a) I

Type I hypersensitivity is also known as an **immediate reaction** and involves immunoglobulin E (IgE) mediated release of antibodies against the soluble antigen. This results in mast cell degranulation and release of histamine and other inflammatory mediators.
Examples include **anaphylaxis and allergic rhino conjunctivitis**.

3. A nurse got needle prick injury from a HIV positive patient while doing blood sampling. Which test would be the confirmatory test to rule out the infection?

   (a) ELISA
   (b) P24 assay
   (c) Western blot
   (d) Blood culture

   Ans. (b) P24 assay

The p24 test **identifies actual HIV virus particles in blood** (p24 is a capsid structural protein which makes up a protein 'shell' on the surface of the HIV virus).

www.medicalcoupons.in

However, the p24 test is generally only positive from about two to three weeks after infection with HIV.

4. A patient comes for organ transplantation. He has a twin brother who is a match donor for
him. Which type of grafting it would be considered as?

   (a) Isograft
   (b) Allograft
   (c) Autograft
   (d) Xenograft

Ans. (a) Isograft

An Isograft is **a graft of tissue between two individuals who are genetically identical**(i.e. monozygotic twins). ... Monozygotic twins have the same major histocompatibility complex, leading to the low instances of tissue rejection by the adaptive immune system.


5. A 83-year-old female suffered with heart disease and with cardiac atrophy. On H/E, perinuclear brown colored deposits were seen. What could be the pigment responsible for this?

   (a) Hemosiderin
   (b) Melanin
   (c) Lipofuscin
   (d) Iron

   Ans. (c) Lipofuscin

Lipofuscin (**age pigment**) is a brown-yellow, electron-dense, auto fluorescent material that accumulates progressively over time in lysosomes of post mitotic cells, such as neurons and cardiac myocytes.

6. A child presents to OPD with complaints of fatigue. On further examination anemia and hepatosplenomegaly are seen. His peripheral blood smear is shown in the given Image. What is the diagnosis?



www.medicalcoupons.in


(a) Thalassemia major
(b) PNH
(c) AOCD
(d) Sickle cell anemia

Ans. Thalassemia → Image shows Target cells


Thalassemia is **an inherited blood disorder that causes your body to have less hemoglobin than normal**. Hemoglobin enables red blood cells to carry oxygen. Thalassemia can cause anemia, leaving you fatigued.

Numerous target cells are present in this patient with hemoglobin E and beta thalassemia trait. Target cells, or **codocytes**, have an excess of cell membrane relative to cell volume. Macrocytic target cells can be seen in liver disease, and microcytic target cells may be seen in thalassemia.

7. The Image below shows origin from which of the following cells?




(a) B cells
(b) CD4 T cell
(c) NK cell
(d) CD8 T cell



www.medicalcoupons.in

Reed–Sternberg cells (also known as lacunar histiocytes for certain types) are distinctive, giant cells found with light microscopy in biopsies from individuals with Hodgkin lymphoma. They are usually derived from **B lymphocytes**, classically considered crippled germinal center B cells.

8. Identify the Image and diagnose accordingly?




(a) IDA
(b) Thalassemia major
(c) Myelodysplastic syndrome
(d) Myelofibrosis

Ans. (d) Myelofibrosis

The presence of tear drop-shaped cells may indicate: **Myelofibrosis**. **Severe iron deficiency**. **Thalassemia major**. **Cancer in the bone marrow**.

A marked increase of dacrocytes is known as dacrocytosis. These tear drop cells are found primarily in **diseases** with bone marrow fibrosis, such as: primary myelofibrosis, myelodysplastic syndromes during the late course of the disease, rare form of acute leukemias and myelophthisis caused by metastatic cancers.


9. Owl eye inclusions are seen in?



(a) HSV
(b) HHV
(c) EBV
(d) CMV

Ans. (d) CMV

CMV is a herpesvirus, has double-stranded DNA, and renders infected cells 2–4 times the size of surrounding cells. These cytomegalic cells contain an eccentrically placed intranuclear inclusion surrounded by a **clear** halo, with an “owl's-eye” appearance.



10. A 50-year-old man presented with gum bleeding. Peripheral smear shows marked leukocytosis with 70% cells showing MPO positivity. What could be the diagnosis?

(a) AML
(b) ALL
(c) CML
(d) CLL

www.medicalcoupons.in

Ans. (a) AML

11. Protein in Alzheimer's?

(a) Apo C17
(b) Tau protein
(c) Keratin
(d) Collagen

Ans. Tau protein

Alzheimer's disease is thought to be caused by the abnormal build-up of proteins in and around brain cells. One of the proteins involved is called **amyloid**, deposits of which form plaques around brain cells. The other protein is called tau, deposits of which form tangles within brain cells.

12. What is the location of lactate dehydrogenase inside the cell?

Nucleus
Cytoplasm
Mitochondria
Ribosomes

Ans. Cytoplasm

Lactate dehydrogenase (LDH) catalyzes the equilibrium reaction of pyruvate to lactate. The activity of serum LDH is due to the presence of the enzyme released from damaged organs and tissues such as liver, heart, skeletal muscle, erythrocytes, etc. because LDH is located in the cytoplasm of the cells. Therefore, the activity of LDH is useful for screening for the existence of cell injuries, estimation of damaged tissues, and evaluation of treatment of diseases.

13. A 10 year old boy presented with a swollen knee after suffering a trauma. Which of the following is the likely condition seen in this patient?

Von Willebrand disease
Factor 8 deficiency
Immune thrombocytopenia
Vitamin K deficiency

Ans. Factor 8 deficiency

Hemophilia is a disease that prevents blood from clotting properly. A clot helps stop bleeding after a cut or injury. In factor VIII deficiency (hemophilia A), **the body**

www.medicalcoupons.in

doesn't make enough factor VIII (factor 8), one of the substances the body needs to form a clot.

SPM

1. Identify the logo given in the image below:



(a) Allopathic private clinic practitioner
(b) First aid
(c) Red cross emblem
(d) Suraksha clinic

Answer: D

Explanation: the above image is the logo for Suraksha clinic. For RTI (reproductive tract infection), STI (sexual tract infection) established by NACO (National Aids Control Organization)

2. Which of the following state or Union territory in India is Rabies free:

A. Sikkim
B. Kerala
C. Jammu & Kashmir
D. Andaman & Nicobar

Answer: D

Explanation: Andaman & Nicobar Islands and Lakshadweep Island from our country are currently rabies-free. As import of animals from these are avoided and hence it was easy to eradicate rabies in these states. Remember water is an effective barrier.

3. You are a medical officer and now you have to check an ANM worker's ability to do immunization who has been given training. She is having cotton, syringe-needle and lunch box after immunization of a child. Which of the following statement is correct?

A. Cotton swab in yellow bag
B. Food waste in blue box
C. Used vial in translucent white container
D. Syringe with needle in translucent white container

www.medicalcoupons.in

Answer: A
Explanation: Soiled waste like cotton, cloth etc., goes in yellow bag



4. NITI Aayog has replaced which of the following:

A. National Board of Examinations (NBE)
B. National Institute of Health & Family Welfare (NIHFW)
C. Medical Council of India (MCI)
D. National Planning Commission (NPC)

Answer: D
Explanation: NITI (National Institute Transforming India). It is actually new name of pre-existing NPC (National Planning Commission). It was established in 1950 but in 1st January 2015 NITI Aayog was formed. Comprising of Prime Minister as the Chair Person and along with him the chief ministers of all the states, Lieutenant Governor of Union territories.

5. In a population, a screening test was used for a disease, from the image below what does B-C represent:




A= Disease onset
B= First possible detection
C= Critical point
D= Usual time of diagnosis
E= final outcome

A. Lead time
B. Screening time
C. Incubation time
D. Latent period

www.medicalcoupons.in

Answer: B
Explanation: Screening time
B-D = lead time also known as advantage gained by a screening test



6. Asha worker in rural areas is designated for how much population?

A. 3000
B. 5000
C. 1000
D. 2000

Answer: C
Explanation: ASHA (Accredited Social Health Activist). A worker under National Rural Health Mission (NRHM) 2005-2012. From 2013 called as NHM (national Health Mission)
Was designated for

- 1 per village
- 1 per 1000
- From 2012 2 per village of a population of 1000

  - She is supposed to be 25–45-year-old female
  - Resident of the same village
  - Should have passed 10th
  - Supposed to get 23 days of training in PHC given by Anganwadi worker (AWW) & Auxillary Nurse Midwife (ANM)
  - She is selected by village panchayat or Gram Sabha
  - Major impact in reduction of infant mortality rate

7. Calculate Neonatal mortality rate from the data given below. Total live births were 4000
Total death among day 0-7 :40
Total death among day 7-28 :40
Total death >than 1 month :40
Perinatal death :40

www.medicalcoupons.in

A. 10
B. 20
C. 30
D. 40
Answer: B
Explanation: A neonate is one who is having an age of 0-28 days. And is Divided into 2

i. Early Neonatal Period = 0-7 days (40 from question)
ii. Late Neonatal Period = 8-28 days (40 from question)

Therefore, neonatal period = early neonatal period + late neonatal period

Formulae for Neonatal mortality rate (NNMR) = $\frac{\text{Neonatal deaths}}{\text{Live birth}} \times 1000$

From the question = $\frac{40 + 40}{4000} \times 1000$

$= \frac{80}{4000} \times 1000$

NNMR = 20.

8. On patients in Rehabilitation, A research study was conducted where impact of addiction among young adults was assessed on their subsequent divorce. Among two groups of study subjects, divorced and non-divorced, history of alcohol/drug use was assessed. What was the type of study done?
A. RCT
B. Cohort study
C. Case Control study
D. Descriptive study
Answer: C
Explanation: here people who are divorced are cases and people who are not divorced are controls and we go in a backward direction and we ask them same question. And we can calculate the strength of association (Odds Ratio).

9. Sample registration system includes all of the following except:

A. Maternal Mortality rate
B. Infant Mortality rate
C. Birth rate
D. Death rate
E. All of the above
Answer: E
Explanation: all of the options mentioned above are being included in the sample registration system. So answer is E

10. Mortality is taken into consideration in which of the following?

www.medicalcoupons.in

A. Gross Reproductive rate (GRR)
B. Net reproduction rate (NRR)
C. Total fertility rate (TFR)
D. Completed family size


Answer: B
Explanation:

- Total fertility rate (TFR) = total no of children born to a woman in her entire reproductive life.
- Gross Reproductive rate (GRR) = total no of girl children born to a woman in her entire reproductive life.
- Net reproduction rate (NRR) = total no of girl children born to a woman in her entire reproductive life taking in account their mortality
- Completed family size is simply a synonym for TFR.

11. For disposal of which of the following, a white bag is used?

A. Anatomical waste
B. Sharp waste
C. Glassware waste
D. Plastic waste

Answer: B
Explanation: A white container, puncture proof and sometimes translucent:



12. For a female with Vaginal discharge, STD color kit code would be?
A. Kit 1 Grey
B. Kit 2 Green
C. Kit 3 White
D. Kit 4 Blue

www.medicalcoupons.in

Answer: B

Explanation

Kit number	Syndrome	Color	Content
1	UD, ARD, Cervicitis	Grey	Tab. Aazithromycin (...) stat Tab. Cefexine 4... 1 stat
2	Vaginitis	Green	Tab. Secnidazo... 2 g (1) stat Cap. Flucona... ...omg (1) stat.
3	Genito ulcerative disease (GUD)- non herpetic	White	Benzathi... Penicill... (2... U)- 1 vial Tab. ...zith...omycin (1g) single dose
4	GUD – non herpetic For patients allergic to penicillin	Blue	Doxc...lin... 100mg BD×15days ...zithromycin 1g (single dose)
5	GUD –non herpetic	Red	Tab. Acyclovir 400m TDS for 7 days
6	Lower abdominal pain (LAP)	Y...llo...	Tab. Cefexime 400m OD stat Tab. Metronidazole BD for 14 days Deoxycycline 100m BD for 14 days
7	Inguinal Bubo	Bl...ck	Tab. Azithromycin 1g OD stat Tab. Doxcycline 100m BD for 21 days

13. The STD kit color for Male urethral Discharge:
    A. Kit 1 Grey
    B. Kit 2 Green
    C. Kit 3 white
    D. Kit 4 Blue

www.medicalcoupons.in

Answer: A
Explanation:

Kit number	Syndrome	Color	Content
1	UD, ARD, Cervicitis	Grey	Tab. Aazithromycin (1g) 1 stat Tab. Cefexine 400 mg 1 stat
2	Vaginitis	Green	Tab. Secnidazole 2g (1) stat Cap. Fluconazole 150mg (1) stat.
3	Genito ulcerative disease (GUD)- non herpetic	White	Benzathine Penicillin (2.4 MU)- 1 vial Tab. Azithromycin (1g) single dose
4	GUD – non herpetic For patients allergic to penicillin	Blue	Doxycycline 100mg BD×15days Azithromycin 1g (single dose)
5	GUD –non herpetic	Red	Tab. Acyclovir 400m TDS for 7 days
6	Lower abdominal pain (LAP)	Yellow	Tab. Cefexime 400m OD stat Tab. Metronidazole BD for 14 days Deoxycycline 100m BD for 14 days
7	Inguinal Bubo	Black	Tab. Azithromycin 1g OD stat Tab. Doxcycline 100m BD for 21 days

14. After which phase of clinical trials, a new drug is introduced in the market?
    A. Phase 1
    B. Phase 2
    C. Phase 3
    D. Phase 4

Answer: C

Explanation:

Phase 1	For safety and toxicity profile of a drug	Done in HHV (healthy human volunteers)
Phase 2	For effectiveness	Done in patients
Phase 3	For comparison trial with existing drug	Also done in patients
Phase 4	For long term, rare side effects	Done in patients

www.medicalcoupons.in

- Phase 4 is also called as Post-Marketing Surveillance and is the longest phase of a trial
- Phase 3 is RCT
- Maximum failure is reported from phase 2
- Maximum tolerated dose (MTD) of a drug is assessed in Phase 1
- Under new guidelines, before phase 1, we have a phase 0, where drug is tested in very small doses in very small number of healthy human volunteers to access for its efficacy and non-toxicity profile also known as phase 0 micro dosing.

15. Parameter used to test for confirmatory diagnosis when a Health worker nurse got an accidental prick with HIV+ needle would be?
A. ELISA
B. p24 Antigen
C. Western Blot assay
D. PCR
Answer: C
Explanation: when we get needle prick injury, we get 0.1-0.3% of transmission, it is also a big risk of transmission
-Screening is done by ELISA. It has very high sensitivity, but it lacks specificity and its main role is to detect antibodies
- if ELISA is positive then confirmatory diagnosis is done by WBA(Western Blot Assay) according to Indian Program, it is a protein based test and it a high specificity, it is based on p24 and GP41
-p24 antigen test can be used in window period before seroconversion
-Another test use din window period is HIV RNA Genomic sequencing (NAAT) Nucleic acid amplification test.

16. One student developed abdominal pain, jaundice and fever. Soon other students presented with similar symptoms, they are all from same boarding school and have been eating from the same canteen. Hostel Medical officer will investigate which of the following parameters to establish diagnosis?
A. IgM for Hepatitis A
B. IgG for Hepatitis A
C. IgM for Hepatitis B
D. IgG for Hepatitis B
Answer: A
Explanation: out of the all types of hepatitis, A & E are the ones with fecal oral transmission and A is the most common cause of hepatitis in children and adults, since we are doing immediate test so it has to be IgM antibody of hepatitis A.
-Hepatitis A is caused by Enterovirus 72 which is from Picornaviridae family. And in children it is sub-clinical. Incubation period of 15-45 days and period of infectivity is 2 weeks prior to Jaundice till 1 week after. Elisa is used as confirmatory to find the antibody.
A Patient is presented with rash and diarrhea for 2 months, from the image given below, find the deficiency

www.medicalcoupons.in

Types Of Viral Hepatitis

	Viral Hepatitis A	Viral Hepatitis B	Viral Hepatitis C	Viral Hepatitis D	Viral Hepatitis E
Agent	Hepatitis A virus (HAV); ssRNA; No envelope	Hepatitis B virus (HBV); dsDNA; envelope	Hepatitis C virus (HCV); ssRNA; envelope	Hepatitis D virus (HDV); ssRNA; envelope from HBV	Hepatitis E virus (HEV); ssRNA; no envelope
Route of Transmission	Fecal-oral	Parenteral, Vertical, Sexual.	Parenteral	Parenteral	Fecal-oral
Age affected	Children	Any age	Adults	Any age	Young adults
Carrier state	Nil	Common	Present	Nil (only with HBV)	Nil
Incubation period	10-50 days (avg. 25-30)	50-180 days (avg. 60-90)	40-120 days	2-12 weeks	2-9 weeks
Chronic infection	No	Yes	Yes	Yes	No
Specific Prophylaxis	Ig and Vaccine	Ig and Vaccine	Nil	HBV vaccine	

17. A Patient is presented with rash and diarrhea for 2 months, from the image given below, find the deficiency?



A. Vitamin B12 deficiency
B. Vitamin C deficiency
C. Niacin deficiency
D. Vitamin D deficiency

Answer:

Explanation: patient is having dermatitis, diarrhea for 2 months, the patient has appearance of CASAL's Necklace. So, we conclude that it is Pellagra.

Pellagra is characterized by 4D's

- Diarrhea
- Dermatitis
- Dementia
- Death

Pellagra is caused due to deficiency of vitamin B3 or Niacin. Frequently seen in Maize/Jowar eaters due to deficiency of Tryptophan which leads to pellagra due to deficiency of niacin. Excess of leucine inhibits the conversion of tryptophan to niacin

www.medicalcoupons.in

(conversion ratio of tryptophan to niacin is 60:1) in the body. Here the main culprit is Leucine.


18. Minimum thickness of Lead Apron required is:
    A. 0.1 mm
    B. 0.2 mm
    C. 0.3 mm
    D. 0.5 mm
    Answer: D
    Explanation:
    - 0.25 mm used for minimal procedures like mammography
    - 0.50 mm is most widely used and is recommended because it attenuates >90% of scattered radiation
    - 1.00 mm becomes very heavy

19. A lactating woman came to OPD 6 weeks after delivery for check-up and to get advice for the contraception, which of the following is not advised?
    A. Combined OCP
    B. Norplant
    C. IUCD
    D. Mini-pill
    Answer: A
    Explanation: Combined OCP suppresses lactation due to its estrogen component. So, we advise the mother to opt for mini-pill (lactation pill) and it has only progesterone in the form of LNG (Levonorgestrel).

20. A low socio-economic female worker who is 22 years-old presents within 8 hours of sexual assault on the 13th day of cycle and asked for emergency contraception. What will be the best advice for the patient?
    A. LNG 1.5 milligram
    B. Misoprostol 600 mcg stat
    C. Mifepristone 3 doses
    D. OCP from 1st day of next cycle
    Answer: A
    Explanation: 13th day of cycle. So, high chances of getting pregnant, out of the following options the best one would be LNG 1.5 milligram i.e., a progesterone only pill-POP (minipill) intraceptive emergency contraceptive used in the program.
    -misoprostol and mifepristone are used in MMA (Medical methods of abortion) utilized after pregnancy has taken place.
    - OCP from next cycle is not recommended because she will be already pregnant

21. Contraindication of the device from the following image includes:

www.medicalcoupons.in



A. Breast feeding
B. Monogamous
C. Hypertension
D. PID

Answer: D

Explanation:

Contra-Indications of IUCD:

- Absolute contraindications
  - Pregnancy
  - PID
  - Vaginal bleeding
  - Ectopic pregnancy
  - Cancer of cervix, uterus, adnexa
- Relative contraindications
  - Anemia
  - Menorrhagia
  - H/O PID
  - Uterine Malformation
  - Unmotivated Female
  - Fibroids

22. A child with an IQ level of 55, He is having which of the following?

A. Mild MR
B. Moderate MR
C. Severe MR
D. Normal MR

Answer: A

Explanation:

$$IQ = \frac{\text{Mental Age}}{\text{Chronological age}} \times 100$$

Mild MR is the most common form of MR in India and Down Syndrome is the most common cause.

- Normal IQ is > or = to70
  - Mild MR is 50-69
  - Moderate MR is 35-49

www.medicalcoupons.in

- Severe MR is 21-34
- Profound MR is < or = to 20

23. In Zika virus, the vector is?
    A. Sandfly
    B. Aedes aegypti
    C. Culex
    D. Anopheles
    Answer: B
    Explanation: Zika is transmitted by 4 ways they are
    - Aedes Mosquito.
    - Sexual route transmission.
    - Blood transmission.
    - Mother to child transmission.

24. Used to prevent Covid-19 among health workers, Relative risk for two vaccines, Relative risk for two vaccines A and B is 0.5 and 2.0 respectively. Identify the graph(s) which represent them correctly?



    A. A & B
    B. A & C
    C. B only
    D. None of the above
    Answer: A
    Explanation: RR risks of 2 vaccines were given. It is used to measure strength of association in a Cohort study. It can be
    RR > 1, Positive association
    RR = 1, No Association
    RR < 1, Inverse Association
    From the options both graphs A & B are correct


www.medicalcoupons.in

25. What is Urban Heart?
    A. Urban Health Equity Assessment and Response Tool
    B. Urban Health Evaluation assessment and Response Tool
    C. Urban Health Equity Assessment and Response Technique
    D. Urban Health Evaluation Assessment and Response Technique
    Answer: A
    Explanation: Tool used by WHO (World Health Organization) to identify and reduce health inequities in the city

Urban HEART
Urban Health Equity Assessment and Response Tool

It has 3 components:
- ❖ Sound Evidence
- ❖ Intersectoral Action
- ❖ Community Participation

It is integrated into local planning cycle



26. The agency that collects data and publishes large scale surveys continuously regarding morbidity, family planning, vital events?
    A. AIIMS
    B. Central Bureau of Health
    C. National Sample Survey
    D. Sample registration system
    Answer: C

Explanation: National Sample Survey taken up by NSSO (National Sample Survey Organization)

-It carries out large scale sample surveys that too in diverse fields
- Mainly based on Household
-started in 1950 -till date 586+ reports have been shared.


www.medicalcoupons.in

27. Prevalence is associated with which of the following?
    A. Sensitivity
    B. Specificity
    C. Positive predictive value
    D. Accuracy
    Answer: C
    Explanation: -positive predictive value (PPV) is directly proportional to Prevalence

PPV α Prevalence

-negative predictive value (NPV) is indirectly proportional to prevalence

$$NPV \alpha \frac{1}{Prevalence}$$



28. Best High-level Disinfectant is which of the following:
    A. Phenol
    B. Glutaraldehyde
    C. Lysol
    D. Cetrimide
    Answer: B
    Explanation:
    - High level disinfectants include
        - Glutaraldehyde
        - 6% H_2O_2
        - Chlorine Di-oxide
        - Peracetic acid
        - useful for
            - Vegetative Microorganisms
            - Lipid/Non-Lipid Viruses

www.medicalcoupons.in

- Mycobacteria
- Fungal Spores
- Few Bacterial Spores

- Intermediate level disinfectants also called as Tuberculocides includes
  - Sodium Hypochlorite (1:50)
  - Phenolic detergent solutions
    - Used in Mycobacteria
    - Most Viruses/ Bacteria
- Low level Disinfectants
  - Quaternary ammonia compounds
  - Sodium Hypochlorite (1:500)
  - Ethyl isopropyl Alcohol
    - Used for some viruses and bacteria

29. Newborn birth rate is 25 per 1000 population in a village with total population of 5000. What is the total number of pregnancies per year?
A. 125
B. 138
C. 145
D. 250
Answer: B
Explanation: First we calculate
Expected births for 1000 = 25
So, for 5000 = 125 births
But in question total no of pregnancies is asked
In India we have about 10% of pregnancy wastage so we add that along with 125 here

So, 125 + 12.5 = 137.5 pregnancies that is from options
138 would be the exact answer

30. Under ESI after certification by medical officer, beneficiaries get 70% wages for 3 months under?
A. Medical Benefit
B. Sickness Benefit
C. Disablement Benefit
D. Dependent's Benefit
Answer: B

www.medicalcoupons.in

Explanation:

Employees State Insurance (ESI)

- Runs under the Ministry of labour
- Union Minister of labour is the chair person
- Employer (3.25%) and Employee (0.75%) both need to contribute to avail the benefits under ESI
- Benefits
  - ✓ For income < 21,000 per/month are beneficiaries
    - Medical Care
    - Sickness benefit (70% wages for 91 days or 3 months)
    - If extended it becomes (80% of wages for 2 years)
    - Disablement benefits 90% wages depending on temporary or permanent disability
    - Dependent's benefit is also 90%

Under ESI Funeral expense 15,000 is also given

31. In a group of 200 Hypertensive patients. First group of 100 follow medical advice whereas second group of 100 follow lifestyle advice. There is test for both the groups after 3 months. Which of the following is best test for significance to compare Blood pressure levels in both the groups?
    A. Paired Student's T test
    B. Unpaired Student's T test
    C. Fischer test
    D. Chi-Square test
    Answer: B
    Explanation: Comparing Quantitative data in two groups, we use Unpaired Student's T test
    If Comparing Quantitative data in same group before and after an intervention we use Paired Student's T test

32. Which of the following is included in NPCDCS?
    A. Bronchial Asthma, Hypertension, CVD
    B. Diabetes, Hypertension, Bronchial Asthma
    C. CVD, Diabetes, Bronchial Asthma
    D. CVD, Cancer, Diabetes, Stroke

    Answer:
    Explanation: Full form **N**ational **P**rogram for prevention and control of **C**ancer **D**iabetes **C**VD's **S**troke **(NPCDCS)**
    Through formation of NCD clinics, this was established in 100+ districts in India

www.medicalcoupons.in



Now 450 districts in country have this and they have included COPD and CKD

33. A doctor is teaching an intern about knee reflex demonstration. What type of learning is this?
    A. Cognitive learning
    B. Affective learning
    C. Psychomotor learning
    D. None of the above
    Answer: C
    Explanation: skills is associated with psychomotor learning

34. Under ICDS, Intersectoral coordination is found with?
    A. NREGA
    B. NRHM
    C. NLEP
    D. NACP
    Answer: B
    Explanation: National rural health mission
    -ICDS-integrated child development services through Anganwadi
    -Maternal and child health care is the main aim
    -Beneficiaries are NRHM
    -Gives nutritional support

35. Blood smear for malaria will be collected by which of the following health worker?

    A. Multipurpose worker Female
    B. ANM
    C. ASHA
    D. Multipurpose worker Male

    Answer: D
    Explanation: Multipurpose health worker male has the main duty of collecting blood smear for malaria.

www.medicalcoupons.in

ANAESTHESIA

1. Which is the following I.V. anesthetic drug used for day care surgeries?

(a) Ketamine
(b) Propofol
(c) Thiopentone
(d) Etomidate

Ans. (b) Propofol

Explanation: Day care anesthesia is involved in small procedures that are done in one day, that is he has to be admitted, operated, recovered and discharged on the same calendar day, so the anesthetic agent should be short acting, easily metabolized and excreted and that is why we choose propofol. It is the anesthetic agent of choice for day care. It has anti-emetic and anti-pruritic properties and has not much side effects.

- Opioid of choice for day care: Remifentanil
- Inhalational agent of choice in day care: Sevoflurane

2. Which of the following drug is causes post-operative delirium and hallucination?

(a) Ketamine
(b) Fentanyl
(c) Thiopentone
(d) Halothane

Ans. (a) Ketamine

Explanation: Waking up from ketamine anesthesia, the name of the phenomenon we see in this condition is Emergence Delirium. Ketamine maintains heart rate, blood pressure, and airway reflexes so can be used in stomach full patients and in emergency during trauma

3. A patient after Gunshot injury presented to emergency department with hypovolemia. Which of the following cannulas can be used for most rapid blood transfusion?

(a) Grey
(b) Orange
(c) Pink
(d) Green


www.medicalcoupons.in



Ans. (b) Orange

Explanation:


Color	Gauge Size	External Diameter (mm)*	Length (mm)*	Water Flow Rate (mL/min)*
Orange	14G	2.1 mm	45mm	~240 mL/min
Gray	16G	1.8 mm	45mm	~180 mL/min
Green	18G	1.3 mm	32mm	~90 mL/min
Pink	20G	1.1 mm	32mm	~60 mL/min
Blue	22G	0.9 mm	25mm	~36 mL/min
Yellow	24G	0.7 mm	19mm	~20 mL/min



4. Local anesthetic agent works by which of the following mechanism?
   a. Blocking Sodium voltage gated channel
   b. Blocking of Potassium channel
   c. Opening Sodium voltage gated channel
   d. Opening of potassium channel
   e. Ans. (a) Blocking Sodium voltage gated channel
   f. Explanation:

www.medicalcoupons.in


Local anesthetics
↓
block the voltage-gated Na+ channels
↓
No entry of Na+ ions into the cell
↓
No depolarization
↓
No generation of action potential
↓
No generation and conduction of impulse to CNS
↓
Local anesthesia



5. What type of anesthesia is required for dental surgery?

   a. Field block
   b. Nerve block
   c. Infiltration
   d. All of the above

   Answer: D
   Explanation: All of the above



- All these kinds of anesthesia can be given in dental surgery
- Local anesthetic agent given just on the site is infiltrative
- Or nerve block for example inferior alveolar nerve block which is given in cleft palate

A field can also be blocked little above the level when we nee to block a bunch of nerves

www.medicalcoupons.in

PSYCHIATRY

1. A child with an IQ level of 55, He is having which of the following ?
   E. Mild MR
   F. Moderate MR
   G. Severe MR
   H. Normal MR
   Answer: A
   Explanation:
   IQ = Mental Age / Chronological age x 100
   Mild MR is the most common form of MR in India and Down Syndrome is the most common cause.
   - Normal IQ is > or = to70
     - Mild MR is 50-69
     - Moderate MR is 35-49
     - Severe MR is 21-34
     - Profound MR is < or = to 20

2. A person with anxiety has a feeling that his parents will die one day and he will have sleepless nights, which of the following disorders describes his condition?
   A. Generalized anxiety
   B. Severe depression
   C. PTSD
   D. Adjustment disorder
   Answer: A
   Explanation: Generalized anxiety disorder: in generalized anxiety disorder We have major and minor criteria for making a diagnosis of generalized anxiety disorder we need 2 major and 3 minor for 6 months. Treatment would be SSRI and behavior therapies
   Major criteria
   - Excessive worry
   - Not able to control

   Minor criteria
   - Restlessness
   - Muscle tension
   - Irritability
   - Sleep disturbance
   - Easy fatigability
   - Decreased concentration

3. A kid with normal learning skills but poor mathematical skills. His IQ is appropriate to his age. He passed the exams with poor marks. What could be the probable diagnosis?

www.medicalcoupons.in

A. Exam phobia
B. ADHD
C. Specific learning disorder
D. Asperger's syndrome
Answer: C
Explanation: in Specific learning disorder (Dyslexia)

- We have reading, writing, mathematical skills, mixed types
- For making a diagnosis of Dyslexia, IQ should be normal for the child
- H/o migration should be ruled out

4. Patient with 6 episodes of depression, hypermania, hypomania in a year with recovery phase in between. which of the following is appropriate diagnosis?
   A. Severe depression
   B. Cyclic dysthymia
   C. Rapid cyclic bipolar disorder
   D. Mania
   Answer: C
   Explanation: by definition, Patient must have at least four different episodes in last 1 year is rapid cycling bipolar disorder.

5. Patient after getting operated for appendicitis developed confusion and hallucination. His wife informed that he takes alcohol occasionally. what could be the most probable cause for his condition?
   A. Delirium
   B. Alcohol hallucination
   C. Acute transient psychotic disorder
   D. Mania
   Answer: A
   Explanation: Delirium
   - Most common organic disorder
   - Diagnosis of delirium
     -Impairment or clouding of consciousness
     -disorientation to time place and person
     -memory deficit
     -remote memory will remain intact
     -perceptual abnormalities in the form of illusion and visual hallucination
     - transient/fragmentary delusion
     -Sun downing phenomenon
     -Increased psychomotor activity
   - Confusion assessment method is used
   - Treated by etiology, low dose of haloperidol(to decrease psychomotor activity)

www.medicalcoupons.in

6. A patient having unilateral headache associated with nausea and vomiting which drug can be used to decrease headache episodes?
   A. Propranolol
   B. Alprazolam
   C. Fluoxetine
   D. None of the above
   Answer:
   Explanation: patient has Migraine
   - Unilateral, pulsating type associated with Nausea, Vomiting, photophobia, phonophobia
   - Precipitated by sun light
   - Travelling and lack of sleep, even hunger precipitates too
   - Treatment – for acute- to abort the attack- NSAids, tryptans
     -for prophylaxis- Propranolol, sodium valproate,topiramate

ORTHO

1. Which of the following is a radio sensitive tumor?
   A. Melanoma
   B. Ewing's sarcoma
   C. Osteosarcoma
   D. Hepatoma
   Answer: B
   Explanation: from the mnemonic **WELMS** we know the radio sensitive tumors are
   - **W**ilms
   - **E**wing's sarcoma
   - **L**ymphoma
   - **M**yeloma
   - **S**eminoma/Dysgerminoma

   We also can remember radio resistant tumors from the **HOMP**
   - **H**epatoma
   - **O**steosarcoma
   - **M**elanoma
   - **P**ancreatic cancers

   So, from the above explanation we can conclude from the options that Ewing's sarcoma is the radiosensitive tumor.


www.medicalcoupons.in

2. What is the following image?



A. Taylor brace splint
B. Milwaukee brace
C. Scoliosis brace
D. Von Rosen splint
Answer: A
Explanation: The given brace is Taylor Brace splint, given for immobilization of thoracic and lumbar spine and in used in traumatic or infected spine

3. Identify the condition given in the image?



A. Galeazzi fracture
B. Monteggia deformities
C. Monteggia Fracture
D. Green stick fracture

www.medicalcoupons.in

Answer: A
Explanation: this a fracture involving disruption of distal radial-ulnar joint with damage to the interosseous membrane and triangular Fibro cartilage component (TFCC). this is Galeazzi Fracture.
-Monteggia fracture is a fracture of ulnar with dislocated radial head


4. A patient comes after injury, from the following X-ray, What is the probable diagnosis?



A. Tibial Fracture
B. Patellar Fracture
C. Knee loose body
D. Femur fracture

Answer: B
Explanation: the image clearly shows fracture of Patella and the treatment is Tension Band wiring

5. Identify the condition shown in the following image?



A. Osteosarcoma
B. Chondrosarcoma
C. Chondroblastoma
D. Ewing's sarcoma

Answer: B
Explanation: - since osteosarcoma is a lytic type
- Chondroblastoma occurs in epiphysis

www.medicalcoupons.in

- Ewing's sarcoma occurs in the diaphysis
- Chondrosarcoma is the answer because it is the calcified lesion shown in the image and femur is the most common site

6. A 30-year-old patient has a history of trauma and pain in right hip with flexion, abduction and external rotation, what is the most likely diagnosis?
   A. Hip posterior dislocation
   B. Hip anterior dislocation
   C. Fracture neck femur
   D. Intertrochanteric fracture
   Answer: B
   Explanation: we conclude Hip anterior dislocation because
   - Hip anterior dislocation: From **FABER F**lexion **Ab**duction **E**xternal **R**otation is seen
   - Hip posterior dislocation: From **FADIR F**lexion **Ad**duction **I**nternal **R**otation is seen
   - Fracture of neck of femur: extracapsular and will have shortening and external rotation
   - Intertrochanteric fracture: extracapsular and will have more shortening and more External rotation

7. Patient has multiple fractures along with blue sclera, what is the most likely diagnosis?
   A. Osteogenesis Imperfecta
   B. Rickets
   C. Scurvy
   D. Paget's disease
   Answer: A
   Explanation: Osteogenesis Imperfecta
   Osteogenesis imperfecta (OI) is an inherited (genetic) bone disorder that is present at birth. It is also known as brittle bone disease. A child born with OI may have soft bones that break (fracture) easily, bones that are not formed normally, and other problems. Blue sclera is also a feature of Osteogenesis Imperfecta. It is the commonest cause of multiple fractures at birth.
8. 60-year-old female patient complaints of pain in knee bilaterally, the pain increases while going up and down the stairs. The x-ray shows Bilateral reduced joint space, what is the most likely diagnosis?
   A. Osteoarthritis
   B. Rheumatoid Arthritis
   C. Ankylosing spondylitis
   D. Hemophiliac arthropathy

   Answer: A
   Explanation: Osteoarthritis

www.medicalcoupons.in

Osteoarthritis is the most common form of arthritis, affecting millions of people worldwide. It occurs when the protective cartilage that cushions the ends of the bones wears down over time. Although osteoarthritis can damage any joint, the disorder most commonly affects joints in your hands, knees, hips and spine.

9. The following clinical image shows?




A. Winging of scapula
B. Erb's palsy
C. Klippel Feil Syndrome
D. Fracture Scapula

Answer: A

Explanation: the image clearly shows winging of scapula, The term 'winged scapula' is used when the muscles of the scapula are too weak or paralyzed, resulting in a limited ability to stabilize the scapula. As a result, the medial or lateral borders of the scapula protrudes from back, like wings.

10. A child with widening of bone has the following x-ray, what is the aetiology?



www.medicalcoupons.in

A. Scurvy
B. Rickets
C. Gout
D. Salter Harris type 2 Injury

Answer: B

Explanation: Rickets will have a wide skeleton, the bones have mineralization defect and cupping and splaying of the bone can be clearly seen and the margins point out called as flaring/fraying, classical characters of Rickets

11. Which of the following nerve is related to medial epicondyle?
    A. Median nerve
    B. Ulnar nerve
    C. AIN
    D. PIN

    Answer: B

    Explanation: Ulnar nerve
    - In upper limb around upper part of humerus= axillary nerve
    - Around the shaft =Radial nerve
    - Behind the medial epicondyle=Ulnar nerve
    - And head of radius=Posterior interosseous nerve

12. The spot diagnosis for the following image is?



    A. Scoliosis
    B. Disc prolapses
    C. Klippel Feil Syndrome
    D. Torticollis


    Answer: C

    Explanation: short neck, lower hairline, short and high scapula which is a classical feature of Klippel Feil Syndrome

www.medicalcoupons.in

SURGERY

1. In the hospital, a patient presents with the following swelling on standing which tends to reduce when he lies down. What could be the probable diagnosis?



A. Para-umbilical Hernia
B. Umbilical Hernia
C. Epigastric Hernia
D. Omphalocele

Answer: A

Explanation:

	UMBLICAL HERNIA	ARA-MBLICAL HERNIA
1	The abdominal contents bulge out through weak umbilical sc-	Ab nial on through linea alba above or el w the umblicus
2	Umblicus is everted	U nblicus becomes cresent shaped
3	The entire fundus of the ac is covered by the um, cus	nly half of the fundus is covered by umblicus and the reminder by adjacent skin
4	May be congenita /acqu d	Always acquired
5	Ascitis is an i- orta precipitati g fac or	Obesity and lax abdominal wall are factors
6	Neck of th s- is v ide	Neck of the sac is narrow
7	Con a d type can wait upto 4 yea s for s ntaneous closure	Always needs surgery

2. A 5-year-old child was brought to the clinic with the swelling shown below which keeps on increasing in size. What should be the appropriate management for this condition?



A. Intralesional steroids
B. Excision
C. Reassurance
D. Conservative

www.medicalcoupons.in

Answer: B
Explanation: A dermoid cyst is a collection of tissue under the skin. It may contain hair follicles, oil, and sweat glands. In some cases, it may contain bone, teeth, or nerves. A dermoid cyst may appear at birth or soon after. Dermoid cysts are often found on the head, neck, or face, most often around the eyes.

- Surgical Excision is the only mode of treatment for dermoid cyst.

3. A 45-year-old female presents with Rapidly enlarging lump in her breast. Firm and mobile on examination with Bosselated surface. Lymph nodes are not involved. What could be the diagnosis?



A. Fibroadenoma
B. Phyllodes tumor
C. Ductal carcinoma
D. Ductal papilloma
Answer: B
Explanation: Phyllodes tumor

- A phyllodes tumor is a rare tumor of the breast. Phyllodes tumors grow in the connective tissue of the breast, called the stroma. This includes the tissue and ligaments that surround the ducts, blood vessels, and lymph vessels in the breast.
- Most phyllodes tumors are benign. They may look very much like common benign breast tumors called fibroadenomas. Often, the pathologist needs to look at the whole tumor under the microscope to make a diagnosis. This is why surgery to remove a phyllodes tumor is recommended, even if it is thought to be benign.

4. An Incision was made over the patient's chest for a surgery and after sometime, he comes with the scar shown in the image. What is the diagnosis?



www.medicalcoupons.in

A. Keloid
B. Hypertrophic scar
C. Contracture
D. Abscess

Answer: A

Explanation: Keloid scar

Hypertrophic scars	Keloid scars
Raised, linear, firm lesion	Raised, hard dermal outgrowth
Develop within 3 months of injury	Develop up to or even beyond 1 year postinjury
Remain within the boundaries of the original wound site	Spread outside wound margins, invading normal skin
Common postburn injury	Any form of dermal injury
Increased collagen type III	High in collagen type I
Not typically associated with pigmented skin	Associated with certain ethnicities (higher in darkly pigmented skin)
Not commonly familial	Strong familial tendency
Affect any age	Common in young age
Can regress spontaneously	Do not regress spontaneously

5. After RTA, a patient presented to emergency and following is done to maintain the airway. What is the procedure shown in the image?



A. Intubation
B. Tracheostomy
C. Thoracotomy
D. Radical neck dissection

Answer: B

Explanation: Tracheostomy

6. A bed ridden patient developed fever, pain and pedal edema, what could be the diagnosis?

A. Cellulitis
B. DVT
C. DIC
D. Pressure ulcer

www.medicalcoupons.in

Answer: B
Explanation: DVT only can cause low grade fever, remaining all in the options can only cause high grade fever.

7. A Patient presented with long standing pain in the anal region. On examination the finding shown is given below. What is the most important thing to look for on per rectal examination of this patient?



A. Internal os of the fistula
B. International haemorrhoids
C. Anal Fissure
D. Fistula in ano

Answer: D
Explanation: Perianal fistula

- A perianal fistula (fistula-in-ano) refers to an abnormal connection between the anal canal and the perianal skin. The majority are associated with anorectal abscess formation, with one third of patients with an anorectal abscess having an associated perianal fistula at the time of presentation.
-


8. Which instrument is shown in the image?



www.medicalcoupons.in

A. Allis Forceps
B. Sponge holder
C. Babcock forceps
D. Retractor
Answer: C
Explanation: The image showed is Babcock forceps

9. What procedure is done in the following image?



A. Phallen's sign
B. Tinel's sign
C. Prayer sign
D. Schamroth sign
Answer: C
Explanation: A positive prayer sign can be is when on examination the patient is unable to approximate the palmar surfaces of the phalangeal joints while pressing their hands together.
Seen in diabetes

10. A woman was presented with the following lesion in the image through which purulent discharge was coming out. The most probable diagnosis?



A. Hydradenitis Suppurativa
B. Tuberculous lymphadenitis
C. Branchial cyst
D. Bezold abscess

www.medicalcoupons.in

Answer: B
Explanation: Tubercular Lymphadenitis

- Tuberculous lymphadenitis is a chronic, specific granulomatous inflammation of the lymph node with caseation necrosis, caused by infection with Mycobacterium tuberculosis or related bacteria.

11. A man came to the hospital presented with vomiting, nausea and fever. He was admitted for appendectomy after diagnosing appendicitis. During surgery, surgeons found an incidental finding as shown in the image below which was present at a distance of 2 feet from ileocaecal junction. What is the most likely diagnosis?



A. Intussusception
B. Intestinal obstruction
C. Meckel's diverticulum
D. Diverticulitis

Answer: C
Explanation: Meckel's diverticulum

- Most common congenital anomaly of small bowel
- It is remnant of Vitellointestinal duct
  - Connects between midgut and yolk sac
  - Normally disappears in 7th week of intrauterine life
- Present in 2% of population
- Lies in ileum 2 feet from ileocaecal junction
- Measures 2 inches in length
- Has a separate blood supply
- May contain ectopic gastric, pancreatic or colonic mucosa

12. Identify the procedure in the following image?



A. Colostomy
B. Loop ileostomy
C. Ascending colostomy
D. Appendectomy

www.medicalcoupons.in

Answer: B
Explanation: loop Ileostomy

- To form a loop ileostomy, a loop of small intestine is pulled out through a cut in your abdomen. This section of intestine is then opened up and stitched to the skin to form a stoma. The colon and rectum are left in place.

13. A cricket player was injured with the ball and suddenly lost consciousness. When one of the teammate comes for helping, he gets up by himself and starts playing. During tea break, the person was again unconscious while his friend saw him, immediately he has been rushed to the hospital. What is the most probable diagnosis?

A. EDH
B. SDH
C. SAH
D. Intracerebral haemorrhage
Answer: A
Explanation: EDH


- The player was in Lucid interval and it is a classical feature of EDH
- CT-scan will show Biconvex shaped hematoma



14. In which of the following is Dohlman's procedure used for?

A. Cricoid webs
B. Pharyngeal pouch
C. Achalasia Cardia
D. Diffuse esophagal spasm
Answer: B
Explanation: used in pharyngeal pouch or Zenker's Diverticulum

www.medicalcoupons.in



15. A young male a chronic smoker, after developing the following condition of finger tips as seen in the Image coms to the clinic. What could be the condition?



A. Medium vessel vasculitis
B. Buerger's disease
C. Scleroderma
D. APLA syndrome

Answer: B
Explanation: Buerger's disease

- Buerger's disease is a rare disease of the arteries and veins in the arms and legs. In Buerger's disease — also called thromboangiitis obliterans — blood vessels become inflamed, swell and can become blocked with blood clots (thrombi)
- This eventually damages or destroys skin tissues and may lead to infection and gangrene. Buerger's disease usually first shows in your hands and feet and may eventually affect larger areas of your arms and legs.
- The exact cause of Buerger's disease is unknown. While tobacco use clearly plays a role in the development of Buerger's disease, it's not clear how it does so. It's thought that chemicals in tobacco may irritate the lining of your blood vessels, causing them to swell.
-

www.medicalcoupons.in


16. A patient presented with the following condition as shown below. what could be the diagnosis?



A. Arterial ulcer
B. Venous ulcer
C. Neuropathic ulcer
D. Pressure ulcer


Answer: B
Explanation: venous ulcer

- Are seen most commonly seen in medial malleolus (Gaiter area)
- They are usually sloping ulcers and floor is covered by granulation tissue
- If it bleeds, we ask the patient to lift the limb and we do not compress

Feature	Ulcer Type			
	Venous	Arterial	Neuropathic Diabetic	Pressure
Underlying condition	Varicose veins, previous deep-vein thrombosis, obesity, pregnancy, recurrent phlebitis	Diabetes, hypertension, smoking, peripheral vascular disease	Diabetes, trauma, prolonged pressure	Limited mobility
Ulcer location	Area between the lower calf and the medial malleolus	Pressure points, toes and feet, lateral malleolus and tibial areas	Plantar aspect of foot, tip of the toe, lateral to fifth metatarsal	Bony prominences, heel
Ulcer characteristic	Shallow and flat, veins, moderate shear, exudate, rough at base with granulation tissue	Punched out and deep, irregular shape, unhealthy wound bed, presence of necrotic tissue, minimal exudate unless infected	Deep, surrounded by callus, insensate	Deep, often macerated
				
Condition of leg or foot	Hemosiderin staining, thickening and fibrosis, eczematous and itchy skin, limb edema, normal capillary refill	Thin shiny skin, reduced hair growth, cool skin, pallor on leg elevation, absent or weak pulses, delayed capillary refill, gangrene	Dry, cracked, insensate, calluses	Atrophic skin, loss of muscle mass
Treatment	Compression therapy, leg elevation, surgical management	Revascularization, antiplatelet medications, management of risk factors	Off-loading of pressure, topical growth factors	Off-loading of pressure; reduction of excessive moisture, shear, and friction; adequate nutrition

www.medicalcoupons.in

17. Identify the Anomaly shown in the picture?



A. Pyloric stenosis
B. Pancreatic Divisum
C. Annular Pancreas
D. Duodenal Atresia

Answer: C

Explanation: Annular Pancreas

- It is due to the failure of rotation of Ventral bud of pancreas.
- Annular pancreas is the most common birth defect seen in the pancreas. The pancreas is an organ that plays an important part in your digestive process. The term annular pancreas means that a ring of extra pancreatic tissue covers the first part of the small intestine (duodenum).

18. A patient has an USG finding of a gall stone abutting the cystic duct with dilatation of Common hepatic duct. What is the most likely diagnosis?


A. Acute cholecystitis
B. Porcelain GB
C. GB polyp
D. Mirizzi Syndrome

Answer: D
Explanation: Mirrizi Syndrome

- Mirizzi syndrome is defined as common hepatic duct obstruction caused by extrinsic compression from an impacted stone in the cystic duct or infundibulum of the gallbladder

www.medicalcoupons.in

MIRRIZI`s syndrome



19. Which of the following fluid is avoided in Intracranial Head injury?

    A. NS
    B. RL
    C. 5% dextrose
    D. Hypertonic solution


    Answer: C
    Explanation: The ideal fluid is Mannitol, even any hypertonic solutions can be used.
    But we cannot use any hypotonic solution or glucose containing solutions because they will cause cerebral edema causing an increase in intra-cranial pressure.

20. A male came to the OPD with an abdominal lump in periumbilical region that moves at right angle to attachment of mesentry. What is your diagnosis?

    A. Chylolymphatic cyst
    B. Pancreatic Pseudocysts
    C. Retroperitoneal Sarcoma
    D. Peritoneal cyst

    Answer: A
    Explanation: Chylolymphatic cyst

    - A chylolymphatic cyst is a rare variant of a mesenteric cyst [1,2]. These cysts present within the mesentery, lined with a thin endothelium or mesothelium and filled with chylous and lymphatic fluid.




www.medicalcoupons.in

21. A new born is found to have herniation of bowel and liver through th Umbilicus, which is covered with a membrane. What is the diagnosis?

A. Gastrochisis
B. Omphalocele
C. Epigastric Hernia
D. Ectopia Vesica

Answer: B
Explanation:



PEDIA

1. A 5-year-old child of 17 kg weight with diarrhea and unable to take the feed. What is the amount of maintenance fluid required in 24 hours?

(a) 1000 ml
(b) 1700 ml
(c) 1350 ml
(d) 2000 ml

Ans (c) 1350 ml

17 kg ----- (10 x 100) + (7 x 50) = 1000 + 350 = 1350 ml

Calculation of fluids by HOLIDAY SEGAR REGMIN
First 0-10 kg = 100ml/kg
Next 10-20 kg= +50 ml/kg
>20 kg = +20 ml/kg

TREATMENT PLAN FOR DEHYDRATION: (Diarrhea Treatment)

www.medicalcoupons.in

No Dehydration	Some Dehydration	Severe dehydration (lethargy)
PLAN A	**PLAN B**	**Plan C**
• ORS 100-200ml/loose stool (ongoing losses 10-20ml/kg)	• Rehydration (75ml/kg in 1st 4 hrs) + maintenance (replacing on going loss- 10-20ml/kg)	• Start IV fluids • Ringer lactate >>Normal saline Give 100 ml/kg as :

Age	30ml/kg	70ml/kg
<1yr	1hr	5 hr
>1yr	30 min	2 ½ hr

2. A mother with the new born is worried for her child as the newborn hasn't passed the meconium till 6 hours of birth. How much time should she wait for the meconium to pass out?

(a) 48 hours
(b) 24 hours
(c) 12 hours
(d) 72 hours

Ans. (a) 48 hours

Maximum time to wait in a newborn to pass out -

24 hours – urine
48 hours – meconium

Delay in passage of meconium:

Hirschsprung disease (Aganglionsis)
Imperforate anus
Hypothyroidism
Meconium ileus [cystic fibrosis]
Small left colon syndrome [Associated with Infant of diabetic mothers]

www.medicalcoupons.in

3. **A boy presents with hemarthrosis of left knee after a history of fall. Deficiency of which factor can lead to the following condition?**



(a) Von-willebrand disease
(b) Vitamin K deficiency
(c) Thrombocytopenia
(d) Factor VIII deficiency

Ans. (d) Factor VIII deficiency

Hemophilia is a disease that prevents blood from clotting properly. A clot helps stop bleeding after a cut or injury. In factor VIII deficiency (hemophilia A), **the body doesn't make enough** factor VIII (factor 8), one of the substances the body needs to form a clot.

Hemophilia:
X linked Recessive
Causes severe forms of bleeding
M/C joint affected: ANKLE
Characteristic lab feature: increased APTT

MEDICINE


1. A 45 year-old female having arthralgia, swallowing difficulty with dry food and gritty feeling under eyelid. She is suffering from NCNC(Normocytic normochromic anemia) with Increased ESR, positive for anti-ANA, Anti-Ro and Rheumatoid factor. Which of the following could be the diagnosis?

(a) Sjogren's syndrome
(b) Rheumatoid arthritis
(c) Serum sickness
(d) SLE

www.medicalcoupons.in

Ans. (a) Sjogren's syndrome

Sjogren's syndrome is **a disorder of your immune system identified** by its two most common symptoms — dry eyes and a dry mouth. The condition often accompanies other immune system disorders, such as rheumatoid arthritis and lupus.



2. Diagnose the given ECG?



(a) Ventricular tachycardia
(b) Ventricular fibrillation
(c) Atrial tachycardia
(d) Atrial fibrillation

Ans. (a) Ventricular tachycardia


www.medicalcoupons.in

Atrial Flutter




- **Organized** electrical activity (Atrial fibrillation is *disorganized*)
- **Atrial rate** ranges from 250-300 beats per minute
- AV node conducts every **2 or 3 atrial impulses**
- Ventricular rate is classically **150 (2:1 conduction)**
- Characteristic **"saw-tooth" pattern**

Atrial Fibrillation



- **No P waves**
- Irregularly **irregular**
- **Uncoordinated** atrial activity
- Most common **sustained dysrhythmia**
- **Variable** ventricular response rate

Ventricular Fibrillation



- Completely **disorganized**
- Immediate cessation of cardiac output and **associated pulse**
- No discernible P waves, QRS complexes, or T waves
- **Incompatible** with life


Ventricular Tachycardia



- **AV dissociation**
- QRS > 120 msec **(wide complex)**
- **Rate** >100 (typically 150-200)
- **Fusion** beats
- **Capture** beats

www.medicalcoupons.in

3. Identify the anomaly shown in the Image?



(a) Pyloric stenosis
(b) Pancreatic divisum
(c) Annular pancreas
(d) Duodenal atresia

Ans. (c) Annular Pancreas

Annular pancreas is a morphological anomaly that results in pancreatic tissue completely or incompletely encircling the duodenum. This condition can cause duodenal obstruction
he pancreas develops from a single dorsal and two ventral buds, which appear as outgrowths of primitive foregut at 5 weeks of gestation. The ventral buds fuse rapidly. In the 7th week of gestation, the duodenum expands, and rotates the ventral bud from right to left, and fuses with the dorsal bud. The ventral bud forms the inferior part of uncinate process and inferior head of pancreas while the dorsal bud gives rise to tail and body of pancreas.

Annular pancreas develops due to failure of the ventral bud to rotate with duodenum, causing encasement of duodenum.

4. A patient comes to emergency with complaints of acute abdominal pain after alcohol intake. Pain was radiating to his back. On further investigation, he was found to have raised level of AST, ALT and amylase. What could be the diagnosis?

(a) HCC
(b) Hepatitis
(c) Acute pancreatitis
(d) Gall bladder stones

Ans. (c) Acute Pancreatitis

www.medicalcoupons.in

The most common causes of acute pancreatitis are gallstones and alcohol-induced pancreatitis. This patients liver enzymes especially ALT and **bilirubin levels are elevated**


Mnemonic for Causes of Acute Pancreatitis

- **I: I**diopathic
- **G: G**allstones
- **E: E**thanol (alcohol)
- **T: T**rauma
- **S: S**teroids
- **M: M**umps (and other infections) / malignancy
- **A: A**utoimmune
- **S: S**corpion stings/Spider bites
- **H: H**yperlipidemia/Hypercalcemia
- **E: E**RCP
- **D: D**rugs

5. A patient with ptosis, anhidrosis and slight drooping of eyelid is presented to the clinic. What do you think the patient is suffering from?
   a. Horner syndrome
   b. Keratoconjunctivitis
   c. Dacryocystitis
   d. Orbital cellulitis

Ans. Horner syndrome

Horner syndrome is a **combination of signs and symptoms caused by the disruption of a nerve pathway from the brain to the face and eye** on one side of the body. Typically, Horner syndrome results in a decreased pupil size, a drooping eyelid and decreased sweating on the affected side of your face.



www.medicalcoupons.in


6. **Which fluid should not be given in severe head injury?**

(a) 5% dextrose
(b) Normal saline
(c) Hypertonic saline
(d) Mannitol

Ans. (a) 5% dextrose

Mannitol, a hypertonic crystalloid solution, is commonly used to decrease brain water content and reduce intracranial pressure (ICP). Hypertonic saline solutions also decrease brain water and ICP while temporarily increasing systolic blood pressure and cardiac output. **Hypo-osmolar solutions, such as 5% dextrose in water, reduce serum sodium and increase brain water and ICP so Hypotonic, low sodium and dextrose-containing fluids should be avoided**

7. Purple striate on the abdomen as shown in the picture given below is a characteristic feature of which of the following given condition?




(a) Cushing's syndrome
(b) Hypothyroidism
(c) SIADH
(d) Addison's disease

Ans. (a) Cushing syndrome

Cushing's syndrome is **a disorder that occurs when your body makes too much of the hormone cortisol over a long period of time**. Cortisol is sometimes called the "stress hormone" because it helps your body respond to stress.

www.medicalcoupons.in



8. A boy presented with blue sclera who has a history of frequent fractures. What could be the diagnosis?

   (a) Scurvy
   (b) Rickets
   (c) Osteogenesis Imperfecta
   (d) Osteomalacia

   Ans. (c) Osteogenesis Imperfecta


Osteogenesis Imperfecta (OI), also known as **"brittle bone disease"**, is an **autosomal dominant** bone disorder characterized by fragile bones that break easily.

Specifically, Osteogenesis Imperfecta is caused by defects in **Type I Collagen,** a key component of bone. Patients present with **recurrent fractures** (distinguish vs. child abuse), **blue sclera** (translucency with visible choroidal veins), **tooth abnormalities**, and **hearing loss** (due to ossicle involvement).

Treatment involves **bisphosphonates** (such as alendronate), which prevent the loss of bone density and help reduce the risk for fracture.

www.medicalcoupons.in

10. Identify the ECG given below?



(a) Atrial flutter
(b) Ventricular tachycardia
(c) Ventricular fibrillation
(d) MI

Ans. (a) Atrial flutter

www.medicalcoupons.in

Atrial Flutter



- **Organized** electrical activity (Atrial fibrillation is *disorganized*)
- **Atrial rate** ranges from 250-300 beats per minute
- AV node conducts every **2 or 3 atrial impulses**
- Ventricular rate is classically **150 (2:1 conduction)**
- Characteristic **"saw-tooth" pattern**

Atrial Fibrillation



- **No P waves**
- Irregularly **irregular**
- **Uncoordinated** atrial activity
- Most common **sustained dysrhythmia**
- **Variable** ventricular response rate

Ventricular Fibrillation



- Completely **disorganized**
- Immediate cessation of cardiac output, **no associated pulse**
- No discernible P waves, QRS complexes, or T waves
- **Incompatible** with life

Ventricular Tachycardia




- **AV dissociation**
- QRS >120 msec **(wide complex)**
- **Rate >100** (typically 150-200)
- **Fusion** beats
- **Capture** beats

www.medicalcoupons.in

www.medicalcoupons.in

11. Identify the condition shown below?




a. Nail pterygium
b. Clubbing
c. Koilonychia
d. None of the above

Ans. Clubbing

www.medicalcoupons.in

www.medicalcoupons.in





Causes of Nail Clubbing - Mnemonic CLUBBING
Cyanotic Heart Disease, Cystic Fibrosis
Lung Cancer, Lung abscess
Ulcerative Colitis
Bronchiectasis
Benign Mesothelioma
Infective Endocarditis, Idiopathic Pulmonary
Neurogenic Tumors
Gastrointestinal Disease

12. A 68 year-old Diabetic patient develops which of the following macro vascular complications after 20 years?


(a) Retinopathy
(b) Neuropathy
(c) Nephropathy
(d) Myocardial Infarction

www.medicalcoupons.in

Ans. (d) Myocardial Infarction

Macrovascular complications of T2DM include coronary heart disease, cardiomyopathy, arrhythmias and sudden death, cerebrovascular disease and peripheral artery disease. Cardiovascular disease is the primary cause of death in diabetic patients. Many clinical studies have shown a connection between T2DM and vascular disease, but almost always other risk factors are present in diabetic patients, such as hypertension, obesity and dyslipidemia.

13. Tracheal shift towards right with left sided pneumothorax. What is the management?



(a) Left sided thoracostomy
(b) Right sided thoracostomy
(c) Conservative
(d) Wrapping the chest tightly

Ans. (a) Left sided thoracostomy

When tracheal deviation is present, the trachea will be displaced in the direction of less pressure. Meaning, that if one side of the chest cavity has an increase in pressure (such as in the case of a pneumothorax) the trachea will shift **towards the opposing side**.
Thoracostomy is a **minimally invasive procedure** in which a thin plastic tube is inserted into the pleural space — the area between the chest wall and lungs — and may be attached to a suction device to remove excess fluid or air. A chest tube may also be used to deliver medications into the pleural space.

www.medicalcoupons.in

14. A HIV patient presents with following presentation. What is the staging of HIV





(a) I
(b) II
(c) III
(d) IV

Ans. (c) III

STAGE I	STAGE 2	STAGE 3	STAGE 4
Asymptomatic	Moderate unexpected weight loss	Unexpected severe weight loss	HIV wasting syndrome
Persistent generalized lymphadenopathy	Recurrent respiratory tract infection	Persistent oral candidisis	Pneumocystic pneumonia
	Herpes Zoster	Pulmonary tuberculosis	Karposi sarcoma
	Angular cheilitis		Extra pulmonary tuberculosis
	Fungal nail infection		

15. JVP Image shows deep x descent and a rapid y descent and elevated over all. What could be the diagnosis?






www.medicalcoupons.in

(a) Constrictive pericarditis
(b) Dilated cardiomyopathy
(c) HOCM
(d) Tamponade

Ans. (a) Constrictive pericarditis

Constrictive pericarditis

- JVP is elevated
- a wave is usually normal
- v wave is usually equal to a wave
- x descent –prominent
- y descent – rapid descent
- Kussmauls sign is usually positive



16. A student who was eating the food of his hostel mess found to have developed hepatomegaly, jaundice. On the same time all other students who were eating from the same mess developed similar symptoms. Which investigation is more appropriate to be done to find out the cause?

(a) IgM for Hepatitis A
(b) IgG for Hepatitis A
(c) IgM for Hepatitis B
(d) IgG for Hepatitis B

Ans. (a) IgM for Hepatitis A

Hepatitis A immunoglobulin M (IgM anti-HAV) antibody test: When a person is first infected with hepatitis A, the body produces IgM anti-HAV antibodies. These antibodies are usually detectable from two weeks after symptoms begin to around six months later.

Hepatomegaly is an enlarged liver, which means it's swollen beyond its usual size. A swollen liver usually is a symptom of another health condition, such as hepatitis.

Depending on the cause of your enlarged liver, you may notice symptoms like:

- Yellowing of the skin or eyes (jaundice)
- Fatigue and weakness
- Nausea
- Weight loss

www.medicalcoupons.in

17. A male patient presented with high grade fever and headache. He was found to have protein level 220 mg/dl and glucose 35mg/dl in CSF. He develops rashes as shown below. What should be the probable cause?



(a) E coli
(b) Listeria
(c) Neisseria meningitides
(d) Cryptococcus

Ans. (c) Neisseria meningitides

Typical CSF abnormalities in meningitis include the following:

- Increased opening pressure (>180 mm water)
- Pleocytosis of polymorphonuclear leukocytes (white blood cell [WBC] counts between 10 and 10,000 cells/µL, predominantly neutrophils)
- Decreased glucose concentration (< 45 mg/dL)
- Increased protein concentration (>45 mg/dL)
- The bacteria rapidly multiply and produce toxins which travel around the body causing damage to blood vessels and organs. As the blood vessels get damaged, **blood starts to 'leak' into the surrounding tissue**, often causing what looks like a 'rash' to appear on the skin.

18. A patient who is normal at rest but has discomfort on activities. Normal activities are slightly affected. The patient comes under which category of NYHA classification?

(a) NYHA I
(b) NYHA II
(c) NYHA III
(d) NYHA IV

www.medicalcoupons.in

Ans. (b) NYHA II

Functional Capacity	Objective Assessment
Class I. Patients with cardiac disease but without resulting limitation of physical activity. Ordinary physical activity does not cause undue fatigue, palpitation, dyspnea, or anginal pain.	**A.** No objective evidence of cardiovascular disease.
Class II. Patients with cardiac disease resulting in slight limitation of physical activity. They are comfortable at rest. Ordinary physical activity results in fatigue, palpitation, dyspnea, or anginal pain.	**B.** Objective evidence of minimal cardiovascular disease.
Class III. Patients with cardiac disease resulting in marked limitation of physical activity. They are comfortable at rest. Less than ordinary activity causes fatigue, palpitation, dyspnea, or anginal pain.	**C.** Objective evidence of moderately severe cardiovascular disease.
Class IV. Patients with cardiac disease resulting in inability to carry on any physical activity without discomfort. Symptoms of heart failure or the anginal syndrome may be present even at rest. If any physical activity is undertaken, discomfort is increased.	**D.** Objective evidence of severe cardiovascular disease.

19. A patient comes to emergency with fever and headache. O/E he has neck stiffness, CSF analysis was done and Opening pressure increased, proteins mild increased, glucose normal, lymphocytes increased. What is the most likely diagnosis?

(a) Tubercular meningitis
(b) Purulent bacterial meningitis
(c) Viral meningitis
(d) Cryptococcus

Ans. (c) Viral meningitis

	Opening Pressure (mm Hg)	Cell Count (WBC Cells/microL)	Protein (mg/dL)	Glucose (mg/dL)
Normal	**8 - 15** (100 - 180 mm of H_2O) *with patient lying on their side*	**0 - 5**	**< 40**	**40 - 70**
Bacterial	↑	↑ **PMNs**	↑	↓
Viral	↑ or **normal**	↑ **Lymphocytes**	↑ or **normal**	**normal**
Fungal/TB	↑	↑ **Lymphocytes**	↑	↓

MEDZCOOL

www.medicalcoupons.in

20. A woman after hysterectomy presents with distended abdomen. What is electrolyte imbalance seen?

(a) Hyperkalemia
(b) Hypokalemia
(c) Hypernatremia
(d) Hyponatremia

Ans. (b) Hypokalemia

The symptoms of **hypokalemia** are nonspecific and predominantly are related to muscular or cardiac function. Weakness and fatigue are the most common complaints. The muscular weakness that occurs with hypokalemia can manifest in protean ways (eg, dyspnea, constipation or abdominal distention, exercise intolerance).

21. Patient with previous history of MI 3 years ago now presenting with tachypnea, tachycardia, S3 heard. What could be the diagnosis?

(a) MI
(b) Pericarditis
(c) Congestive heart failure
(d) Pulmonary embolism

Ans. (c) Congestive heart failure

Results **from increased atrial pressure leading to increased flow rates**, as seen in congestive heart failure, which is the most common cause of a S3.

Risk factors for heart failure include:

- Coronary artery disease. Narrowed arteries may limit your heart's supply of oxygen-rich blood, resulting in weakened heart muscle.
- Heart attack. ...
- Heart valve disease. ...
- High blood pressure. ...
- Irregular heartbeats. ...
- Congenital heart disease. ...
- Diabetes. ...
- Some diabetes medications.

www.medicalcoupons.in

22. Identify the condition due to the Image shown below?



(a) Hypercholesterolemia
(b) Hypertriglyceridemia
(c) Hyper alpha lipoproteinemia
(d) Hyper beta lipoproteinemia

Ans. (a) Hypercholesterolemia

Hypercholesterolemia, also called **high cholesterol**, is the presence of high levels of cholesterol in the blood.

Some types of hypercholesterolemia lead to specific physical findings. For example, familial hypercholesterolemia (Type IIa hyperlipoproteinemia) may be associated with xanthelasma palpebrarum (yellowish patches underneath the skin around the eyelids), arcus senilis (white or gray discoloration of the peripheral cornea), and xanthomata(deposition of yellowish cholesterol-rich material) of the tendons, especially of the fingers. Type III hyperlipidemia may be associated with xanthomata of the palms, knees and elbows

23. A 12-year-old child having proteinuria 4+ and edema and features associating with nephrotic syndrome without hematuria. Which drug would you prefer?

(a) Prednisolone
(b) Cyclosporine
(c) Cyclophosphamide
(d) Methotrexate

Ans. (a) Prednisolone

The term "nephrotic syndrome" refers to a distinct constellation of clinical and laboratory features of kidney disease. It is specifically defined by the presence

www.medicalcoupons.in

of **heavy proteinuria**(protein excretion greater than 3.5 g/24 hours), hypoalbuminemia (less than 3.5 g/dL), and peripheral edema.

Nephrotic syndrome is a condition where the kidneys leak protein from the blood into the urine. When untreated, children can suffer from serious infections. In most children with nephrotic syndrome, this protein leak resolves with cortic steroid drugs (prednisone, prednisolone) **reducing the risk of serious infecti n**

24. A 65-year-old patient presents with dementia and loss of cognition. Which is the most common cause for this condition?

(a) Alzheimer's
(b) Pick's
(c) Parkinson's
(d) CJD

Ans. (a) Alzheimer's

Alzheimer disease is a neurodegenerative disease that is signified by cognitive decline, memory loss, and erratic behavior. Alzheimer's disease is **the most common type of dementia**. It is a progressive disease beginning with mild memory loss and possibly leading to loss of the ability to carry on a conversation and respond to the environment. Alzheimer's disease involves parts of the brain that control thought, memory, and language.

25. A patient is allergic to wheat and unable to eat it. O/E, he was tested positive with Anti-Endomysial Antibodies. What could be the diagnosis?

(a) Celiac sprue
(b) Acrodermatitis enteropathica
(c) Tropical sprue
(d) Thiamine deficiency

Ans. (a) Celiac sprue

IgA anti-endomysial antibodies are found in 90% coeliac patients. They are a **very specific & sensitive marker for coeliac disease and dermatitis** herpetiformis. The anti-endomysial antibody test has been suggested to be 98% sensitive and 98% specific for clinical or subclinical coeliac disease.
Celiac disease, also known as celiac sprue or gluten-sensitive enteropathy, is **a chronic disorder of the digestive tract** that results in an inability to tolerate gliadin, the alcohol-soluble fraction of gluten. Gluten is a protein commonly found in wheat, rye, and barley.

www.medicalcoupons.in

26. A 75-year-old female has a problem in walking long distances. O/E bruit heard in knee. On X-ray imaging, reduced joint space and osteophytes were seen. Most probable diagnosis is?

(a) Osteoarthritis
(b) Rheumatoid arthritis
(c) Psoriatic arthritis
(d) Gouty arthritis

Ans. (a) Osteoarthritis


Osteoarthritis is the most common form of arthritis, affecting millions of people worldwide. It occurs when the protective cartilage that cushions the ends of the bones wears down over time.

Osteoarthritis symptoms often develop slowly and worsen over time. Signs and symptoms of osteoarthritis include:

- **Pain.** Affected joints might hurt during or after movement.
- **Stiffness.** Joint stiffness might be most noticeable upon awakening or after being inactive.
- **Tenderness.** Your joint might feel tender when you apply light pressure to or near it.
- **Loss of flexibility.** You might not be able to move your joint through its full range of motion.
- **Grating sensation.** You might feel a grating sensation when you use the joint, and you might hear popping or crackling.
- **Bone spurs.** These extra bits of bone, which feel like hard lumps, can form around the affected joint.
- **Swelling.** This might be caused by soft tissue inflammation around the joint.
- **X-rays.** Cartilage doesn't show up on X-ray images, but cartilage loss is revealed by a narrowing of the space between the bones in your joint. An X-ray can also show bone spurs around a joint.

www.medicalcoupons.in

27. Comment on the image shown below?



a) Prayer sign
b) Phalen sign
c) Tinel Sign
d) Schamroth sign

Ans. Prayer sign
Inability to extend the proximal interphalangeal joints of the fourth and fifth fingers - Prayer sign
"Prayer sign," a feature of diabetic cheiroarthropathy. It is also known as diabetic stiff hand syndrome or limited joint mobility syndrome. It is found in both type 1 and type 2 diabetic patients. The prevalence increases with duration of diabetes. This condition is associated with and predictive of diabetic microvascular complications.

28.. An old man presents with complaints of breathlessness and occasional episodes of chest pain. General physical examination shows high-volume pulse and BP of 150/70 mm Hg. On cardiovascular examination, the apex beat displaced 1.5cm lateral to Mid clavicular line. On auscultation diastolic murmur is heard. What is the diagnosis?

a) Aortic insufficiency
b) Mitral insufficiency
c) Pulmonary insufficiency
d) Tricuspid insufficiency

Ans. Aortic insuffiency

Lateral and/or inferior displacement of the apex beat usually indicates enlargement of the heart, called **cardiomegaly**.

Aortic regurgitation (AR), also known as aortic insufficiency, is a form of valvular heart disease that allows for the retrograde flow of blood back into the left ventricle. Aortic regurgitation is characterized by **cardiomegaly on the chest radiographs**, which is predominantly caused by **LV enlargement**

www.medicalcoupons.in

29. A patient was asked to copy the clock face made by the physician. The image on right is made by the
patient. Which is correct about this image?






a) Apraxia
b) Agraphia
c) Hemianopia
d) Hemi-neglect

Ans. Apraxia

Apraxia is **a neurological disorder** characterized by the inability to perform learned (familiar) movements on command, even though the command is understood and there is a willingness to perform the movement. Both the desire and the capacity to move are present but the person simply cannot execute the act.

The clock-drawing test is used for screening for **cognitive impairment and dementia** and as a measure of spatial dysfunction and neglect. It was originally used to assess visuo-constructive abilities but we know that abnormal clock drawing occurs in other cognitive impairments.

30. A 30-year-old patient presents with acute abdomen. X Ray Abdomen was done in standing position. Which of the following is correct about management of this condition?



a) Drain the peritoneal fluid
b) X Ray guided drainage of peritoneal accumulation
c) Exploratory laparotomy
d) Insert Flatus tube and repeat X ray

www.medicalcoupons.in

Ans. Exploratory Laparotomy

Exploratory laparotomy has been the main stay of the treatment in patients with **acute abdomen**. Laparoscopic management has previously been considered as a relative contraindication for acute small bowel obstruction and peritonitis.


31. The following CT abdomen shows



a) Pan Cake Kidney
b) Polycystic kidney
c) Horse Shoe kidney
d) Renal cell cancer

Ans. Horseshoe Kidney
Horseshoe kidney when the 2 kidneys join (fuse) together at the bottom to form a U shape like a horseshoe. It is also known as **renal fusion**. The condition occurs when a baby is growing in the womb, as the baby's kidneys move into place. Horseshoe kidney can occur alone or with other disorders.

32. Comment on the diagnosis based on X Ray abdomen image shown here



a) Congenital hypertrophic pyloric stenosis
b) Duodenal atresia
c) Congenital diaphragmatic hernia
d) Transient Tachypnoea of newborn

Ans. Duodenal Atresia

www.medicalcoupons.in

Duodenal atresia is a **condition in which the first part of the small bowel (the duodenum) has not developed properly**. It is not open and cannot allow the passage of stomach contents.
The **double bubble sign** is seen in infants and represents dilatation of the proximal duodenum and stomach. It is seen in both radiographs and ultrasound, and can be identified antenatally

33. A 60-year-old with restricted ambulation and prolonged bed rest due to acute lumbago presented with severe pain in left leg below the knee. On examination discoloration of left calf with significantly increased girth is noted. What is the clinical diagnosis?

a) Acute lymphangitis
b) Superficial migratory thrombophlebitis
c) Deep vein thrombosis
d) Milroy disease

Ans. Deep vein thrombosis

Deep vein thrombosis (DVT) occurs when a blood clot (thrombus) forms in one or more of the deep veins in your body, usually in your legs. Deep vein thrombosis can cause leg pain or swelling but also can occur with no symptoms.

You can get DVT if you have certain medical conditions that affect how your blood clots. A blood clot in your legs can also happen if you don't move for a long time, such as after you have surgery or an accident, when you're traveling a long distance, or when you're on bed rest.

DVT signs and symptoms can include:

- Swelling in the affected leg. Rarely, there's swelling in both legs.
- Pain in your leg. The pain often starts in your calf and can feel like cramping or soreness.
- Red or discolored skin on the leg.
- A feeling of warmth in the affected leg.

34. A cricket player was hit on the head with the cricket ball and was fine and then collapsed suddenly. He woke up to play the match brilliantly but was later found unconscious in the dressing room.Which of the following is correct about this presentation?

www.medicalcoupons.in

a) Extradural hemorrhage
b) Subdural hemorrhage
c) Cortical venous thrombosis
d) Sub arachnoid hemorrhage

Ans. Extradural Hemorrhage

THE LUCID interval following head trauma and unconsciousness is described classically in **epidural hematomas**, the lucid interval occurs **after the patient is knocked out by the initial concussive force of the trauma and then temporarily recovers**, before lapsing into unconsciousness again when bleeding causes the haematoma to expand past the extent for which the body can compensate.

35. Boy baby is brought at 3 weeks of age with recurrent episodes of non-bilious vomiting. Diagnosis is?

a) Duodenal atresia
b) Divarication of recti
c) CHPS
d) Peri-umblical hernia

Ans. CHPS

Congenital Hypertrophic pyloric stenosis (CHPS) is the idiopathic thickening of gastric pyloric musculature which later results in progressive gastric outlet obstruction.

Clinical presentation

Symptoms typically clinically manifests between 2 to 12 weeks of age. The presentation is typical with non-bilious projectile vomiting. The hypertrophied pylorus is palpated as an olive-sized mass in the right upper quadrant. A succussion splash can be heard hours after the last meal. Due to the loss of hydrochloric acid in the gastric contents from persistent vomiting, there is electrolyte imbalance, specifically the characteristic hypochloremic metabolic alkalosis.

The baby in early stage of the disease remains hungry and sucks vigorously after episodes of vomiting. Delay in diagnosis can lead to dehydration, malnutrition, metabolic alterations, and lethargy. The pyloric stenosis can be associated with Turner syndrome, tracheo-esophageal fistula, esophageal atresia and trisomy 18.

Diagnosis

The diagnosis is easily made if the presenting clinical features are typical, with projectile vomiting, visible peristalsis, and a palpable pyloric tumor. An enlarged pylorus, described as an "olive," can be palpated in the right upper quadrant or epigastrium of the abdomen.

www.medicalcoupons.in

Ultrasound is the modality of choice, and Muscle wall thickness 3 mm or greater and pyloric channel length 14 mm or greater are considered abnormal in infants younger than 30 days.

Barium meal Upper GI study reveals delayed gastric emptying, peristaltic waves (caterpillar sign), elongated pylorus with a narrow lumen (string sign). The entrance to the pylorus looks like beak (beak sign)

36. An 8-year-old child presents with a localized painless swelling slowly growing in size since childhood. Which is best treatment for this condition?



a) Excision
b) Chemotherapy
c) Intralesional steroids
d) Radiotherapy

Ans.Excision

A dermoid cyst is a congenital choristoma of the orbit. A choristoma is a benign tumor consisting of histologically normal cells occurring in an abnormal location. Dermoid cysts consist of keratinized epithelium and adnexal structures such as hair follicles, sweat glands, and sebaceous glands. They slowly enlarge with age.

Signs

- Palpable mass
- Ptosis
- Proptosis
- Strabismus
- Diplopia worsening proptosis with mastication
- Globe displacement
- Restriction in extraocular movements
- Inflammation
- Orbitocutaneous Fistula

www.medicalcoupons.in

Small, asymptomatic cysts may not require treatment. They may stabilize or even decrease in size over years. However, some surgeons opt for early excision to avoid the risk of traumatic rupture in the future

37. A young Guy comes to your OPD with the complaint that he develops pain in penis during erection and cannot have sex. Which is the likely diagnosis?

a) Paraphimosis
b) Phimosis
c) Hypospadias
d) Peyronie disease

Ans. Peyronie disease

Peyronie's disease is a **noncancerous condition resulting from fibrous scar tissue that develops on the penis and causes curved, painful erections**.
Peyronie's disease is caused by repeated penile injury, typically during sex or physical activity.
Penises vary in shape and size and having a curved erection isn't necessarily a cause for concern. In Peyronie's disease, the bend is significant, and may occur along with pain or interfere with sexual function.

PHARMACOLOGY

1. Which of the following is a SERM?

(a) Ethyl estradiole
(b) Tamoxifene
(c) Anastrozole
(d) Mifepristone

Ans. (b) Tamoxifen

SERM DRUGS

- **Tamoxifen**
- Tamoxifen analogs
  → Toremifine, Droloxifene, Idoxifene
- Fixed ring compounds :
  → Raloxifene, Lasofoxifene, Aroxif, Miproxifene, Levormeloxifene, EM652

2. Which among the following is a monoclonal antibody used for treatment of cancer?

www.medicalcoupons.in

(a) Cisplatin
(b) Rituximab
(c) Vincristine
(d) Cyclophosphamide

Ans. (b) Rituximab

Rituximab is **the first MAb approved for cancer therapy**. Clinical trials indicate that rituximab is efficacious and safe for recurrent or chemotherapy resistant, B-cell, low-grade NHL.

3. Find out the best treatment for the given condition?

(a) Valacyclovir

(b) Imipenem

(c) Carbapenem

(d) Penicillin

Ans. (a) Valacyclovir

Valacyclovir is used to **treat herpes virus infections**, including herpes labialis (also known as cold sores), herpes zoster (also known as shingles), and herpes simplex (also known as genital herpes) in adults. It is also used to treat chickenpox and cold sores in children.

4. A patient was taking Theophylline for Bronchial asthma. After starting another drug, symptoms of theophylline toxicity started appearing. Which is the likely drug to cause theophylline toxicity?

(a) Rifampicin
(b) Erythromycin

(c) Valproate
(d) Phenobarbitone

Ans. (b) Erythromycin

www.medicalcoupons.in

www.medicalcoupons.in

Using erythromycin together with **theophylline may increase the effects of theophylline**

5. Antidote used for heparin toxicity?

(a) Vitamin K
(b) Protamine sulphate

(c) Ascorbic acid
(d) N acetylcysteine

Ans. (b) Protamine sulphate

Despite of the low therapeutic index, **protamine** is the only registered antidote of heparins. The toxicology of protamine depends on a complex interaction of the high molecular weight, a cationic peptide with the surfaces of the vasculature and blood cells.

6. A 65-year-old patient presented with dribbling of urine with urgency. He was started on prazosin therapy, but started developing postural hypotension. Which of the following is the better alternative drug for this patient?

(a) Tamsulosin
(b) Phenoxybenzamine

(c) Terazosin
(d) Timolol

Ans. (a) Tamsulosin

- **With prazosin and other alpha1 (postsynaptic) adrenergic blockers**, first dose hypotension due to arterial and venous dilatation is well recognised and is minimised by using a low starting dose. With chronic use, postural hypotension **may persist**.
- **Tamsulosin is well tolerated and has minimal effects on blood pressure**; tamsulosin 0.4 mg has the lowest potential to reduce blood pressure and causes less symptomatic orthostatic hypotension than terazosin.

7. A patient presented with hospital acquired pneumonia. On testing he was found to be allergic to Penicillin G. Which of the following drug is likely to be safe in this patient?

(a) Methicillin

(b) Cefotitan

www.medicalcoupons.in

(c) Imipenam

(d) Aztreonam

Ans. (d) Aztreonam

Aztreonam **can safely be used in all patients** with a penicillin or cephalosporin allergy with the exception of ceftazidime as they share a similar side chain.

8. A new drug is introduced in the market after which phase of clinical trial?

(a) I
(b) II
(c) III
(d) IV

Ans. (c) III

New Drug Application (NDA) is done following the Phase III Clinical Trials, the drug manufacturer analyzes all the data from the studies and files an NDA with the FDA (provided the data appear to demonstrate the safety and effectiveness of the drug). The NDA contains all of the data gathered to date about the drug.

9. A mother is suspected to have a premature delivery. Which of the following drug could be helpful to attain the lung maturity before birth?

(a) Aspirin
(b) Dexamethasone
(c) MgSO4
(d) NSAID

Ans. (b) Dexamethasone

Betamethasone and dexamethasone are the most widely studied corticosteroids, and they generally have been preferred for antenatal treatment to accelerate fetal organ maturation.

10. All of the following conditions get worsened after timolol maleate 0.5% administration except?

(a) Bronchial asthma
(b) Hypertension
(c) Depression
(d) Diabetes/Glaucoma

www.medicalcoupons.in

Ans. (b) HTN

It works by blocking the action of certain natural chemicals in your body such as epinephrine on the heart and blood vessels. **This effect lowers the heart rate, blood pressure, and strain on the heart**. This medication is also used to prevent migraine headaches. **So timolol doesn't worsen hypertension but used to lower it .**

11. Which anti-tussive should not be prescribed for the patients who are driving?


(a) Codeine
(b) Ambroxol
(c) Diphenhydramine
(d) Hydrocodone

Ans. (c) Diphenhydramine

First-generation antihistamines, such as diphenhydramine, are known to affect driving performance.

PHYSIOLOGY

1. Identify the movement taking place in the Intestine?



(a) Segmented movement
(b) Peristaltic movement
(c) Passive diffusion
(d) Intersegmental contraction

Ans. (b) Peristaltic movement

Peristalsis, involuntary movements of the longitudinal and circular muscles, primarily in the digestive tract but occasionally in other hollow tubes of the body, that

www.medicalcoupons.in

occur in progressive wavelike contractions. Peristaltic waves occur in the esophagus, stomach, and intestines.

2. A patient after trauma came to the ER. On neurological examination, he was found to have normal handwriting skills but could not speak. Where could be the lesion for his clinical condition?


(a) Wernicke's area
(b) Temporal area
(c) Broca's speech area
(d) Primary motor cortex

Ans. (c) Broca's speech area

Symptoms of Broca's aphasia include:

- poor or absent grammar
- difficulty forming complete sentences
- omitting certain words, such as "the," "an," "and," and "is" (a person with Broca's aphasia may say something like "Cup, me" instead of "I want the cup")
- more difficulty using verbs than nouns correctly
- difficulty articulating sounds and words
- difficulty repeating what has been said by others
- trouble with writing sentences
- difficulty reading
- problems with full comprehension
- difficulty following directions
- frustration


3. **Pressure – volume graph given and asked to find out A accordingly:**




(a) Aortic valve closing
(b) Aortic valve opening
(c) Mitral valve opening
(d) Mitral valve closing

www.medicalcoupons.in

Ans. (b) Aortic valve opening



4. Which part of nephron does water rea. sorp .ion takes place more:




(a) A
(b) B
(c) C
(d) D

Ans. (c) C

The proximal convoluted tubule is where a majority of reabsorption occurs. About 67 perc nt of the water, Na^+, and K^+ entering the nephron is reabsorbed in the proxim convoluted tubule and returned to the circulation.

www.medicalcoupons.in



Structure of a nephron

5. Inhibition of Na +/ K+ ATPase pump results in which of the following?


(a) First depolarize and then hyperpolarize
(b) Hyperpolarisation
(c) Depolarisation
(d) Repolarisation

Ans. (c) Depolarisation

Na +/ K+ ATPase pump is essential for the maintenance of Na^+ and K^+ concentrations across the membrane. ... Inhibition of this pump, therefore, causes **cellular depolarization resulting not only from changes in Na^+ and K^+ concentration gradients, but also from the loss of an electrogenic component of the resting membrane potential.**

MICROBIOLOGY

1. A patient complaints of frequent perianal pruritus. On microscopic examination, following Image was seen. Find out the most probable causative organism?



www.medicalcoupons.in

(a) Ascaris lumbricoides
(b) Enterobius vermicularis
(c) Trichuris Trichura
(d) Ancylostoma duodenale

Ans. (b) Enterobius vermicularis

Enterobius vermicularis, also called pinworm, is one of the most common nematode infections in the world. Humans are the only natural host for this infection. *Enterobius vermicularis* is an organism that primarily lives in the ileum and cecum.
Once *E. vermicularis* eggs are ingested, they take about 1 to 2 months to develop into adult worms which happens in the small intestine. These do not usually cause any symptoms when confined to the ileocecal area. The female adult worms and ova migrate to the anal area mostly at night time and deposit thousands of eggs in the perianal area. This migration causes pruritus. Eggs hatch near the anal area causing perianal pruritus.

2. A patient with history of travel to Ohio shows lung symptoms. What could be the most common cause?

(a) Blastomycosis
(b) Histoplasmosis
(c) Mucormycosis
(d) Coccidiodomycosis

Ans. (a) Blastomycosis

- Blastomycosis is an infection caused by a fungus called *Blastomyces*. The fungus lives in the environment, particularly in moist soil and in decomposing matter such as wood and leaves. *Blastomyces* mainly lives in areas of the United States and Canada surrounding the Ohio and Mississippi River valleys and the Great Lakes.
- Blastomycosis is **a fungal infection caused by inhaling Blastomyces dermatitidis spores**. If it involves only the lungs, it is called pulmonary blastomycosis. Only about half of people with the disease have symptoms, which can include fever, cough, night sweats, muscle pains, weight loss, chest pain, and feeling tired.

3. Rice water stools are characteristic of:

(a) Shigella
(b) Cholera
(c) E.coli
(d) Salmonella

www.medicalcoupons.in

Ans. (b)Cholera

The passage of profuse "rice-water" stool is characteristic of **severe cholera**. The stool is watery with flecks of mucous and has the appearance of water in which rice has been washed.

4. Which of the following is high level disinfectant?

(a) Gluteraldehyde
(b) Alcohol
(c) Lysol
(d) Phenol

Ans. (a) Gluteraldehyde

High-level disinfectants/liquid sterilants are to be used as immersion products and not for surface disinfection. Examples are glutaraldehydes, special hydrogen peroxide, and special peracetic acid products. Items submerged should be limited to those that cannot be heat-sterilized (such as heat-sensitive plastic items).

5. Identify the organism stained with India Ink stain?

(a) Chlamydia
(b) Histoplasma
(c) Cryptococcus
(d) Aspergillus

Ans. (c) Cryptococcus

India ink capsule stain is used to demonstrate cell capsules through microscopic examination. This procedure is used to detect presence of encapsualted species, e.g. **Cryptococcus neoformans**, which causes cryptococcosis in humans.

6. **A pregnant woman with malaria presents in front of the doctor. What could be the causative organism**?

www.medicalcoupons.in

(a) P. vivax
(b)P. falciparum
(c) P. ovale
(d) P. malariae

Ans. (b) P. falciparum

Pregnant women are susceptible to malaria during pregnancy. **Plasmodium falciparum**, which sequesters in the placenta, causes the greatest disease, contributing significantly to maternal and infant mortality.

7. **PrP deposition is seen in**?

(a) CJD
(b) Rabies
(c) Parkinson's disease
(d) Alzheimer's disease

Ans. (a) CJD

Creutzfeldt-Jakob disease (CJD) is caused by an abnormal **infectious protein in the brain called a prion.** (prion protein PrP)

8. **A patient presented with colicky abdominal pain, diarrhea and no history of fever. What could be the diagnosis?**

(a) E. coli
(b) Entamoeba histolytica
(c) Shigella
(d) Giardia lamblia

Ans. (b) Entamoeba histolytica

Amoebiasis or amoebic dysentery is an infection caused mainly by the **protozoan parasite Entamoeba histolytica.** Symptoms may include lethargy, loss of weight, colonic ulcerations, abdominal pain, diarrhea, or bloody diarrhea.

9. **A male patient presented with bloody diarrhea. On Investigation of stool sample, the organism was found to be gram negative, non-motile and oxidase negative. What could be the cause?**


www.medicalcoupons.in

(a) Shigella
(b) E. coli
(c) Salmonella
(d) Yersinia pestis

Ans. (a) Shigella

Shigellae are Gram-negative, nonmotile, facultatively anaerobic, non-spore-forming rods.

10. A post covid-19 patient started to develop nasal congestion, black crusts in nose, and blackish discoloration of skin on nasal and cheek area. Which of the following micro-organisms is the causative agent ?



(a) Aspergillus
(b) Fusarium
(c) Mucormycosis
(d) Pencillium


Ans. (c) Mucormycosis

www.medicalcoupons.in

Mucormycosis is **a fungal infection caused by** a group of microorganisms belonging to the phylum Glomeromycota. Once classified under rare fungal disease, Mucormycosis is regrouped under emerging pathogens. They are ubiquitous, found mainly in soil and decaying organic matter.
During the second wave of the COVID-19 pandemic in India, an unprecedented surge in cases of mucormycosis was observed: immune dysregulation caused by the SARS-CoV-2 and the use of broad-spectrum antibiotics and corticosteroids—particularly in patients with poorly controlled diabetes with ketoacidosis

ENT

1. **An infant presented with inspiratory stridor and the clinical picture shows omega shaped epiglottis. The probable diagnosis is?**



(a) Acute epiglottitis
(b) RLN palsy
(c) SLN palsy
(d) Laryngomalacia

Ans. (d) Laryngomalacia

This is the well-known "omega shaped" epiglottis in **laryngomalacia**.
Laryngomalacia is **a congenital softening of the tissues of the larynx (voice box) above the vocal cords**. This is the most common cause of noisy breathing in infancy. Another common finding of laryngomalacia involves the posterior or back part of the larynx, where the arytenoid cartilages or the mucosa/tissue over the arytenoid cartilages can collapse into the airway and cause airway obstruction.

www.medicalcoupons.in

2. Identify the Image shown below:



(a) Rhinophyma
(b) Rhinosporidiosis
(c) Rhinoscleroma
(d) Rhinosinusitis

Ans. (a) Rhinophyma (aka potato nose)

Rhinophyma is **a skin disorder characterized by a large, red, bumpy or bulbous nose**. It can occur as part of phymatous rosacea. The exact cause of rhinophyma is unknown, but it's considered a subtype of severe rosacea. This condition is significantly more common in men, especially between the ages of 50 to 70 years.

3. What is the best management for Atrophic Rhinitis?

(a) FESS
(b) Young's operation
(c) Nasal steroid spray
(d) Immunotherapy

Ans. (b) Young's operation

Young's operation is a surgery designed for the **treatment of atrophic rhinitis**, first described by Austen Young in 1967. Young's operation is a surgery designed for the **treatment of atrophic rhinitis**, first described by Austen Young in 1967.

4. True statement about BERA?

(a) Used to detect early hearing impairment
(b) Invasive test
(c) It is a subjective test
(d) None of the above

www.medicalcoupons.in

Ans. (a) Used to detect early hearing impairment

Brainstem-evoked response audiometry (BERA) is a simple, noninvasive, objective test **for early identification of hearing impairment in children and neonates**. It can be used as a screening test and is useful in newborns, infants, and other difficult-to-test patients.

5. Laryngeal crepitus is seen in all except?

(a) Tongue cancer
(b)Pneumonia
(c) Laryngitis
(d) Post cricoid carcinoma

Ans. (d) Post cricoid carcinoma

Cause of decreased bone conduction on audiogram?

(a) Ossicular dislocation
(b) Cochlear pathology
(c) Fixation of foot plate of stapes
(d) EAC pathology

Ans. (b) Cochlear pathology

6. **The cause of Nasopharyngeal carcinoma is?**

(a) HSV
(b) EBV
(c) HZV
(d) Parainfluenza virus

Ans. (b) EBV

The nasopharynx is located at the very back of the nose near the Eustachian tubes. Nasopharyngeal carcinoma is more common in Southeast Asia and is frequently, but not always, caused by **Epstein-Barr virus (EBV)**.

www.medicalcoupons.in

7. Identify the procedure shown below?



(a) Hemlich maneuver
(b) Epleys maneuver
(c) Semonts maneuver
(d) Cottles maneuver

Ans. (d) Cottles maneuver

The Cottle's maneuver (CM) is a **test in which the cheek on the side to be evaluated is gently pulled laterally with one to two fingers to open the valve**. This test is used to determine if the most significant site of nasal obstruction is at the valve or farther inside the nasal cavity.

8. What is the most common causative agent in croup?

a) Measles virus
b) Parainfluenza virus
c)Respiratory syncytial virus
d) Corynebacterium diphtheriae

Ans. (a) Parainfluenza Virus

Croup is commonly caused by virus (75% of cases), less commonly by bacteria. **In most cases the causative agent is parainfluenza virus type 1 and 2 (a member of the paramyxovirus family).** Croup can also be caused by adenovirus ,measles virus, respiratory syncytial virus. The most common bacteria associated with croup are Staphylococcus aureus, Haemophilus influenzae, Streptococcus pneumonia and Moraxella catarrhalis.

9. A patient presented to ER after fracture of base of skull with CSF Rhinorrhea. On further investigation the following image was obtained. What is the name of this sign shown below?


www.medicalcoupons.in



(a) Schwartz sign
(b) Brown sign
(c) Halo sign
(d) Phelp sign

Ans. (c) Halo sign

A “halo” or “ring” sign, occurs when cerebrospinal fluid **(CSF) mixes with blood on an absorbent surface**. The blood forms a spot in the center and a lightly stained ring forms a halo around it.



10. The most appropriate investigation for JNA is?

(a) PET
(b) CT
(c) Angiography
(d) X-ray

Ans. (b) CT

To confirm a diagnosis of **Juvenile nasopharyngeal angiofibroma**, **your doctor will order a CT scan or MRI**. These scans help doctors determine the exact location of the tumor and see how far it has spread.

11. A child presented with a high arched palate and mouth breathing and failure to thrive. The probable diagnosis is?

www.medicalcoupons.in

(a) Tonsil hypertrophy
(b) Adenoid hypertrophy
(c) Lingual tonsil hypertrophy
(d) Turbinate hypertrophy

Ans. (b) Adenoid hypertrophy

Adenoid hypertrophy is **an obstructive condition related to an increased size of the adenoids**
Adenoid facies refers to the atypical appearance of facial features, and can result from persistent adenoid hypertrophy. Commonly, adenoid facies is known as "long face syndrome", and is characterized by a long, lean face with an open mouth. With adenoid facies, individuals typically present with **increased mouth breathing, an arched palate,** underdeveloped upper jaw bones (i.e. hypoplastic maxilla), a short upper lip, elevated nostrils, and dental crowding of the front teeth. It most commonly occurs in children and often presents with chronic nasal obstruction.

12. A patient presented with pulsation of the tonsil. The cause is?

(a) Aneurysm of the common carotid
(b) Aneurysm of the external carotid
(c) Carotid body tumor
(d) ICA

Ans. (d) ICA

Pulsating Tonsil Due to **Medial Displacement of the Internal Carotid Artery**

RADIOLOGY

1. Thickness of Lead Apron?

(a) 0.1
(b) 0.2
(c) 0.5
(d) 0.01

Ans. (c) 0.5

Lead aprons are the primary radiation protective garments used by personnel during fluoroscopy. The radiation protection provided by a lead apron is approximately the same as **0.25- to 1-mm thick lead**. An apron with 0.5-mm thickness can attenuate approximately 90% or more of the scatter radiation.

www.medicalcoupons.in

2. A female visits OPD with her 6-week child presenting with vomiting and abdominal pain. X-ray of the infant shows?



(a) Jejunal atresia
(b) Diaphragmatic hernia
(c) Duodenal atresia
(d) Pyloric stenosis

Ans. (c) Duodenal Atresia → Image shows Double Bubble Sign

The double bubble sign is a classic radiographic manifestation **of duodenal obstruction**, the cause of which could be intrinsic (such as duodenal atresia, duodenal stenosis or duodenal web) or extrinsic (such as annular pancreas or rotational anomalies).

3. A two-weeks child brought to the emergency room by her mother with the symptoms of non-bilious vomiting and lump in the epigastric area. What is the most appropriate diagnosis?

(a) Congenital pyloric stenosis
(b) Jejunal atresia
(c) Duodenal atresia
(d) Esophageal atresia

Ans. (a) Congenital pyloric stenosis

Pyloric stenosis should be suspected in any young infant with severe vomiting. Infants with pyloric stenosis classically present with projectile, **non bilious vomiting.** Vomiting may be intermittent or occur after each feedingOn physical exam, palpation of the abdomen **may reveal a mass in the epigastrium**. This mass, which consists of the enlarged pylorus, is referred to as the 'olive'.

www.medicalcoupons.in

4. Patient comes to emergency room with vomiting and colicky abdominal pain. On investigation following Image is obtained. Likely diagnosis is?



(a) Ectopic kidney
(b) Horse-shoe kidney
(c) Pan cake kidney
(d) Cross fused kidney

Ans. (b) Horse-shoe kidney

Horseshoe kidney is **a condition in which the kidneys fuse (bind) together at the bottom**, forming a "U" shape or horseshoe shape. Children who have horseshoe kidney have one "fused" kidney instead of 2 separate kidneys.


5. A patient presents with hard palpable gall bladder with pain in right hypochondrium. On performing USG following Image was seen. What could be the diagnosis?



(a) Multiple gall stones
(b) Choledocolithiasis
(c) Hydatid cyst
(d) Gall bladder cancer

www.medicalcoupons.in

Ans. (a) Multiple gall stones



Gallbladder with multiple gallstones. Arrows indicate stones with associated posterior acoustic shadowing

6. Identify the radiograph shown below?



a) Popcorn calcification
b) Egg shell calcification
c) Ground Glass Appearance
d) Miliary opacities

Ans. Egg shell calcification

Peripheral calcification of lymph nodes, "eggshell calcification," commonly occurs in patients with **sili**

www.medicalcoupons.in

cosis and coal-worker's pneumoconiosis. Sarcoidosis, postirradiation Hodgkin disease, blastomycosis, and scleroderma are other reported causes.


7. **A 5 year old child has been brought to hospital with complaints of discomfort in the hip joint, limping while walking and decreased range of motion in hip. The following radiograph was taken, what will be the diagnosis ?**



(a) Slipped capital femoral epiphysis
(b) Perthes disease
(c) Myositis ossificans
(d) Meyer's dysplasia

Ans. (b) Perthes disease

Perthes disease is a **rare childhood condition that affects the hip**. It occurs when the blood supply to the rounded head of the femur (thighbone) is temporarily disrupted. Without an adequate blood supply, the bone cells die, a process called avascular necrosis.



www.medicalcoupons.in


8. A 6-day-old neonate with respiratory distress showed the following findings on the chest X-ray. What is the diagnosis?



(a) Congenital cystic adenomatoid malformation
(b) Bronchogenic cysts
(c) Hyaline membrane disease
(d) Congenital diaphragmatic hernia


Ans. (d) Congenital Diaphragmatic Hernia

Congenital diaphragmatic hernia (CDH) occurs **when there is a hole in the diaphragm**, which is the thin sheet of muscle separating the chest from the abdomen. When this gap forms during a fetus's development in the womb, the bowel, stomach or even the liver can move into the chest cavity.



www.medicalcoupons.in

9. A patient presenting with breathlessness and pedal edema gives a history of MI few years back. A chest radiograph is taken and shown below. Likely diagnosis is?




(a) Pericardial effusion
(b) Constrictive pericarditis
(c) Cardiac tamponade
(d) Dilated cardiomyopathy

Ans. (a) Pericardial effusion

Pericardial effusion is **the build up of extra fluid in the space around the heart**. If too much fluid builds up, it can put pressure on the heart. This can prevent it from pumping normally. A fibrous sac called the pericardium surrounds the heart. This sac consists of two thin layers.

Pericardial effusion symptoms include:

- Shortness of breath or difficulty breathing (dyspnea)
- Discomfort when breathing while lying down (orthopnea)
- Chest pain, usually behind the breastbone or on the left side of the chest
- Chest fullness
- Swelling in the legs or abdomen



www.medicalcoupons.in


10. **A patient of NPC (Nasopharyngeal carcinoma) presented with the following finding as shown below. Which of the following sign is marked by the arrow?**



(a) Griesenger sign
(b) Phelp sign
(c) Holman miller sign
(d) DODD sign

Ans. (c) Holman miller sign


The Holman-Miller sign (also called the antral sign) is seen in **juvenile nasopharyngeal angiofibroma**; it refers to **the anterior bowing of the posterior wall of the maxillary antrum** as seen on a lateral skull radiograph or cross-sectional imaging





www.medicalcoupons.in

11. Identify the diagnosis on Imaging as shown below?



(a) Incomplete mole
(b) Hydatidiform mole
(c) Retained products of conception
(d) PCOD

Ans. (b) Hydatidiform mole

A **molar pregnancy** also known as hydatidiform mole is a rare complication of pregnancy characterized by the abnormal growth of trophoblasts, the cells that normally develop into the placenta. There are two types of molar pregnancy, complete molar pregnancy and partial molar pregnancy.


Radiographic features

Ultrasound

- enlarged uterus
- may be seen as an intrauterine mass with cystic spaces without any associated fetal parts,**the multiple cystic structures classically give a "<u>snowstorm</u>" or "<u>bunch of grapes</u>" type appearance.**
- may be difficult to diagnose in the first trimester
  - may appear similar to a normal pregnancy or as an empty gestational sac
  - <50% are diagnosed in the first trimester
- bilateral <u>theca lutein cysts</u> may also be seen on ultrasound
- color Doppler interrogation may show high velocity with a low impedance flow

A 27 year old married women comes to hospital with complains of irregular menstruation, hirsutism and not able to conceive ,the following usg image was observed in that patient?

www.medicalcoupons.in




(a) Ovarian hyperstimulation syndrome
(b) Ovarian cyst
(c) PCOS
(d) Endometriosis

Ans. (c) PCOS

- Polycystic ovary syndrome (PCOS) is a **condition in which the ovaries produce an abnormal amount of androgens**, male sex hormones that are usually present in women in small amounts. The name polycystic ovary syndrome describes the numerous small cysts (fluid-filled sacs) that form in the ovaries.

The string of pearls (or beads) sign in the ovary is the **ultrasound sign of multiple small (<5 mm) follicles peripherally located in the gonad**. It is typically seen in polycystic ovaries, and is thought to represent abnormal accumulation of immature follicles.

12. A cattle handler from a village with cough, hemoptysis and chest pain underwent a chest radiograph and is shown below. What is the probable diagnosis?



www.medicalcoupons.in


(a) Hydatid cyst
(b) Anthrax
(c) Byssinosis
(d) Abscess/Aspergillosis

Ans. (a) Hydatid cyst

- Cystic echinocccosis (CE), also known as hydatid disease, is **caused by infection with the larval stage of Echinococcus granulosus**, a 2–7 millimeter long tapeworm found in dogs (definitive host) and sheep, cattle, goats, and pigs (intermediate hosts).
- **Cough, hemoptysis, and chest pain** are the most common symptoms in patients with hydatid disease of the lungs.

OPHTHALMOLOGY

1. A tumor in the anterior pituitary causing pressure over optic chiasma will present as?



(a) Homonymous hemianopia
(b) Bitemporal hemianopia
(c) Heteronymous hemianopia with central sparing
(d) Monocular vision loss

Ans. (b) Bi-temporal Hemianopia

When there is compression at optic chiasm, the **visual impulse from both nasal retina are affected**, leading to inability to view the tempora, or peripheral, vision. This phenomenon is known as bitemporal hemianopsia.

www.medicalcoupons.in

2. Ophthalmic surgical Instrument shown in the Image is used for surgery of ?



(a) Entropion
(b) Ectropion
(c) Chalazion
(d) Bitot spot

Ans. (c) Chalazion

A chalazion is **a red bump on your eyelid**. It is sometimes called an eyelid cyst or a meibomian cyst. It slowly forms when an oil gland (called a meibomian) becomes blocked.



A chalazion forceps is a self-retaining type of thumb forceps that has discoid ends. One arm has a flattened plate at its end, which is round and solid. The other arm has a matching ring that fits into the opposite arm's solid plate. There is an adjustable screw at the middle of the plates, which can be tightened until it has a firm and fine grip to the cyst or lesion.

3. Corneal transparency is maintained by?

(a) Chondratin sulphate
(b) Keratan sulphate
(c) Hyaluronic acid
(d) Dermatan sulphate

www.medicalcoupons.in

Ans. (b) Keratan sulphate

Keratan sulfate (KS) also plays a role in maintaining the transparency of the cornea by regulating the **organization of collagen fibrils** and maintaining corneal hydration. Alteration of corneal KS may lead to reduced hydration and corneal opacification. KS is also implicated in corneal wound repair.

4. **5-year-old suffering with chronic epiphora has appeared in the clinic. What is the next step of management?**

(a) Laryngeal probing and massaging
(b) Dacrocystorhinostomy
(c) Tarsal plate fracture
(d) Lacrimal sac massage

Ans. (b) Dacryocystorhinostomy

- Epiphora is **an overflow of tears onto the face**, other than caused by normal crying. It is a clinical sign or condition that constitutes insufficient tear film drainage from the eyes, in that tears will drain down the face rather than through the nasolacrimal system.
- Dacryocystorhinostomy (DCR) is **a surgery that creates a new path for tears to drain between your eyes and your nose**. You may need this surgery if your tear duct has become blocked.

5. **Beta Blocker is avoided in the below conditions except?**

(a) Glaucoma
(b) Peripheral vascular disease
(c) COPD
(d) Diabetes

Ans. **Glaucoma**

- Glaucoma is a **group of eye conditions that damage the optic nerve**, the health of which is vital for good vision. This damage is often caused by an abnormally high pressure in your eye.
- Beta Blockers decrease the **pressure** inside your eyes by reducing how much fluid (aqueous humor) is produced in the eyes. Reducing pressure in the eyes, slows down optic nerve damage which greatly decreases the rate of vision loss.

www.medicalcoupons.in

6. **Patient wearing contact lens since two years develops redness and decreased vision. What could be the cause?**

(a) Bacterial corneal ulcer
(b) Giant cell papillary
(c) Fungal keratitis
(d) Conjunctivitis

(e) Ans. Bacterial corneal ulcer

- Contact lens wear is a common predisposing factor in **Bacterial keratitis** and is one of the two preventable risk factors for corneal infection in a working age population.
- A bacterial corneal ulcer, or bacterial keratitis, is **an infection of the corneal stroma that causes rapid visual loss and pain**. Infectious corneal ulcers need to be treated as soon as possible to preserve vision.

7. **Identify the condition?**




(a) Chronic melanoma
(b) Retinoblastoma
(c) HSV keratitis
(d) Geographic ulcer

Ans. (b) Retinoblastoma

www.medicalcoupons.in

Retinoblastoma is **an eye cancer that begins in the retina** — the sensitive lining on the inside of your eye. Retinoblastoma most commonly affects young children, but can rarely occur in adults.



8. A patient presented with anterior uveitis in OPD. Which of the following can be associated with his condition?

(a) HLA B5
(b) HLA B27
(c) HLA B7
(d) HLA DR4

Ans. (b) HLA B27

Acute anterior uveitis (AAU) is the most common form of uveitis, accounting for approximately 90% of all cases. **Half of all cases of AAU are HLA-B27 positive**. The disease is typically acute in onset, unilateral, no granulomatous inflammation involving the iris and ciliary body, with a tendency to recurrent attacks.

9. A patient with miosis, anhidrosis and mild drooping of eyelid is presented to the clinic. What is the suspected diagnosis?




www.medicalcoupons.in

(a) III cranial nerve palsy
(b) Horner's syndrome
(c) VI cranial nerve palsy
(d) IV cranial nerve palsy

Ans. (b) Horner's syndrome

Horner syndrome is a **combination of signs and symptoms caused by the disruption of a nerve pathway from the brain to the face and eye** on one side of the body. Typically, Horner syndrome results in a decreased pupil size, a drooping eyelid and decreased sweating on the affected side of your face.

10. What is not correct regarding following Image?



(a) Occurs more commonly in sunny areas
(b) Amniotic membrane transplant
(c) It causes myopia
(d) Iron deposition is found


Ans. (c) It causes myopia

The give condition in the image is pterygium.

Pterygium
→ Triangular overgrowth of subconjunctival tissue.
→ Wedge shaped, pink -white translucent membrane with apex extending into cornea.
→ Stocker's line - iron line at leading edge
Risk factors - Exposure to UVrays, hot, dry weather, wind, dust
Symptoms
→ Irritation, lacrimation, FB sensation, decreased vision , difficulty in contact lens wear
Treatment
→ Excision with conjunctival autograft
→ Amniotic membrane transplant with fibrin glue / sutures.

www.medicalcoupons.in

11. A 15-year-old boy presents to clinic with difficulty in coordination of movements and abdominal pain. On further examination copper deposition in a ring form is seen in cornea. Most likely diagnosis?



(a) Keratitis
(b) Munson sign
(c) Wilson's disease
(d) Neuroblastoma

Ans. (c) Wilson's disease

Wilson's disease is a **genetic disorder in which excess copper builds up in the body**.
Individuals with Wilson disease, copper deposits in the front surface of the eye (the cornea) form a green-to-brownish ring, called the Kayser-Fleischer ring.

OBG

1) A 55-year-old female presents with complaints of post coital bleeding. She is mother of 6 children and has a history of early marriage. On colposcopic examination, inflammed cervix is seen which bleeds on touch. What is the most probable diagnosis?




(a) Cervical cancer
(b) Cervical cyst
(c) Endometrial carcinoma
(d) Vulval carcinoma

Ans. (a) Cervical Cancer

Cervical Cancer is a malignant tumour of the lower-most part of the uterus (womb) that can be prevented by PAP smear screening and a HPV vaccine.
Symptoms include bleeding in between periods and after sexual intercourse. Foul smelling white discharge and low back pain or lower abdominal pain may also occur. In some cases there may be no symptoms.


www.medicalcoupons.in

2) Identify the procedure being done in the Image?



(a) Leopold's Ist maneuver
(b) Leopold's IInd maneuver
(c) Leopold's IIIrd maneuver
(d) Leopold's IVth maneuver

Ans. (c) Leopold's IIIrd maneuver



Maneuver I: The uterine contour is outlined. The fundus is palpated with the fingertips of both hands facing toward the maternal xiphoid cartilage. This should allow the identification of the fetal parts in the upper pole (fundus) of the uterus.

Maneuver II: Once an assessment is made of the fetal part present in the uterine fundus, the hands are placed at either side of the maternal abdomen. With this maneuver, the examiner will be able to determine the location of the fetal back.

Maneuver III: Using one hand, the examiner will grasp the presenting part between the thumb and fingers. This is done on the lower abdomen, a few centimeters above the symphysis pubis. This will allow the examiner to develop a further identification of the presenting part and assessment of its engagement.

Maneuver IV: This last maneuver resembles the first one, but instead of facing the fundus, the examiner faces the pelvis of the patient. The palms of both hands are placed on either side of the lower maternal abdomen, with the fingertips facing toward the pelvic inlet. This maneuver should allow the identification of the fetal parts in the lower pole of the uterus.

www.medicalcoupons.in

3) Look at the Image and diagnose the condition?



a. Kleinfelter
b. Down's
c. CAH
d. Turner's

Ans. D - Turner's syndrome

Turner syndrome is a **genetic disorder affecting girls and women**. The cause of Turner syndrome is a completely or partially missing X chromosome. Turner syndrome symptoms include short stature and lack of breast development and periods. Treatment for Turner syndrome may include hormone therapy.

- Broad chest.
- Cubitus valgus, where the arms point out slightly at the elbows.
- Dental problems.
- Eye problems, such as a lazy eye or drooping eyelids.
- Scoliosis, when the spine curves sideways.
- Low hairline at the back of the neck.
- Many skin moles.
- Missing knuckle in a particular finger or toe, making the digit shorter.
- Narrow fingernails and toenails.
- Small lower jaw.
- Swelling of the hands and feet.
- Unusually short, wide neck or webbed neck (extra skin folds).

www.medicalcoupons.in

4) An 8-weeks pregnant lady with spotting since the start of her pregnancy. On USG, gestational sac corresponds to gestational age. What is the probable diagnosis?

(a) Missed abortion
(b) Threatened abortion
(c) Inevitable abortion
(d) Incomplete abortion

Ans. (b) Threatened abortion

- **Threatened Abortion**
- It is a clinical entity where the process of abortion has started but has not progressed to a state from which **recovery is impossible**

✓ C/o - **Slight bleeding**
✓ O/E - Uterus **- corresponds to gestational age**, Cx - Internal os **is closed**
✓ USG - **live foetus** with subchorionic haemorrhage

5) Treatment for Abortion within 9 weeks?

(a) Mifepristone + Misoprostol
(b) Methotrexate
(c) Misoprostol
(d) Danazol

Ans. (a)Mifepristone + Misoprostol

METHODS OF PERFORMING MEDICAL TERMINATION OF PREGNANCY

➢ FIRST TRIMESTER

✓ **Medical**
- Mifepristone and Misoprostol, PGE,

✓ **Surgical**
- Menstrual regulation
- Vacuum Aspiration (MVA/EVA)
- Suction evacuation and/or curettage
- Dilatation and evacuation (Rapid method/Slow method)

➢ SECOND TRIMESTER

✓ **Medical**
- **Prostaglandins : GE1 (Misoprostol), 15 methylPGF2a (Carboprost), PGE2 (Dinprostone)** and their analogues (used-intravaginally, intramuscularly or intra-amniotically)

✓ **Surgical**
- Dilatation and evacuation
- Intrauterine instillation of hyperosmotic solutions
- Intra-amniotic hypertonic urea (40%), saline (20%)
- Extra-amniotic—**Ethacrydine lactate, Prostaglandins (PGE2, PGF2a)**
- Hysterotomy (abdominal)— less commonly done

➢ Most suitable method for 2nd trimester abortions is by prostaglandins

➢ **Mifepristone + Misoprostol**
- 200 mg of mifepristone is given orally on day 1 followed 2 days / 48 hrs later by

www.medicalcoupons.in

- Vaginal misoprostol 800 mcg (4 tablets of 200mcg each) is placed in the posterior fornix of vagina
-
- The combipack of 1 tablet mifepristone and 4 tablets misoprostol (i.E. 5 tablets) is approved by DGHS, govt, of india.

- **Mifepristone blocks progesterone receptors** in the endometrium which leads to disruption of the embryo, production of prostaglandins and a decrease in human **chorionic gonadotropin levels.**
- Combination of **mifepristone and misoprostol** is effective for inducing abortion within **63 days of amennorhea**

6) 35-year-old female, working in a factory presents within 8 hours after a sexual assault by a male colleague. She is at day 13 of her menstrual cycle. What should be the emergency contraceptive of choice in her case?

(a) Levonorgesterol of 1.5 mg single dose
(b) DMPA
(c) Misoprostol followed by methotrexate
(d) Ethyl estradiol

Ans. (a) Levonorgesterol of 1.5 mg single dose

The American College of Obstetricians and Gynecologists recommends a single dose of levonorgestrel 1.5 mg for emergency contraception within 72 hours of unprotected intercourse as one option. As another option, they recommend two doses of levonorgestrel 0.75 mg may be effective when taken 12-24 hours apart.

Levonorgestrel 1.5mg tablets can be used at any time during the menstrual cycle unless menstrual bleeding is overdue.

7) Identify the Image?




a. Hydrosalpinx
b. Long para ovarian cyst
c. PID
d. Bicornuate uterus

www.medicalcoupons.in

Ans: (a) Hydrosalpinx

Hydrosalpinx is **the blockage of a woman's fallopian tube** caused by a fluid buildup and dilation of the tube at its end. Most often it occurs at the fimbrial end of the tube next to the ovary, but it can also occur at the other end of the tube that attaches to the uterus.



8) A G2P1 presents with history of child with down's syndrome. Now her POG is 10 weeks. What is the best possible method of Investigation?

(a) Amniocentesis and Chorionic villi sampling
(b) Triple marker
(c) Quadraple marker
(d) USG

Ans. (a) Amniocentesis and Chorionic villi sampling

Karyotyping of cells obtained by either amniocentesis or CVS(Chorionic villus sampling) is the standard and definitive means of diagnosing aneuploidy in fetuses. The risk that a woman will give birth to an infant with Down syndrome increases with age.
Chorionic villus sampling can reveal whether a baby has **a chromosomal condition**, such as Down syndrome, as well as other genetic conditions, such as cystic fibrosis.
Amniocentesis — Amniocentesis is **used most commonly to identify chromosomal problems such as Down syndrome**. When the fetus is known to be at risk, it can detect other genetic diseases like cystic fibrosis, Tay-Sachs disease and sickle cell disease.

9) Which of the following strain of HPV is more commonly associated with cervical cancer?

(a) HPV 6
(b) HPV 11
(c) HPV 16
(d) HPV 8

Ans. (c)HPV 16

Two HPV types (16 and 18) cause 70% of cervical cancers and pre-cancerous cervical lesions. There is also evidence linking HPV with cancers of the anus, vulva, vagina, penis and oropharynx.

www.medicalcoupons.in

10) 25 year old married man came to infertility clinic, having azoospermia. normal sized testis, FSH and testosterone levels also normal. What could be the reason?

(a) Obstruction of Vas Deferens
(b) Kallman syndrome
(c) Klinifelter syndrome
(d) Impotency

Ans.(a) Obstruction of Vas Deferens

Obstructive azoospermia: This type of azoospermia means that there is a **blockage or missing connection in the** epididymis, vas deferens, or elsewhere along your reproductive tract. You are producing sperm but it's getting blocked from exit so there's no measurable amount of sperm in your semen.

In some men, **the epididymis becomes blocked**, preventing sperm from entering the vas deferens and getting into the ejaculate. A blockage can occur on one side or on both sides. If a man has an epididymal obstruction on one side, he may suffer a lowered sperm count.

11) A female in the labor room - head has not reached to the ischial spines with 3 cm cervical dilatation in 3 hours will be considered at which stage of labor?


(a) Latent phase
(b) Active phase
(c) Second stage
(e) Third stage

Ans. (a) Latent phase

Latent phase
• It is the preparatory phase of cervical dilatation • It is 8 hrs in primi and 4hrs in multi* • The ***cervix dilates only 3 cms during the latent phase****

12) Image showing a contraceptive device in uterus. The placement is best seen with:

(a) Laparoscopy
(b) Hysteroscopy
(c) Laparotomy
(d) Hysterectomy



Ans. (b) Hysteroscopy

A hysteroscopy is **a procedure used to examine the inside of the womb (uterus)**. It's carried out using a hysteroscope, which is a narrow telescope with a light and camera at the end. Images are sent to a monitor so your doctor or specialist nurse can see inside your womb.
The hysteroscope is inserted into the vagina, through the cervix, and into the uterus, allowing the surgeon to view, take a biopsy or to use special instruments to treat the underlying condition.
Hysteroscopy may also be **used to remove adhesions, polyps or a malpositioned IUD** (intrauterine device).

13) A woman with secondary amenorrhea with history of curettage for abortion. FSH 7 IU/ml. What is your likely diagnosis?

(a) Pituitary failure
(b) Ovarian failure
(c) Uterine synechiae
(d) Pregnancy

Ans. (c) Uterine synechiae

Uterine Synechiae are usually a consequence of uterine trauma caused by: **Infections**. **Prolonged use of intrauterine devices (IUDs) Post-miscarriage or postpartum curettage**.
Uterine synechiae are associated with increased risk of preterm premature rupture of membranes, placental abruption, and malpresentation. Uterine synechiae are **intrauterine adhesions around which chorioamniotic membranes wrap during pregnancy** to produce amniotic sheets or shelves.


14) Identify the Image given below?

(a) Dichorionic Monoamniotic
(b) Monochorionic Monoamniotic
(c) Dichorionic Diamniotic
(d) Monochorionic Diamniotic



www.medicalcoupons.in

Ans. (c) Dichorionic Diamniotic



15) Drug used for Endometriosis?

(a) Raloxifene
(b) Danazole
(c) Gosarelin
(d) tamoxifene

Ans. (b) Danazole

Endometriosis is an **often painful disorder in** which tissue similar to the tissue that normally lines the inside of your uterus — the endometrium — grows outside your uterus. Endometriosis most commonly involves your ovaries, fallopian tubes and the tissue lining your pelvis.

TREATMENT

- Treat all patients - regardless of clinical profile – even asymptomatic pts **– progressive disease – asymptomatic →symptomatic (30-60% patients in 1 yr.**
- **Chronic disease** with ↑ recurrence rate (even with hormonal and surgical treatment)

➢ MEDICAL MANAGEMENT
- Aim - Relief from pain

○ Pseudo pregnancy drugs - decidualization of endometrium – **COC** (Combined Oral Contraceptive), **Progesterone**

Pseudomenopause drugs - atrophy of endometrium - **Danazol, GnRH agonists**

16) A patient presented with secondary amenorrhea after 5 years of her marriage. O/E she was found to have retroverted uterus with some tenderness and swelling near the posterior wall of vagina. What is the appropriate diagnosis?

(a) Posterior wall fibroid
(b) Cervical fibroid
(c) Endometriosis
(d) Adenomyosis

Ans. (c) Endometriosis

A **retroverted uterus** (**tilted uterus**, **tipped uterus**) is a uterus that is oriented posteriorly, towards the back of the body.
A retroverted uterus is due to a disease such as endometriosis

www.medicalcoupons.in

The other factors that can cause the uterus to become retroverted. Pelvic surgery, pelvic adhesions, endometriosis, fibroids, pelvic inflammatory disease, or the labor of childbirth can change the position of the uterus to retroverted.

17) Women with ovarian malignancy, what are all the procedures that can be done except?

(a) Debulking surgery
(b) iv chemotherapy
(c) Intraperitoneal chemo therapy
(d)radiotherapy

Ans. (d) Radiotherapy

Radiation therapy in ovarian cancers has been considered an outdated concept for many years, mainly due **to toxicity and failure to show benefit in terms of survival**. Chemotherapy has been extensively used after surgery for these cancers and it has almost replaced radiation therapy as an adjuvant treatment.

18) A 14-year-old girl presented with primary amenorrhea, complains of urinary retention, pain in the abdomen. On history talking, she explains that she has been suffering from cyclical abdominal pain for last 6 months. O/E following image is seen. What is the diagnosis?



(a) Imperforate hymen
(b) MRKH
(c) Klinefelter syndrome
(d) CAH

Ans. (a) Imperforate hymen

Imperforate hymen is **when the hymen covers the whole opening of the vagina**. Imperforate hymen is the most common type of blockage of the vagina. Imperforate hymen is something a girl is born with.

www.medicalcoupons.in

19) A 30-year nulliparous lady being married for 5 years came to infertility center for treatment. She is having secondary amenorrhea for last 45 days. Doctor had done test for her ovulation in which she is ovulating regularly and normally. Under what parameters you can confirm that ovulation has taken place?

(a) High FSH
(b) Basal body temperature < 0.5
(c) High LH
(d) High progesterone

Ans. (d) High Progesterone

In conception cycles, properly timed mid-luteal progesterone levels are **over 10 ng/mL**. **Progesterone** is also often used to assess ovulation after induction of ovulation.

20) A 14-year-old girl presented with heavy bleeding per vaginum for last 10 days. What should be done to know the cause other than pregnancy test?

(a) Coagulation test
(b) LFT
(c) KFT
(d) D-dimer test

Ans. (a) Coagulation test

Coagulation tests **measure your blood's ability to clot**, and how long it takes to clot. Testing can help your doctor assess your risk of excessive bleeding or developing clots (thrombosis) somewhere in your blood vessels.

21) A woman who is lactating came to OPD 6 months after delivery for checkup and to get advice for contraception. She still wants to continue breast feeding for her child. Which is not to be adviced?

(a) Combined OCP
(b) POP
(c) IUCD
(d) Mini pill

Ans. (a) Combined OCP

Contraceptives which **contain estrogen have been linked to reduced milk supply and early cessation of breastfeeding even when started after milk supply** is well established and baby is older.

22) A 32-year-old female came for regular check up found with ovarian cyst of 6*6 cm. What's the next line of management?

(a) Laparoscopic cystectomy
(b) CA-125 follow up
(c) Laparotomy and Ovariotomy
(d) Combined OCP and follow up

www.medicalcoupons.in

Ans. (d) Combined OCP and follow up

Ovarian cysts are growths that develop on or inside the ovaries. There are several types of cysts. Functional cysts are common, not related to a disease and form as a result of ovulation. Disease-related cysts can include those that result from polycystic ovary syndrome and ovarian cancer.

Functional ovarian **cysts generally go away without treatment**. Your healthcare provider may give you medications containing hormones (such as birth control pills) to stop ovulation and prevent future cysts from forming. If you do not ovulate, you will not form functional cysts.

23) A primigravida at 24-weeks of gestation comes to the hospital with an appointment with a report of RBS of 126mg/dl. Fasting and OGTT after 2 hours was found to be 200mg/dl. On performing the USG the fetus was found to be 2.5kg in weight. What should be the best management for her?

(a) Reassure
(b) Call her to check again at 34 weeks
(c) Admit, start insulin and check for response
(d) Start insulin and send her home

Ans. (c) Admit, start insulin and check for response

Gestational diabetes mellitus (GDM), defined as any degree of glucose intolerance with onset or first recognition during pregnancy, is associated with adverse outcomes in mothers and offspring

Target glucose values in women with GDM are **≤ 95 mg per dL** (5.3 mmol per L) with fasting, ≤ 140 mg per dL (7.8 mmol per L) one-hour postprandial or ≤ 120 mg per dL (6.7 mmol per L) two-hour postprandial.

GDM Pregnant women should be managed by **Medical Nutrition Therapy (MNT)**, and **insulin therapy**/ metformin as required **and check for response**. In the postpartum period, OGTT should be repeated at 6 weeks after delivery if blood sugar <140 mg/dL, then women should be referred to NCD clinic for Post Prandial Blood Sugar (PPBS) testing annually.

24) What is the contraindication of the given device?



(a) HTN
(b) PID
(c) Obesity
(d) Anemia

Ans. (b) PID
ABSOLUTE CONTRAINDICATIONS

- Pregnancy (current, recurrent, or recent (within past 3 months))
- Pelvic inflammatory disease (PID) or sexually transmitted infection (STI)
- Puerperal sepsis
- Immediate post-septic abortion
- Severely distorted uterine cavity
- Unexplained vaginal bleeding
- Cervical or endometrial cancer
- Malignant trophoblastic disease
- Copper allergy (for copper IUDs)

www.medicalcoupons.in

25) A 40cc fetal material was detected on USG and planned for evacuation. But the catheter goes deep into the uterus without resistance. What should be the next step?

(a) Mifepristone and misoprostol
(b) Wait and watch
(c) Continue the procedure
(d) Immediate laparoscopy to be done

Ans. (d) Immediate laparoscopy to be done

Uterine perforation is a **potential complication of all intrauterine procedures** and may be associated with injury to surrounding blood vessels or viscera (bladder, bowel). In addition, uterine perforation and associated complications can result in hemorrhage or sepsis. So in order to control the bleeding immediate laparoscopy should be done.

26) Which of the following is true about true conjugate?

(a) From sacral promontory to posterior border of pubic symphysis
(b) From sacral promontory to anterior border of pubic symphysis
(c) From sacral promontory to upper border of pubic symphysis
(d) From sacral promontory to lower border of pubic symphysis

Ans. (c) From sacral promontory to upper border of pubic symphysis

THE TRUE PELVIS
It is composed of inlet, cavity, and outlet.
The Pelvic Inlet (Brim)

Boundaries

- Sacral promontory,
- alae of the sacrum,
- sacroiliac joints,
- iliopectineal lines,
- iliopectineal eminencies,
- upper border of the superior pubic rami,
- pubic tubercles,
- pubic crests and
- upper border of symphysis pubis.

Diameters

- Antero -posterior diameters:
    - Anatomical antero-posterior diameter (true conjugate) = 11cm

from the tip of the sacral promontory to the upper border of the symphysis pubis.


27) Which among the following is the most dependent part in a sitting female?

(a) Hepatorenal pouch
(b) Paracolic gutter
(c) Pouch of douglas
(d) Recto-uterine pouch

Ans. (c) Pouch of douglas

The broad ligaments reflect over the uterus, fallopian tubes, and parametrial uterine vessels and serve as the anterior boundary of the rectouterine pouch of Douglas. **The cul-de-sac** or the pouch of douglas is the most dependent portion of the peritoneal cavity and collects fluid, blood, abscesses, and intraperitoneal drop metastases.

28) Identify the usg image shown below?



a) Hydatiform mole
b) PCOS
c) Milliary TB
d) Ectopic pregnancy

Ans. (a) Hydatidiform mole

Hydatidiform mole (HM) is **a rare mass or growth that forms inside the womb (uterus) at the beginning** of a pregnancy. It is a type of gestational trophoblastic disease (GTD).
Snowstorm sign in obstetric imaging is classically seen in complete Hydatiform mole. It is characterized by the presence of many hydropic villi which gives the ultrasonographic appearance of a central heterogeneous mass having a solid, hyperechoic area and interspersed with a multitude of cystic areas and filling the entire uterine cavity.

DERMATOLOGY

1. Which sign is shown in the given picture?



A. Auspitz sign
B. Dermographism
C. Koebner's Phenomenon
D. Nikolsky's sign

www.medicalcoupons.in

Answer: C
Explanation: The Koebner phenomenon describes the appearance of new skin lesions of a pre-existing dermatosis on areas of cutaneous injury in otherwise healthy skin. It is also known as the Köbner phenomenon and isomorphic response.

2. From the given image is shown which nail finding that is the characteristic feature of Psoriasis Vulgaris?



A. Subungual keratosis
B. Auspitz sign
C. Oil drop sign
D. Onycholysis

Answer: A
Explanation: Subungual hyperkeratosis refers to **the accumulation of scales under the nail plate**, which is detached and uplifted. The nail bed often appears thickened. The condition results from excessive proliferation of keratinocytes and failure to shed off from the stratum corneum.
The nail findings in Psoriasis can be remembered by a simple mnemonic POISON
-P= Pitting
-OI=Oil drop sign
-S=Subungual hyperkeratosis
-O=Onycholysis
-N=Nail plate dystrophy

3. Identify the following Image:



A. Leishmania Cutanea
B. Lepromatous Leprosy
C. Post Kala Azar Dermatitis
D. Tuberculoid Leprosy

Answer: B
Explanation: There is saddle nose deformity and loss of eyebrows (lateral madarosis) and we can see multiple nodular lesions over the face leading us to the answer Lepromatous Leprosy.

www.medicalcoupons.in

4. Which of the following is best treatment for the following condition?



A. Valacyclovir
B. Imipenem
C. Carbapenem
D. Penicillin

Answer: A
Explanation: we can see from the image that the lesion has clear midline demarcation and is vesicular lesions on erythematous base leading us to the diagnosis Herpes zoster

- It is a reinfection of varicella virus
- Painful condition
- Appears as a vesicular lesion on erythematous base on dermatomal distribution
- Treatment is Acyclovir, Valacyclovir

5. A patient who has anti-convulsant medication presented with rashes which is shown in the following image. What could be the diagnosis?



A. Urticaria
B. Fixed drug eruption
C. Angioedema
D. Cellulitis

Answer: B
Explanation: in the question there is a history of drug intake, and the patient developed rashes after taking a drug leading us to a diagnosis of Fixed drug eruption. In angioedema and urticaria we can see a much bigger edematous swelling

- In fixed drug eruption, after taking drug a rash develops that heals with hyperpigmentation
- And after when the drug is retaken, the rash appears on the same site and this will heal with hypopigmentation.

www.medicalcoupons.in

6. Identify the image given below?



A. Carbuncle
B. Follicular abscess
C. Cellulitis
D. Erysipelas

Answer: A
Explanation: a furuncle is infection of hair follicle, single pus forming infection of a single hair follicle but whereas a carbuncle is also a pus forming infection of contiguous hair follicles

7. A patient presented with hypopigmented rash with satellite lesions as shown in the image given below?



A. BT
B. BL
C. LL
D. TT

Answer: A
Explanation: satellite lesions and finger like projections (Pseudopodia) are seen in BT type of leprosy
-In BB, we can see Swiss Cheese appearance or inverted saucer appearance
-other dermatological disorders where we can see satellite lesions are

- BT Leprosy
- Sporotrichosis
- Candidiasis

www.medicalcoupons.in

8. A person is with chronic alcoholism, which deficiency causes the following as shown in the image?



A. Riboflavin
B. Ascorbic acid
C. Biotin
D. Niacin

Answer: D
Explanation: patient is having dermatitis, diarrhea for 2 months, the patient has appearance of CASAL's Necklace. So, we conclude that it is Pellagra.
Pellagra is characterized by 4D's

- Diarrhea
- Dermatitis
- Dementia
- Death

Pellagra is caused due to deficiency of vitamin B3 or Niacin. Frequently seen in Maize/Jowar eaters due to deficiency of Tryptophan which leads to pellagra due to deficiency of niacin. Excess of leucine inhibits the conversion of tryptophan to niacin (conversion ratio of tryptophan to niacin is 60:1) in the body. Here the main culprit is Leucine.

9. Identify the following image?



A. Verrucous vulgaris
B. Elephantiasis
C. Leprosy
D. Leishmaniasis

www.medicalcoupons.in

Answer: B
Explanation: we can see unilateral swelling in lower limb,

- In elephantiasis, there will be a chronic lymphatic obstruction
- Transmitted by culex
- Once the microfilaria enters the body, they convert into adult worms and they finally destruct the lymphatics causing unilateral, non-pitting kind of an edema of the lower limbs
- Skin changes: Hypertrophy, verrucous changes and pigmentary changes

10. A truck driver with history of lesions on penis undergoing treatment presented with rashes. What is the diagnosis?




A. Secondary syphilis
B. Drug eruption
C. TEN
D. Steven Johnson syndrome

Answer: A
Explanation: these rashes are typical for secondary syphilis along with the history of lesions in the penis is leading us to the diagnosis of Secondary syphilis.

- In (Primary) 1^0 Syphilis – Hard Chancre, Painless indurated genital ulcer
- If not treated properly patient will have (secondary) 2^0 Syphilis

www.medicalcoupons.in

JUNE 2021

1. **Identify the muscle marked 'X' in the given image?**




a) Styloglossus
b) Genioglossus
c) Hyoglossus
d) Palatoglossus

Ans. B

Muscles of Tongue	
Intrensic M/s 4 pairs	**Extrinsic M/s 4 pairs**
→ Superior Longitudinal M/s	• Genioglossus
→ Inferia Longitudinal M/s	• Styloglossus
→ Transverse M/s	• Palato glossus (vagus nerve)
→ Verticle M/s	• Hyoglossus
Q1. Largest Muscle	Genioglossus
Q2. Safty muscle of tongue	Genioglossus
Q3. Not a true Muscle of tongue	Palatoglossus

2. Arrange a, b, c, d in sequence?



www.medicalcoupons.in


a) A - Cordate nucleus B - Thalamus C - Lentiform Nucleus D – External capsule
b) A-Cordate nucleus B-lentiform nucleus C-thalamus D-external capsule
c) A-Thalamus B-lentiform nucleus C-cordate nucleus D-external capsule
d) A- Thalamus B-cordate nucleus C-lentiform nucleus D-external capsule

Ans. A

3. Injury to bulgar urethra leads to extravasation of urine in all except ?
a) Deep perineal pouch
b) Superficial perineal pouch
c) Scrotum
d) Thigh

Ans. A

4. All of the following are correct except?



a) Superior thoracic artery
b) Thoracoacromial artery
c) Lateral thoracic artery
d) Posterior circumflex humeral artery

Ans. D


1) given artery : AXILLARY ARTERY
Axillary artery is divided into 3 parts by PECTORILIS MINOR MUSCLE
1ST PART
Superior thoraxic artery
2nd part
Thoracoacromial artery
Lateral thoracic artery
3rd part
Anterior circumflex humeral artery
Posterior circumflex humeral artery
Subscapular artery

www.medicalcoupons.in

Corticospinal tract crosses at

a) Mid brain
b) Pons
c) Lower medulla
d) Upper medulla

Ans. C



Functions:


Motor Functions
Skilled Voluntary monents (coriting)

6. Sutures was done to the finger during the surgery procedure. after the patient complains that he is un able to
flex the middle finger. this can be due to injury of which of the following?

a) Median nerve
b) Flexor pollicis longus
c) Flexer digitorum profundus
d) Flexor retinaculum

Ans. A

Also called : LABOURES NERVE
Root value: C5 C6 C7 C8 T1
Sign:pointing ind



Benedictis hand sign



www.medicalcoupons.in

Test for nerve: PEN TEST : Muscle- abductor pollicis brevis



7. A new born present with the congenital anomaly of cleft lip. Which is the embrynological development deftct?

a) Non fusion of palatine shelves
b) Non fussion of medial nasal process &maxillary process
c) Non fussion of lateral nasal process & maxillary process
d) Separation of nasal septem&vomer from palatine process

Ans. B

8. A patient came to the physician with complains of chest pain which is radiating to the left arm , which nerve will causes pain?

a) Musculocutaneous nerve
b) Intercostobrachial nerve
c) Medial cutaneous nerve of arm
d) Medial cutaneous nerve of forearm

Ans. B

. A patient of covid-19 requires oxygen therapy, which of the following cylinders should be used to provide oxygen to this patient?

a.  b.  c.  d. 

a) O2 [Black body with white shoulders] pressure 2000 psi (140 kg/cm^2)
b) N2O [Blue body]→ 760 psi
c) Entonox [Blue body with white shoulder] → 2000 psi
d) CO2 [Brown]

Ans. A

www.medicalcoupons.in

10. A copd patient in ICU was on oxygen therapy with mask with 2 liter of oxygen. He was found to be Hypo-
ventilating. His PCO2 on blood gas analysis was 64mmHg. Which of the following should be done?

a) Increase the FIO2
b) Decrease the FIO2
c) Start BIPAP
d) Intubate and start on ventilator

Ans. C

- ✓ ABG [Blood gas analysis] ➪po2 : < 50 (normal)
  ➪pco2 :>50 (normal) (PH < 7.3)➞ causes tachypnea
  ↓
  ➪ Then RR will be :> 35 bpm [normal : 12-20 bpm]
  ➪tidal volume :< 5ml/kg [normal : 6-8 ml/kg]

- ✓ Ventilator

Non – Invasive Invasive [Intubate]

Non-invasive ventilator :

1. Continuous positive airway Pressure (CPAP) [Alveoli always) open so that gas exchange takes place continuously
even during expiration]

2. Bilevel – positive airway Pressure [BIPAP] positive pressure during inspiration and expiration ↓

(Means better distribution of gases)

3. pressure support ventilation (PSV)

- ✓ NOTE : BI-PAP >> CPAP

Contraindications to NPPV	**Predictors of Success of NPPV**
Cardiac or respiratory arrest	Ability to cooperate
Hemodynamic instability	Good mask fit/minimal air leak
Inability to protect airway or clear secretions	Moderate hypercarbia (PaCO2 45-92 mmHg)
Uncooperative or agitated	Moderate acidemia (pH <7.35)
Facial surgery, deformity, or trauma	Clinical improvements within first two hours

www.medicalcoupons.in

11. A young patient underwent surgery under GA with O2, N2O and Desflurane. Just after extubation, the patient started to desaturate. On auscultation, there are crepitations. What can be the most common cause?

a) Pleural effusion (No crepitations)
b) ARDS (long term intubation)
c) Negative pressure pulmonary edema
d) Atelectasis (no crepitations)

Ans. C

- ✓ It happened because of laryngospasm (Upper airway obstruction) due to secretions
- ✓ ***Rx***: by giving positive pressure breath & Suxamethomium (to extubate secretions)

12. Which of the following anesthetic agent is best for cardiac patient?

a) Halothane
b) Isoflurane → it minimally ↓ CO
c) Nitrous oxide
d) Sevoflurate

Ans. B


	INHALATIONAL	INTRAVENOUS	MUSCLERELAXANT
1.Cardiac patient	Isoflurane	Ethomidate	Vecuronium
2.Asthma/COPD	Sevoflurane (max dilution	✓ Propofol (controlled) ✓ Ketamine (uncontrolled)	Any one
3.Hepatic	Sevoflurane (minimally decrease the Hepatic blood flow) Desflurane	Propofol	Cis – atracurium/ Atracurium
4.Renal		Propofol	Cisatracurium / Atracurium
5.Neuro surgeries	sevoflurane	Thiopentone	Any one can use

13. Which of the following structure is not encountered during epidural anesthesia?

a) Supraspinous ligament
b) Interspinous ligament
c) Ligament flavum
d) Dura

www.medicalcoupons.in

Ans. D



- ✓ Epidural:- 5 parts pierced
- ✓ Spinal: 5 parts + Dura &Arachorid

14. In endotracheal tube what determines the resistance?

a) Curvature
b) diameter
c) Texture
d) Length

Ans. B

- ✓ Resistance to flow → R = 8nL/Π r⁴ ⇨↑resistance →↑length , ↑radius→↑resistance ↓↓

(L- length , r – radius)

15. 7 A patient was given halothane for general anaesthesia. He developed rigidity and hyperthermia. These symptoms are most likely due to which ion?

a) Na+
b) k+
c) ca2+
d) cl-

Ans. C

- ➢ Malignant hyperthermia :
  - ✓ Malignant hyperthermia is a hypermetabolic muscle disorder due to genetic mutation in the ryanodine receptor
  - ✓ Mutation is seen most commonly in chr19
  - ✓ Mutation causes uncontrolled release of ca+ from sarcoplasmic reticulum causing intense muscle contractions and subsequent complications of hypermetabolism

www.medicalcoupons.in

16. Macrocytic anemia case with normal methyl-malonyl with increaselevel of homocysteine Diagnosis?

a) B9 deficiency
b) B12, 9 deficiency
c) Iron deficiency
d) Anemia of chronic disease

Ans. A

• Vit B12 (Cobalamine) Ilium – (termaind) Vit B9 / (Folic Acid)  Jejurum	Vit B12 Folate →DHF ↑ H_2	• Homocystein methyle transferavo • Methu melonyle Co-A mutain DHF Reductars ↑ THF → DNA & RNA H_2	• Megaloblastic anemia • Peripheral nspropathy • Megaloblastic Anemia • Neural tube defects	Only B12 – source : mineal pratus B12 if seen in Vegetarian Megablastic anemia B12 / B9 Mechameture acid / Ⓝ methameture acid

17. A newborn develops vomiting, irritability, and lethargy several days afterbirth. His mother states that his diapers smell like "burned sugar". Laboratory studies confirm the diagnosis, and the patient is started on the appropriate dietary restrictions with subsequent improvement in his symptoms. The defective enzyme responsible for this patient's condition

a) Branched chain keto acid dehydrogenase
b) Pyruvate dehydrogenase complex
c) Glutathione peroxidase
d) Glucose-6-phosphatase

Ans. A

18. A 28 year old male has undergone surgery where would debriment wasperformed and (wound was not healing which one of the following vitamindeficiency is related with delayed would healing?

a) Vitamin C
b) Vitamin B2
c) Vitamin A
d) Vitamin D

Ans. A

19. Lesch Nyhan syndrome is reported due to enzyme defect of?

a) Adenosine Deaminase
b) Xanthine Oxidase
c) Hypoxanthine Guanine Phospho Ribosyl Transferase
d) Purine Nucleoside Phosphorylase

Ans. C

www.medicalcoupons.in

20. Which of the following protein doesn't have a quaternary structure?

a) Hemoglobin
b) Collagen
c) Insulin
d) Myoglobin

Ans. D

21. A 8 month old child presents to with hypoglycemia in between fe... On examination there was hepatomegaly and liver was palpable1 cm below subcostal margin. The child also presents with lacticacidosis, ketosis and xanthomas on the buttocks. What's the most likely enzyme deficiency leading to this?

a) Branching enzyme
b) Glucose 6 phosphatase
c) C. Glycogen synthase
d) Phosphofructokinase

Ans. B

SIGNS AND SYMPTOMS of Glycogen storage disease type 1-
Glycogen accumulation in liver and renal tubule cells(hepatomegaly &renomegaly)
Hypoglycemia
Lactic acidemia
Ketosis
Hyperlipemia

Disease	Enzyme Deficient
Carbohydrates	
Type I/Von Gierke's Disease	Glucose – 6-phosphatase
Type II/Pompe's Disease	Acid Maltase
Type III/Cori's Disease	Debranching Enzyme
Type IV / Anderson's Disease	Branching Enzyme
Type V/Mc Arlde's Disease	Muscle Glycogen Phosphorylase
Type VI/ Her's Disease	Hepatic Glycogen Phosphorylase
Essential Fructosuria	Fructokinase
Fructose Intolerance	Aldolase B
Galactossemia	Galactose-1-phosphatase uridyl transferase

22. A newborn develops vomiting, irritability, and lethargy several days afterbirth. His mother states that his diapers smell like "burned sugar". Laboratory studies confirm the diagnosis, and the patient is started on the appropriate dietary restrictions with subsequent improvement in his symptoms. The defective enzyme responsible for this patient's condition

a) Aromatic acid decarboxylase
b) Phenylalanine Hydroxylase
c) Branched Chain Alpha keto acid dehydrogenase complex
d) Isovaleryl CoA dehydrogenase

Ans. C

Protein's	
Phenylketonuria (Mousy/Musty odour)	Phenylalanine Hydroxylase


Alkaptonuria / Black Urine Disease	Homogentisate Oxidase
Maple Syrup Urine Disease (Burnt Sugar Odour)	Branched Chain Keta Acid Dehydrogenase/Decarboxylase

www.medicalcoupons.in

23. A boy presents with the stretchable skin & flexible fingers which he can reach his forearm following. What's the most likely condition?

a) Osteogenesis imperfecta
b) Ehlers Danlos Syndrome
c) Chondrodysplasia
d) Marfan syndrome

Ans. B



24. Most common type of Collagen seen in fibrous cartilage of joints?

a) Type I
b) Type II
c) Type lll
d) Type IV

Ans. B

Types	Locations
Type I	Bone, skin, dentin, cornea, blood vessels, fibrocartilage and tendon
Type II	Cartilaginous tissues
Type III	Skin, ligaments, blood vessels and internal organs
Type IV	Basement membrane in various tissue
Type V	Blood vessel wasll, synovium, corneal stoma, tendon, lung, bone cartilage and skeletal muscle

25. A 73-year-old woman comes to the physician because of a 2-monthhistory of diffuse weakness and tingling of her arms and legs Neurologicexamination shows weakness of the extensor and flexor muscles of the lower extremities. Sensation to vibration and position is decreased in allextremities. This patient most likely has a deficiency of which of the following vitamins?

a) Vitamin B1
b) Vitamin B2
c) Vitamin B6
d) Vitamin B12

www.medicalcoupons.in

Ans. D

Vitamin	Name	Deficiency
Vitamin B1	Thiamine	Deficiency causes beriberi. Symptoms of this disease of the nervous system include weight loss, emotional disturbances
Vitamin B2	Riboflavin	Deficiency causes ariboflavinosis. Symptoms may include cheilosis (cracks in the lips), high sensitivity to sunlight,
Vitamin B3	Niacin	Deficiency causes pellagra. Symptoms include aggression, dermatitis, insomnia, weakness, mental confusion, and diarrhea
Vitamin B5	Pantothenic Acid	Deficiency can result in acne and paresthesia, although it is uncommon
Vitamin B6	Pyridoxine	Deficiency may lead to microcytic anemia because pyridoxyl phosphate is the cofactor for heme synthesis, depression, dermatitis
Vitamin B7	Biotin	Deficiency does not typically cause symptoms in adults but may lead to impaired growth and neurological disorders in infants
Vitamin B9	Folic Acid	Deficiency results in a macrocytic anemia, and elevated levels of homocysteine. Deficiency in pregnant women can lead to birth defects

26. Xeroderma Pigmentosa, the type of DNA repair defect

a) Nucleotide excision repair
b) Base excision repair
c) Double stranded break repair
d) Transcription coupled repair

Ans. A

DNA REPAIR MECHANISMS:

1. Exonucleolytic Proofreading
2. Nucleotide Excision Repair -
    a. Xeroderma Pigmentosa
3. Strand Directed Mismatch Repair
    a. Hereditary NON Polyposis Colon
    b. Cancer (Lynch Syndrome)
4. Base Excision Repair
5. Double Strand Break Repair
6. Transcription Coupled Repair
    a. Cockayne's Syndrome

27. A 25 old female presents to medicine OPD with acuteyearabdominal pain for which almost exploratory laparotomy was done. Heis also accompanied by neuropsychiatric symptoms and the urinesample collected turns to black with time in room temperature) Mostlikely diagnosis is

a) Porphyria cutanea tarda
b) Acute intermittent porphyria
c) Congenital erythropoietic porphyria
d) Variegate porphyria

Ans. B

www.medicalcoupons.in

Summary of Major findings in Porphyrias

Enzyme Involved[a]	Type, Class, and OMIM Number	Major Signs and Symptoms	Results of Laboratory Tests
1. ALA synthase 2 (ALAS2), EC 2.3.1.37	X-linked sideroblastic anemia[a] (erythropoietic) (OMIM 301300)	Anemia	Red cell counts and hemoglobin decrease
2. ALA dehydratase EC 4.2.1.24	ALA dehydratase deficiency (hepatic) (OMIM 125270)	Abdominal pain, neuropsychiatric symptoms	Urinary ALA and coproporphyrin III increased
3. Uroporphyrinogen I synthase[d] EC 2.5.1.61	Acute intermittent porphyria (hepatic) (OMIM 176000)	Abdominal pain, neuropsychiatric symptoms	Urinary ALA and PBG[e] increased
4. Uroporphyrinogen III synthase EC 4.2.1.75	Congenital erythropoietic (erythropoietic) (OMIM 263700)	Photosensitivity	Urinary, fecal, and red cell uroporphyrin I increased
5. Uroporphyrinogen decarboxylase EC 4.1.1.37	Porphyria cutanea tarda (hepatic) (OMIM 176100)	Photosensitivity	Urinary uroporphyrin I increased
6. Coproporphyrinogen oxidase EC 1.3.3.3	Hereditary coproporphyria (hepatic) (OMIM 121300)	Photosensitivity, abdominal pain, neuropsychiatric symptoms	Urinary ALA, PBG, and coproporphyrin III and fecal coproporphyrin III increased
7. Protoporphyrinogen oxidase EC 1.3.3.4	Variegate porphyria (hepatic) (OMIM 176200)	Photosensitivity, abdominal pain, neuropsychiatric symptoms	Urinary ALA, PBG, and coproporphyrin III and fecal protoporphyrin IX increased
8. Ferrochelatase EC 4.99.1.1	Protoporphyria (erythropoietic) (OMIM 177000)	Photosensitivity	Fecal and red cell protoporphyrin IX increased

28. Storage form of neutral lipids in our liver is

a) Cholesterol
b) Cholesterol ester
c) Triglycerides
d) Fatty acids

Ans. C

29. A 25 year male presents with hypercholesterolemia and increasedrisk of heart disease. His elder brother also had similar findings andsuccumbed to this. Workup reveals a congenital LDL receptordeficiency, The most likely defect will be?

a) Primary familial hypercholesterolemia
b) Abetalipoproteinemia
c) Familial combined Hyperlipidemia
d) Apolipoprotein E

Ans. A

30. Which of the following is a cofactor for Glutathione Peroxidase?

a) Zinc
b) Copper
c) Selenium
d) Calcium


Ans. C

www.medicalcoupons.in

7. Cofactors

Mineral	Function
Cobalt	Constituent Vitamin B12
Chromium	Potentiate the action of insulin
Flouride	Constituent of bone and teeth
Iodine	Thyroid Hormone Synthesis
Molybdenum	Cofactor for Xanthine Oxidase
Selenium	Cofactor for Glutathione Peroxidase and Deiodinase
Zinc	Cofactor for Carbonic Anhydrase Lactate dehydrogenase Alcohol dehydrogenase Alkaline Phosphatase
Manganese	Cofactor for Arginase Carboxylase Kinase Enolase

31. A 35 years old male returns back to India from China. On immigrationhe was found to have slightly elevated temperature. One of the followingscreening test was used for this?




a) CBNAAT
b) Dimer
c) Real time RT-PCR
d) CRP

Ans. C

32. Marasmus is reported in Indian Children mainly due to?
a) Due to decreased intake of carbohydrate
b) Due to decreased intake of protein
c) Due to decreased intake of fat
d) Due to decreased intake of vitamins

Ans. B

www.medicalcoupons.in



Kwashiorkor	Marasmus
It develops in children whose diets are deficient of protein.	It is due to deficiency of proteins and calories.
It occurs in children between 6 months and 3 years of age.	It is common in infants under 1 year of age.
Subcutaneous fat is preserved.	Subcutaneous fat is not preserved.
Oedema is present.	Oedema is absent
Enlarged fatty liver.	No fatty liver.
Ribs are not very prominent.	Ribs become very prominent.
Lethargic	Alert and irritable.
Muscle wasting mild or absent.	Severe muscle wasting
Poor appetite.	Voracious feeder.
The person suffering from Kwashiorkor needs adequate amounts of proteins.	The person suffering from Marasmus needs adequate amount of protein, fats and carbohydrates.

33. Which vitamin deficiency leads to cognitiveimpairment in old age?

a) Vitamin A
b) Vitamin B 6
c) Vitamin B 12
d) Vitamin C

Ans. A

Vitamin A – Retinol/Retinoic acid
Vitamin B 6 - Seizures in children
Vitamin B 12 - Methylcobalamine& cognitive impairment
Vitamin C – Iron absorbtion

Clinically important vitamins and vitamin deficiency states


Name	Sources	Deficiency	Investigations
Fat-soluble			
Vitamin A	Liver, milk, butter, fish oils	Xerophthalmia, night blindness, keratomalacia, follicular hyperkeratosis	Serum retinol
Vitamin D	Synthesised in skin	Rickets, osteomalacia	Plasma 25(OH)D/ 1,25$(OH)_2$D
Vitamin E	Vegetables, seed oils	Haemolytic anaemia, ataxia	Plasma vitamin E
Vitamin K	Green vegetables, dairy products	Coagulation disorder	Coagulation assay ± plasma vitamin K
Water-soluble			
Thiamin (B_1)	Cereals, green beans, pork	Beri-beri, Wernicke–Korsakoff syndrome	RBC transketolase, whole-blood vitamin B_1
Riboflavin (B_2)	Milk	Glossitis, stomatitis	RBC glutathione reductase, whole-blood B_2
Niacin (B_3)	Meat, cereals	Pellagra	Urinary metabolites
Pyridoxine (B_6)	Meat, fish, potatoes, bananas	Polyneuropathy	Plasma pyridoxal phosphate or RBC transaminase activation coefficient
Biotin	Liver, egg yolk, cereals	Dermatitis, alopecia, paraesthesiae	Whole-blood or urine biotin
Folate	Liver, milk	Anaemia, neural tube defects during gestation	RBC folate
Vitamin B_{12}	Animal products	Anaemia, neurological degeneration	Plasma B_{12}
Vitamin C	Fresh fruit, vegetables	Scurvy	Ascorbic acid (plasma: daily intake; leucocyte: tissue stores)

www.medicalcoupons.in

34. An alcoholic patient with neurological symptoms. There is deficiency of vitamin ?

a) B1
b) A
c) B9
d) B12

Ans. A



- B1- Thiamine
- Deficiency seen in chronic alcoholics
- Wernicks Encephalopathy
- G-global confusion
- O-opthalmoplegia
- A-ataxia
- Korsakoff psycosis
- B9-folic acid
- B12-methylcobalamine

35. A patient presented with the following lesion. Likely diagnosis?




a) Hypomelanosis of ito
b) Vitiligo (segmental)
c) Nevus of ito
d) Becker's nevus

www.medicalcoupons.in

Ans. B

Segmental vitiligo

Vitiligo vulgaris	**Most common type**
Segmental vitiligo	Seen in children Most frequent in trigeminal nerve distribution
Generalized vitiligo	Extensive lesions
Acrofacial vitiligo	Periorificially and acral parts
Lip- tip vitiligo	Lips, tip of penis, vulva, nipples
Vitiligo universalis	Widespread ,multiple endocrinopathies



36. A female patient presented with a multiple warty lesions around vulva. The lesions are gradually increasing.
On examination they are soft and fissured didn't bleed on touch. What is the lilely diagnosis?



Condyloma accuminata

a) Condyloma acuminata
b) Bowen disease
c) Condyloma lata
d) Hemorrhoids

Ans. A

Condyloma accuminata : Caused by HPV 6 and 11
Occurs on penis and vulva, in the urethra and vagina, at the uterine cervix, and in the perianal region.
May become large and invasive


www.medicalcoupons.in

Condyloma lata: Seen in secondary syphilis.

37. A female student presented with utricaria following intake of seafood.she wa[illegible]s a non-sedating drug since this
is her exam time.which is the preferred antihistamine for this patient?

a) Diphenhydramine
b) Fexophenidine
c) Promethazine
d) Chlorphendimethate

Ans. B
Fexophenidine



38. What could be [illegible]e diagnosis for this image where exclamation mark is seen in trichoscopy?





www.medicalcoupons.in

a) Alopecia areata
b) Telogen effluvium
c) Trichotillomania
d) Anagen effluvium

Ans. A
Alopecia areata
Patchy alopecia with exclamation mark hair (image)
Subtypes:
- Ophiasis→ alopecia along margin of scalp
- Alopecia totalis→ loss of hair on total scalp
- Alopecia universalis→ loss of hair on whole body

Features:
- Sharply defined alopecia
- Exclamation mark hair
- Sparing of grey /white hair
- Going white over night phenomenon
- First site affected: scalp
- Nails: hammered brass(fine stippled pitting)

Treatment: few lesions → topical minoxidil, topical corticosteroids, topical psoralen + UVA
Extensive lesion → oral corticosteroids, oral psoralen + UVA

39. A 70 years old patient develop the following lesion over face since 2 years. What is the diagnosis?



a) Basal cell carcinoma
b) Squamous cell carcinoma
c) Malignant melanoma
d) keratoacanthoma

Ans.
Basal cell carcinoma
Underlying cause :
- Albinisum
  Arsenic
- Psoralens + UVA
- UV rays

www.medicalcoupons.in

- Tar
- Xeroderma pigmentosum

C/F:
Middle aged/ elderly males
Rodent ulcer ulcerated nodule with rolled, pearly, beaded edges.
Superficial BCC
Morphoeic BCC
Pigmented BCC

Site: FACE (except superficial BCC)
Palisading arrangement
Treatment: surgical excision
Keratoacantoma: This shows keratin deposits in the center.



40. A young lady presents with lacy lesion in the oral cavity. What is the diagnosis?



a) Candidiasis
b) Psoriasis
c) Lichen planus
d) Oral leukoplakia

Ans. C
Oral lichen planus→Asymptomatic/ complain of intolerance of spicy food.
White, reticulated lacy pattern on the buccal mucosa, tongue and gingival.
Erosive gingivitis.

41. Patient presented with skin lesions and crusting. What kind of organism caused this lesion?



a) Gram positive cocci
b) Gram negative cocci
c) Gram positive bacilli
d) Gram negative bacilli

www.medicalcoupons.in

Ans. A
Impetigo →streptococcus aureus and streptococcus pyogens
Characteristic feature: honey -coloured crust




42. Which of the following is an unpaired cartilage of larynx?
a) Arytenoid
b) Thyroid
c) Cuneiform
d) Corniculate

Ans. B

*** Paired**
Arytenoid
Cuneiform
Curricular

Unpaired
- Thyroid
- Cricoid:- Complete ring cartilage
- Epiglottis



43. Which of the following laryngeal cartilage doesn't calcify with age:
a) Thyroid
b) Cricoid
c) Epiglottis
d) Croniculate

Ans. C
Epiglottis is a fibro elastic / soft cartilage will **not** ossify with age

www.medicalcoupons.in

44. A 40 years old trumpet blower has developed a right sided swelling in upper part of neck. The swelling keeps changing in size . the image shown in x-ray find the appropriate diagnosis of the condition ?



a) Throglossal cyst
b) Brachial cyst
c) Laryngocoele
d) Pharyngeal pouch

Ans. C

Laryngocele⇨

- ✓ It is the dilation of the saccule of laryngeal ventricle
- ✓ On imaging, these lesions are generally characterized as well-defined,thin-walled, fluid or air-filled cystic lesions in the paraglottic space


45. Which of the following is the m/c site of head and neck cancer in India?

a) Nasopharynx
b) Oropharynx
c) Oral cavity
d) Hypo pharynx

Ans. C

M/c site of oralcavity ca in India:-
Buccal mucosa (Gingivobuccal sulcus)
Head &Neck:- Oral cavity cancer
Over all:- Breast cancer

46. Identify the structure market:-



www.medicalcoupons.in

a) Spheroid sines
b) Frontal sines
c) Maxillary sines
d) Ethmoid sines

Ans. C
Maxillary sines is the largest sines of all 4 sines
Maxillary sines drains into MIDDLE MEATUS

47. Identify the name of this x – ray shown in given they:-



a) Cold well's view
b) Waters view
c) Schuller's view
d) Towne's View


Ans. B
* Water view →Occipto mental view (also Done with closedmouth)
*Cald well's view Occipto frontal view (Frontal & ethmoid sinus more clearly)



*** Schuller's View**

*** Towne's View**

D x – ray mastoid view




www.medicalcoupons.in

48. Which of the following is the most common craw of neonatal and paediatric inspiratory stridor?

a) Laryngesmalacia
b) Laryngeal web
c) Congenital subglottic – stenosis
d) Laryngeal cyst

Ans. A

- Laryngomalacia is **m/c congenitaialanomaly** of laryngs
- Weakness of supraglottis which leads to lnspiroy stridor shaped lary.. es om ga



49. Which of the following laryngeal muscle is t.. ab..uct.. r of vocal cord:-

a) Cricothyroid
b) Thyroartenoid
c) Posterior crico arytenoid
d) cateralcricoartenoid

Ans. C

NERVE SIPPLY

Only: 1 abductor:- Posterior cric.. a.. tenoid — Recurrent laryngeal nerve


Adductor:-
- Cricot..yroid — Superior laryngeal nerve
- L.. al c.. oarytenail
- Inte.. aryteroid — Recurrent laryngeal nerve
- Th..yroa..ytenoid (ext part) — Recurrent laryngeal nerve

Tenser mu..cri.. Cricothyroid
V..ocals (int part of thyroarytenoid)

50. What is he.. exact anatomical location of Killian's dehiscence?

a) Between sup..ior and inferior constrictor muscle
b) Between Thyropharyngeus and cricopharynges muscle fibers
c) Infer..r cor..trictor muscle
d) P..twe..n middle and inferior constrictor muscle

A.. B




www.medicalcoupons.in

51. A 4 years old child has respiratory distress, fever, inspiratory stridor, x – ray neck lateral view shows view
shows thumbsign on neck what could be causing organism?

a) Harmophilus influenza
b) Streptococcus pneumonia
c) Para influenza virus
d) Staphylococcus aureus

Ans. B

Dx: ACUTE EPIGLOTTITIS



Epiglottitis

- ✓ THUMB-PRINT SIGN is seen
- ✓ Epiglottitis may be seen radiographically on a lateral neck x-ray
- ✓ Epiglottic swelling(supraglottic) of three to four times normal is seen at the base of the tongue and appears as if someone left a thumbprint


52. 7 years old child is complaining of hearing loss and sense of full ness in the ear. The tympanometey show type
B curve what is the diagnoris in this child?

a) Chronic suppurativeotive media
b) Secretory otitis media
c) Acute suppurative otitis media
d) Otitis externa

Ans. B

M/c:- in school going child
2 – Prominent symptom CHL
Fullners in the ear

TYPE A curve: NORMAL
TYPE B curve: serous /secretor otitis media
TYPR As curve:oto Sclerosis
TYPE C curve : Carcinoma (middle ear obstruction)
TYPE Ad curve: oscoles Disruption



www.medicalcoupons.in

53. A 17 year old female patient developed earache on right side post tonsillectomy. Which nerve is responsible for the ear pain?

a) Glossophrayngealnerve
b) Vague Nerve
c) Trigeminal nerve
d) Facial nerve

Ans. A

Ear ache right side post tonsillectomy – Glossophrayngeal nerve
Nerve lies in the bed tonsil Glossophrayngeal nerve
Sensory supply to middle ear – **Tympanic branch Glossophrayngeal nerve**

54. A neonate suffering from bilateral complete choanal atresia has respiratory distress since birth. What is the cause of respiratory difficulty in this neonate?

a) Less lung Volume
b) Associated vocal cord palsy
c) Newborns are obligatory nasal breathers
d) Insufficient lung surfactant


Ans. C

55. A 40year old diabeticwoman tested covid 19 positive and she started to have nasal congestion, black crusts in nose, and blackish discoloration of skin on nasal bridge and cheek area. What could be the cause?

a) Mucormycosis
b) Allergic fungal sinusitis
c) Nasal vestibulitis
d) Maxillary sinusitis

Ans. A





www.medicalcoupons.in


zygomycetes→seen in terminally ill patients
R → Rhizopus
A→Absidia
M→Mucor
Tissue biopsy :H&E stain (aseptate hyphae, obtuse angled)
Biopsy material on SDA→ cultural growth + LPCB→ Rhizopus
Mucor mycosis is commonly seen in:
DM
Frequent blood tansfusion
Chronic dialysis

56. A patient of Presbyacusis has been found to have high frequency sensorineural hearing loss on audiometry.
Which of the following part of inner ear is affected in this patient?
a) Apexof cochlea
b) Basal turn of cochlea
c) Utricle
d) Saccule

Ans. B

57. Which of the following arteries is the main blood supply of Tonsil?
a) Descending palatine artery
b) Ascending palatine artery
c) Tonsillar branch facial artery
d) Ascending pharyngeal artery

Ans. C



58. A 50 year old diabetic patient has presented with severe ear ache and blood stained eardischarge. The examinationshows granulations in the external auditory canal What is the possible diagnosis in this case?
a) Malignant otitis externa
b) Diffuse otitis externa
c) Otomycosis
d) Chronic suppurative otitis media

www.medicalcoupons.in

Ans. A




Young diabetic : black nose – MUCORMYCOSIS
Old diabetic : EAR – MALIGNENT OTITIS EXTERNA

m/c/c –PSEUDOMONOS

59. A child aged 2 years, presented with severe SNHL was prescribed hearing aids, but showed no improvement. Which is the next line of management?

a) Fenestration surgery
b) Conservative
c) Stapes mobilization
Cochlear implant

Ans. D



Cause of death in Judicial hanging?

a) Asphyxia
b) Brain hypoxia
c) Cervical vertebral fracture
d) Vagal inhibition


www.medicalcoupons.in

Ans. C

Judicial hanging ⇨Hang man's # ⇨$C_2 - C_3$ vertebra#

C 2:Pars inter articular part of vertebra





61. Cherry red colour post-mortem staining is associated with which poisoning ?

a) H2s
b) CO
c) Nitrate
d) Hydrogen cyanide

Ans. B

CO: Cherry red





H2s: Bluish greennitrate :redish brown:
Hydrogen cyanide:BRIGHT RED

62. A baby patient comes to casualty with history of unknown poisoning by some seed/drug, on examination
patient presented with Dilated pupil, Dry mouth, Dilirium and decreased, secretions, skin scratches,
Hallucination poisoning should the Dr suspect

a) Dhatura
b) Poppy seeds
c) Mushroom
d) Cannabis

www.medicalcoupons.in

Ans. A

- Dathura contains anti-cholinergic substances that cause these symptoms
- Mushroom contains cholinergic substances
- Poppy seeds contains opium- morphine like substance

Symptoms Resemble Atropin poisoning
↓
Active principle(**hyoscine**) in Dhatura poisoning



63. A person died due to starvation, which is correct in such case during autopsy?

a) Enlarged heart
b) Enlarged stomach
c) Distended gall bladder
d) Expanded lungs

Ans. C

When a person have his food. When if reaches Duodenium it stimulates to release of **CCK** which helps in contraction of gall bladder and Bile is released.

- As the person is in starvation:- bile secritons will not enter to deodinum so, gall bladder distended.
- All visceral organ size ↓↓, Except BRAIN
- Lungs will get collaps
- Cause of death is starvation:- Circulatory failure

↓

d/t:- Brown atrophy of heart

64. A person has committed a criminal offence, caught by police & later brought to a criminal court but the court directs police to transfer the case to Juvenile courts if it was proved that accused is of _____ years of age?

a) 21
b) 1[illegible]
c) 25
d) [illegible]

Ans. B

Child < 18 ⇨ according to Juvinile Justice Board.
Abdament in act:- any Juvinal of 16-18 years do's Heanes offence like (Rape, murder) will come under adult.


www.medicalcoupons.in

65. In Barberious test, yellow colourneede shaped crystals are seen under microscope. This is done to detect
which stain?

a) Sputum
b) Semen
c) Blood
d) CSF


Ans. B

Boy friend:



1) Barberious
2) Florena = Semen

Blood:-T-Tetamans
T – Takayama



66. A farmer was brought to casualty with pinpoint pupil increased secretions and garlicky ordor, which drug to
be used by physiatrics for treatment?

a) Oximes
b) Atropine
c) N-acetylcysteine
d) Physostigmile

Ans. B

- Organophosphate poisoning(OP): pinpoint pupil,garlicodour,increase secretions
- DOC for OP: atropine
- DOC for angiodiazepine: Flumazenil
- Paracetmolpoisoning : N-acetylcysteine is used
- Note: oximes are used in OP but it is not the drug of choice

OPC – poisonics (y? :- farmer, pinpoint pupil, garlicky order)

www.medicalcoupons.in

↑↑↑, Ach at Receptor level
Controlled by:-Atropine

→Physostigmine:- Doc:- DHATHUR poisoning
Oximes :- may be used but in 2nd choice
N – acetyl cysteine:- Antidote :- for Paracetmolo poisoning / Acetamrinaphin

67. A childaccidentaly consumes iron syrup used by the mother, Antidote to be used in such poisoning is?
a) BAL
b) EDTA
c) Desferrioxamine
d) Trientene

Ans. C
Chelatorfor :copper poisoning &wilsons disease: TRIENTENE

68. Gutter fracture of skull bone is associated with impact of?
a) Sharp object
b) Blunt object
c) Fall from height
d) Bullet

Ans. D
Tanguntial gun shot:- # of only outer table of skull



69. In an unconscious patient with head injury requiring emergency Sx. If there are no relatives to give consent the doctor should?
a) Operate without consent
b) Should not operate
c) Wait for the arrival of relatives
d) Take consent from the hospital authority

Ans. A
IPC: **92** state DOCTOR**CAN**operate with out consent in case of emergency

70. A 40-years old patient was brought by police in unconscious state to casualty with history of unknown poisoning, on examination patient had needle track ulcers over upper limbs, Bugs crawling on skin, patient had Hallusions with scratch marks, increased blood pressure, increased heart rate and respiratory rate of 30/ minute, which poisoning should Dr. suspect?
a) Cocaine
b) Heroin
c) Cannabis
d) LSD

www.medicalcoupons.in

Ans. A

Option we can easily eliminate by:
HEROIN : respiratory rate decreases
CANNABIS: BP decreases (due to :vomiting&diatthia)
LSD : is taken through oral rout so percutianeous ulcers are not found over upper limb

71. Partial hanging mimics like ligature striangulation, How?

a) Hyper extended neck
b) Transverse mark over neck
c) Hand touches ground
d)Glow & stocking marks

Ans. B

LIGATURE MARK DIFFERENCE BETWEEN

PARTIAL HANGING COMPLETE HANGING



72. First tooth to appear?

a) Upper lateral incisors
b) Lower lateral incisors
c) Upper central incisors
d) Lower central incisors


Ans. D

Lower central incisors :6-10 months
Upper central incisors :8-12 months
All teeth appear by 2.5 years

73. Which of the following immunoglobulins is a pentamer?


a) IgM
b) IgA
c) IgE
d) IgG

Ans. A



www.medicalcoupons.in

74. Identify the organism:



a) HIV
b) Herpes
c) Hepatitis B
d) Adenovirus

Ans. A

Family :Retroviridae
Subfamily: lentivirnae
It is a ssRNA virus
3 structural gene
→gag : matrix protein
→pol: polymerase
→Env : envelop, Gp41,Gp120
HIV affects CD4+ve T-cells by coreceptors CXCR4
HIV affects macrophages by coreceptors CCR5
Most sensitive test for HIV: ELISA
Most specific test for HIV: HIV RNA PCR/WESTERN BLOT
Mother to child transmission –30%
Blood transfusion –90%
Needle stick injury—0-5%

75. A 5 year old child presents with fever, bull's neck as shown in the image. Identify the disease.




a) Mumps
b) Lyme's disease
c) Diphtheria
d) Measles

www.medicalcoupons.in

Ans. C

Diphtheria(klebs loeffler's bacilli)
Gram positive bacilli
Chinese letter pattern/cuneiform arrangement is seen.




Staining—ponder's, albert's, neisser's (contents of albert's stain: toluidine blue,malachite green, glacial acetic acid, kl + l_2)
Pseudomembrane formation is seen → greyish site →M/C site faucial (tonsillar)
Bull's neck(neck edema)
Chronic complication : myocarditis, polyneuropathies
Dx:
LSS(loeffler's serum slope) ,culture media,enriched media(used for early diagnosis of diphtheria)
Potassium tellurite agar(best selective media)
Macleod's media (diphtheria toxin)


76. Most common viral agent causing pneumonia in < 1 year of age:

a) RSV
b) Parvovirus
c) HHV-6
d) Influenza virus

Ans. A

RSV(Respiratory Syncitial Virus) –Rx in <1 year of age.
Diagnosis :cytopathic effect (syncytium form)—MN giant/mØ.
DOC—Ribavarin.

Safety-pin appearance is seen in



Yersinia pestis

Explanation: yersinia pestis causes Plague
Other organisums which show safety-pin appearance:

www.medicalcoupons.in

1) Klebsiella granulomatis
2) Burkholderia mallei
3) Burkholderia pseudomallei
4) Vibrio parshemolyticus
5) Haemophilus ducreyi

77. The agent of the disease shown in the image is not found in which body fluid?



a) Saliva
b) Urine
c) Fluid from the leg
d) Ascitic fluid

Ans. A


78. Which among the following infection doesn't form a membrane like covering over the tonsils?

a) Diphtheria
b) Infectious mononucleosis
c) Streptococcal tonsilitis
d) Ludwig angina

Ans. D

Ludwig's angina is a rare skin infection that occurs on the floor of the mouth, underneath the tongue. This bacterial infection often occurs after a tooth abscess, which is a collection of pus in the center of a tooth.

79. A female with the following symptoms of sudden high fever, associated with headache and vomiting along with nuchal rigidity, presented to the hospital. CSF analysis of this patient showed the growth of this organism aa gram negative diplococci shown in the image,identify the organism:




a) Neisseria meningitidis
b) Streptococcus pneumoniae
c) Staphylococcus aureus
d) Mycobacterium tuberculosis

www.medicalcoupons.in

Ans. A

Neisseria meningitidis: Gram negative cocci(catalase/oxidase positive)
Ferrments glucose and maltose(capsulated)
Cultured media : thayer martin media, muller hinton agar
Transport media : stuwart amile media
Complications: water-house friderichsen syndrome(B/L adrenal haemorragic necrosis)
Deficiency of C5-C9(MAC): Prone for neisserial infections.
DOC: Ciprofloxacin(carrier), ceftrioxan(cases)


80. A covid patient presented with fever along with nasal congestion and unilateral headache. Later he developed
facial pain and numbness along with blackish discharge, KOH amount reveals nonseptate hyphae at right
angkes or obtuse angle, what is the most probable diagnosis?



a) Mucormycosis
b) Aspergillosis
c) Bacterial orbital cellulitis
d) Cavernous sinus thrombosis

Ans. A

81. Identify the organisum given in the smear image.




a) Plasmodium Falciparum
b) Plasmodium vivax
c) Plasmodium ovale
d) Plasmodium malariae


www.medicalcoupons.in

Ans. A

Plasmodium vivax	Plasmodium falciparum
Young aged RBC involved	All age RBC are involved
↑ in size of RBC	Normal size RBC
Schuffner's dot	Maurer's dot
Gametocyte stage	Banana shaped gametocytes
Single ring with in RBC	Multiple ring within RBC
Reddish brown pigment	Dark brown pigment



82. Identify the organisum causing diarrhea



a) Rotavirus
b) Giardia lamblia
c) Cryptosporidium
d) Adenovirus

Ans. B

Giardia lamblia → fatty foul smelling diarrhea and fat malabsorption.
Fecal-oral route transmission
Dx: string test
Falling leaf motility
4 pairs of flagella , tennis racket shape
DOC: Metronidazole, tinidazole

www.medicalcoupons.in


83. Orchitis is an important complication of which infection?

a) Mumps virus
b) Adeno virus
c) Polio virus
d) Pox virus

Ans. A

Myxo virus
Period of communicability: 4-6 days before and 1 week after.
Secondary attack rate: 86%
Mode of transmission : droplets
C/F: pain, swelling in parotid region, earache.
Complication: aseptic meningitis(children), orchitis(M/C extra salivary manifestation in adults), leading cause of pancreatitis in children.
Vaccin: jerry-lynn stain, live attenuated

84. An immunocompromised patient came to the hospital with severe diarrhea. Flectal swab revealed the following image. Identify the causative organism.



a) Cryptosporidium parvum
b) Rotavirus
c) Giardia lamblia
d) Adenovirus

Ans. A

Cryptosporidium parvum → causes diarrhea in AIDS patients.
Oocysts and cyst are infecti
Dx:Acid fast oocyts
DOC: Cotrimoxazole

85. A 10 year old child is diagnosed with amblyopia in one eye. What will be the best treatment ?

a) Observation
b) Penalisation
c) Occlusion
d) None of the above

Ans. C

Occlusion of the good eye to force the use of lazy eye.



www.medicalcoupons.in


86. A 3 months old baby presents with mucous exuding from eye on pressing the lacrimal sac. What treatment should be given?

a) Lacrimal sac massage
b) Syringing
c) Probing
d) DCR

Ans. A

Lacrimal sac /Pediatric massage/Crigglar sac massage: only till 1 year






1-2 years: syringing
After 2-3 years : probing (bowman's lacrimal probe)
After 3 years : DCR

87. Which lesion of the visual pathways produces a macular sparing field defect?

a) Optic chiasma
b) Occipital lobe
c) Optic nerve
d) Optic tract

Ans. B

Occipital lobe is supplied by 2 arteries: posterior cerebral artery and middle cerebral artery in which posterior cerebral artery is more often not damaged. And middle cerebral artery still supplies so we see macula is spared.



88. A child with history of malnutrition is examined and the following finding is noted. What is the most likely diagnosis?


a) Megalocornea
b) Buphthalmos
c) Anterior staphyloma
d) Corneal degeneration



www.medicalcoupons.in

Ans. C

Due to malnutrition corneal perforation occurs which further leads to staphyloma.
Protrusion of the outer tunic eyeball w.r.t equator :
Anterior to equator —anterior staphyloma
Posterior to equator – posterior staphyloma
Along the equator – equatorial staphyloma



89. A young male complains of watery eyes and itching and on eversion of upper eyelid the following picture is seen. What is the diagnosis?



a) Bacterial conjunctivitis
b) Trachoma
c) Vernal keratoconjunctiviti
d) Angular conjunctivitis

Ans. C

In question-- Itching is mostly seen in allergic conditions and we see cobble stone appearance in image, above features are seen in vernal keratoconjunctivitis.
Other options:
Bacterial conjunctivitis : no itching is seen
Trachoma: follicle formation is seen
Angular conjunctivitis : seen in chronic malnurised patients.

90. Which of the following is the gold standard measuring IOP?

a) Schiotz
b) Tonopen
c) Pulse air tonometer
d) Goldmann applanation tonometer



Ans. D

Goldmann applanation tonometer is used to measure IOP by touching the cornea.


www.medicalcoupons.in

91. What is the treatment of choice for recurrent pterygium?

a) Simple excision
b) Observation
c) Excision with conjunctival auto graft
d) Treatment with Mitomycin C

Ans. C



Excision with conjunctival auto graft



92. A lady present with a sudden painful loss of vision which worsens on eye movement. RAPD is present on perimetry a central scotoma is revealed. What is the diagnosis?

a) Optic neuritis
b) Papilledema
c) Retinoblastoma
d) CRVO

Ans. A

	
In optic neuritis (blurred disc margin)	**In retrobulbar neutritis(clear cut margin)**

Other options:

Papilledema –no loss of vision.
Retinoblastoma – causes leukocoria, squint, glaucoma.
CRVO - painless loss of vision.

93. A patient presents with history of trauma with vegetable matter a few days back, with pain and photophobia.
What is the most likely cause?

a) Herpes simplex keratitis
b) Acanthamoeba keratitis
c) Bacterial keratitis
d) Fungal keratitis

www.medicalcoupons.in

Ans. D



Other options:
Herpes simplex – painless ulcer
Acanthamoeba keratitis – painfull but due to contact lens

94. A child presents with proptosis, pain and loss of vision [illegible], the following picture is seen. What is the most likely diagnosis?




a) Orbital cellulitis
b) Stye
c) Preseptal cellulitis
d) Blepharitis

Ans. A
Orbital cellulitis— proptosis, chemosis, congestion, pupil dilation and inflammation are seen in image.
Generally orbital cellulitis starts with a simple cold and cough→ ethmoid sinusitis →medial wall of orbit.

95. A boy gets punched in the eye. Which of the following is the most likely occurrence?
a) Subluxation of the lens
b) Inferior orbital wall fracture
c) Medial wall fracture
d) Lateral wall fracture

Ans. B
Blow out fracture-- Inferior wall gets fractured more often because it gets supported by maxillary sinus which is cavernous.
(Tear- drop sign)

www.medicalcoupons.in



96. A patient with VA >1/60, but <3/60 in better eye. What type of blindness does he have?

a) Low vision
b) Economic blindness
c) Social blindness
d) Manifest blindness

Ans. C

Comparison of WHO and NPCB definitions


WHO-ICD	VISUAL ACUITY	NPCB
LOW VISION		
Category (1)	<6/18 - 6/60	Low vision
Category (2)	<6/60 - 3/60	Economic / Legal blindness
BLINDNESS		
Category (3)	<3/60 – 1/60	Social blindness
Category (4)	<1/60-perception of light	Manifest blindness
Category (5)	No perception of light	Absolute blindness

97. In which of the following, V-Y procedure is used?


a) Ectropion
b) Entropion
c) Symblepharon
d) Ankyloblepharon

Ans. A


Ectropion

Ectropion	**Outward turning of lid margin.** 

www.medicalcoupons.in

Entropion	**Inward turning of lid margin** 
Symblepharon	**Adhesion between palpebral and bulbar conjunctiva** 
Ankyloblepharon	**adhesion of the edges of upper eyelid with the lower eyelid** 

98. A patient presents with history of DM of more than 20 years duration and painless loss of vision with the given fundus findings. What is the most likely diagnosis?



a) NPDR
b) PDR
c) CRVO
d) CRAO

Ans. A
NPDR
Other options:
PDR— not the answer because no new blood vessels seen in the image.
CRVO—no tomato splash appearance in image.
CRAO—could see o*cular spot.*


www.medicalcoupons.in

99. A lady presents with ptosis, on eating and sipping of fluids her ptosis decreases. What is the most likely diagnosis?

a) Mechanical ptosis
b) Horner's syndrome
c) Complicated ptosis
d) Blepharophimosis syndrome

Ans. C

Complicated ptosis—Marcus Gunn Jaw- winking syndrome: a retraction of the ptotic lid in conjunction with stimulation of the ipsilateral pterygoid muscle



100. A newborn child presents with watering and photophobia. O/E the following picture is seen. What is the diagnosis?




a) Buphthalmos
b) Congenital cataract
c) Megalocornea
d) Congenital dacryocystitis

Ans. A

Other options:

Congenital cataract—absence of water and megalocornea
Megalocornea—no watering
Congenital dacryocystitis—absence of megalocornea

101. A patient presents with complicated cataract and on slit lamp findings are shown in the given image. What is the type of cataract seen in this case?



www.medicalcoupons.in

a) Anterior subscapular cataract
b) Nuclear cataract
c) Posterior subscapular cataract
d) Cortical cataract

Ans. C


102. Refractive error found here is:



a) Myopia and astigmatism
b) Hypermetropia
c) Myopia
d) Astigmatisum

Ans. A
Myopia and astigmatisum due to irregular surface

103. All of the following are true about below procedure, except:




a) It can be done with patient being in any position
b) There is no stereopsis
c) Can be used in cases of hazy ocular media
d) The retinal periphery cannot be examined.

Ans.
The image shown here is direct ophthalmoscopy.

www.medicalcoupons.in

104. A patient presents with ocular motion defect. The position of the eyeball is shown in the image below.
Identify the nerve involved in following defect.




a) Occulomotor nerve
b) Optic nerve
c) Trochlear nerve
d) Abducens nerve

Ans. A

105. Person with one month H/o hip pain, true limb shortening and diagnosed as a case of tuberculosis. Which stage is it ?
a) Late arthritis
b) early arthritis
c) early synovitis
d) advanced arthritis

Ans. A
- ✓ Stage 1 : synovitis stage : hip in safe posture (i.e abduction and external rotation)
- ✓ Stage 2 : early arthritis : hip is in adduction and internal rotation
- ✓ Stage 3 : late arthritis : joint space distructed (TRUE LIMB SHORTENING)
- ✓ Stage 4 : advance arthritis : acetabulum will move freely

106. X-ray and clinical photo is given, which nerve is most commonly damaged ?



a) Deep peroneal nerve
b) Sciatic nerve
c) Femoral nerve
d) Saphenous nerve

Ans : B
Posterior hip dislocation:
- ✓ FADIR ⇨flexion,adduction,internal rotation
- ✓ Limb shortening

www.medicalcoupons.in

- ✓ Sciatic nerve damage
- ✓ Lesser trochanter is not seen clearly

Anterior hip dislocation:

- ✓ FABER⇨flexion,abduction, external rotation
- ✓ Femoral nerve damage
- ✓ Lesser trochanter is seen clearly



1)POSTERIOR HIP DISLOCATION



2)ANTERIOR HIP DISLOCATION


107. Hormones and endogenous substances that are involved in regulation of calcium balance in the body are ?

a) Vitamin D,PTH,Insulin
b) Vitamin D,PTH, Calcitonin
c) Vitamin D,Calcitonin,TSH
d) Vitamin D,PTH,TSH

Ans. B

	ca +	po4
I. Vit d	↑↑	
II. PTH	↑↓	
III. Calcitonin	↓↓	

108. A patient from bihar present with teeth and skeletal image as shown. most likely cause is ?




a) Osteopetrosis
b) Osteoporosis
c) Osteosclerosis
d) Osteomalacia

Ans. C

- ✓ Dental flourosis
- ✓ Normal value of flourosisis : 0.5 - 0.8 ppm
- ✓ Upper limit : 1- 1.5 ppm


www.medicalcoupons.in



109. A alcoholic patient slept with compression. Around hi[illegible]m region and had injury to radial nerve. Most likely
this is which type of injury ?

a) Neuropraxia
b) Neurotmesis
c) Axonotmesis

Ans: A





110. A 40 year old [illegible]male patient having history of bee allergy came to the emergency room with complaints o[illegible]
swelling of face [illegible]nd dyspnea after a bee sting . Which of the following inflammatory mediators plays a major
role

a) S[illegible]roto[illegible]in
b) B[illegible]ady[illegible]in
c) His[illegible]amine
d) P[illegible]aglandin

An[illegible] C

www.medicalcoupons.in






111. When stained with Congo red and visualized under the polarising microscopewhich of the followingsubstance would show apple green birefringence


a) Lipids
b) Amyloid
c) Collagen
d) Calcium

Ans. B



www.medicalcoupons.in

112. 24year oldfemale came with complains ofneurological deficits. Examination reveals pallorHerblood
picture is given below. What is the most like diagnosis



a) Folate deficiency
b) Vitamin B12 deficiency
c) Thalassemia
d) Iron deficiency anemia


Ans. B

Vitamin B12 deficiencyanaemia occurs when a lack of **vitamin B12** causes the body to produce abnormally large **red blood cells** that cannot function properly.

113. A 35 year old lady presented with slow growing thyroid swelling for the past 10 months . Histological
examination was done and the microscopic image of the biopsy had the presence of Orphan Annie eye nuclei.
What is the patient suffering from

a) Follicular carcinoma of thyroid
b) Papillary carcinoma
c) Anaplastic carcinoma
d) Medullary carcinoma

Ans. B




Orphan Annie eye appearance in Papillary Carcinoma thyroid

www.medicalcoupons.in

114. Supravital staining is used for ?

a. Red blood cells
b. Platelets
c. Reticulocytes
d. White blood cells

Ans. C



RETICULOCYTES

- **Reticulocytes** are young , premature, non nucleated red blood cells, contain reticular material (RNA) that stain gray blue.
- Reticulum is present in newly released blood cells for 1-2 days before the cell reach its full mature state.
- RBCs have six stages :
  1) Pronormoblast
  2) Basophilic normoblast
  3) polychromatophilic normoblast
  4) orthochromic normoblast
  5) Reticulocyte
  6) mature red blood cell

PRINCIPLE

- Whole blood is incubated with **supra-vital staining** (new methylene blue). The vital stain causes the ribosomal and residual RNA to coprecipitate with the filamenting mitochondria and ferritin masses in living young erythrocytes to form dark-blue clusters and filaments (reticulum).
- Smears of this mixture are then prepared and examined. The number of reticulocytes in 1000 red blood cells is determined. This number is divided by 10 to obtain the reticulocyte count in percent.

115. Small or medium sizedsterile vegetations found in either or both sides heart valve commonly associated with SLE patients is characteristic of which of the following disease?

a) Non bacterial thrombotic endocarditis
b) Rheumatic heart disease
c) Infective endocarditis
d) Libmann Sachs endocarditis

Ans. D

www.medicalcoupons.in






Figure 12-24 Comparison of the four major forms of vegetative endocarditis. The rheumatic fever phase of rheumatic heart disease (RHD) is marked by small, warty vegetations along the lines of closure of the valve leaflets. Infective endocarditis (IE) is characterized by large, irregular masses on the valve cusps that can extend onto the chordae (see Fig. 12-25A). Nonbacterial thrombotic endocarditis (NBTE) typically exhibits small, bland vegetations, usually attached at the line of closure. One or many may be present (see Figure 12-26). Libman-Sacks endocarditis (LSE) has small or medium-sized vegetations on either or both sides of the valve leaflets.

116. A person with hepatic jaundice, deranged liver function tests and presence of a clinical finding as shown alongside



a) Huntington chorea
b) Wilson's disease
c) Leigh syndrome
d) Hemochromatosis

Ans. B



www.medicalcoupons.in

117. A teenager presents with features of liver disease, tremor and poor incoordination, mask like facies, and
psychiatric symptoms. Which of thefollowing is implicated in the pathogenesis of this condition?
a) ATP 7A gene on chromosome 13
b) ATP 7B genechromosome 13
c) ATP 7C gene on chromosome 13
d) ATP 7B gene on chromosome 6


Ans. B

HNF 1-alpha- Hepatic Adenoma
ATP7B- Wilsons disease
Fibrillin- Marfan syndrome
COL1A1- Osteogenesis imperfecta


WILSON DISEASE-ATP 7B genechromosome 13
HEMOCHROMATOSIS-HFE genechromosome 6P
ALPHA1-ANTITRYPSIN DEFICIENCY- SERPINA 1 genechromosome 14
CYSTIC FIBROSIS – CFTR genechromosome 7


118. A child presented with history of pallor with fatigue, petechial bleeding and fever since last 7 days. On
examination he was havingenlargement of liver and spleen with sternal tenderness. Which of the following
is the likely explanation for the symptoms in the child?
a) Aplastic anemia
b) Chronic myeloid leukemia
c) Acute lymphoblastic leukemia
d) Acute myelogenous leukemia

Ans. C









www.medicalcoupons.in

119. A person is working with 30 year history of working in cardboard manufacturing and develops
breathlessness and likely to be associated withmottling in lungs is

a) Bysinossis
b) Bagasossis
c) Asbestosis
d) Nasopharyngeal carcinoma

Ans. B

Bagassosis is occupational disease of lung caused by inhalation of Bagasse or sugarcain dust

- First reported in India in cardboard manufacturing firm near Kolkata
- Asbestosis: Cement, roof tiling
- Byssisinosis: Cotton industries


Agent	Disease	Typical Exposure
Coal Dust	Anthracosis	Coal miners
Asbestos	Asbestosis	Mining, installation of insulation
Silica	Silicosis	Sand blasting, Stone cutting
Bagasse	Bagassosis	Cardboard, paper and sugar industry
Cotton, hemp	Byssinosis	Textile manufacturing

120. A tall male teenager presented with gynecomastia . On examination,absence of secondaryhe has average
cognition characteristics a deep voice, beard, and male distribution of pubic hair. His genetic analysis was
performed and he had the presence of abarrbodv. What is the likely diagnosis of this condition?

a) Turner syndrome
b) Klinefelter syndrome
c) Swyer syndrome
d) Down syndrome


Ans. B

Klinefelter-xxy-and-turner-xo-syndrome



www.medicalcoupons.in

121. A 16 year old boy had abdominal discomfort on the left side andhe went to his phy ician. His father had a
similar condition during his teenage and later developed some GI malignancy. He th n under went a
colonoscopy which is shown alongside. What doyou conclude about his condition?



a) Familial adenomatous polyposis
b) Cancer colon
c) Ulcerative colitis
d) Lynch syndrome

Ans. A
FAP

- **>100 polyps are seen.**
- **Autosomal dominant**
- **Mutation in APC gene (5q21)**

	Genetic Mutati n	*Clinical Findings*
Adenomas		
Familial adenomatous polyposis (FAP)	APC gene	1. 100s-1000s of colonic polyps that inevitably become malignant
Gardner syndrome	APC g...	1. Adenomatous gastrointestinal polyps 2. Dental and ocular abnormalities 3. Soft tissue tumors, osseous abnormalities 4. Extraintestinal malignancies
Turcot syndrome	...or ...match-repair ...nes (hMLH1 or ...PM...2)	1. Colonic adenomatous polyps 2. Brain tumors
Hamartomas		
Peutz Jeghers syndrom...	STK11 or LKB1	1. Intestinal hamartomatous polyps 2. Mucocutaneous melanotic pigmentation 3. Pancreatic, breast, ovarian, testicular, thyroid malignancy
Juvenile polypos... ...ndrom...	SMAD 4 or BMPR 1A	1. Multiple juvenile polyps within the gastrointestinal tract 2. Increased risk of colon, stomach, and duodenal cancer
Cowden sy...drome ...nd Ban...evan-...ley-Ruvalcaba sync...m	PTEN	1. Gastrointestinal hamartomas 2. Mucocutaneous lesions 3. Macrocephay and neurologic abnormalities 4. Breast, thyroid, and genitourinary cancers

1... 2. A 30 year male patient has a complaint of a painless testicular mass. His blood sample ...ealed non elevated
alp...a fetoprotein levels. Which of the following is the most likely cause of his testicular mass?
a) Yolk sac tumor
b) Classical seminoma
c) Teratoma
d) Choriocarcinoma

www.medicalcoupons.in

Ans. B

Seminoma

- The commonest variety of testicular tumour
- Adults are the usual target (4th and 5th decade)
- Right > Left Testis
- Starts in the mediastinum: compresses the surrounding structure.
- Patients present with painless testicular mass
- 30 % have metastases at presentation, but only 3% have symptoms related to metastases
- 90% of patients stain for PLAP.
- AFP is normal

- Classification
  a) Classical – 85%
  b) Anaplastic – 5-10%
  c) Spermatocytic

123. An elderly male has been working in the dye industry for 40years.He develops somefrequency and urgency,
Bladder symptoms like hematuria. What is the likely subtype of the conditionpresent in him?
a) Transitional cell cancer
b) Adenocarcinoma
c) Squamous cell cancer
d) Small cell cancer

Ans. A

Box I. confirmed or suspected risk factors for transitional cell carcinoma	
Smoking	Increases risk up to five times
Analgesics	Mainly associated with renal pelvis transitional cell carcinoma, but also bladder tumors
Occupation	Workers in aniline dye, rubber and chemical industries due to exposure to β-naphthylanine (which in the liver is converted to carcinogen that must be activated in the bladder). These workers need regular bladder checks.
Cyclophosphamide	Can cause bladder cancer in the long term (although used for cancer treatment)
Schistosomiasis	Can cause chronic inflammation and metaplasia (squamous) of the bladder mucosa (leading to squamous cell carcinoma)
Chronic infections/inflammation	Some authorities believe that any chronic inflammatory may predispose to cancer

124. Which of the following is the likely diagnosis in a lady with extremely low TSH and high T3 and T4?
a) Subclinical hypothyroidism
b) Subclinical hyperthyroidism
c) Graves disease
d) Reidel thyroiditis

www.medicalcoupons.in

Ans. C

Table 1: Differences Between Hashimoto's Thyroiditis and Graves' Disease

Autoimmune disorder	Hashimoto's thyroiditis[2,4]		Graves' disease[2,5]	
Antibodies	• Antithyroid peroxidase antibodies • and/or Antithyroglobulin antibodies • (Normally no anti-TSH receptor antibodies)		• Anti-TSH receptor antibodies • (Antithyroid peroxidase antibodies (in 80% of patients)) • (Antithyroglobulin antibodies (in 50%))	
Thyroid state	Progresses into hypothyroidism		Hyperthyroidism	
Serum TSH	Variable	Progresses into hypothyroidism	Low	Hyperthyroidism
Free T3	Variable		High	
Free T4	Variable		High	
Clinical features	• Hypothyroid, • Euthyroid, • or (more rarely) Hyperthyroid		• Hyperthyroid signs and symptoms with sometimes pretibial myxedema, often thyroid eye disease	
Thyroid ophthalmopathy	• Rarely exophthalmos (<2%), • More frequently hypothyroid signs, such as suborbital edema		• Exophthalmos (in 25%), • Other signs: lid lag, eyelid edema, chemosis, extraocular muscle weakness	
Thyroid gland	• Thyroid atrophy and (micro) nodules are more common, • Occasionally goiter		• Goiter	
Histological findings	• Mononuclear cells, mostly lymphocytes (especially T cells)		• Activated follicular cells • Increased vascularization	
	• Destroyed thyroid follicles		• Hyperplastic follicles with colloidal absorption	
Thyroid ultrasound	• Reduced echogenicity (progresses to atrophic hypovascular thyroid)		• Increased echogenicity (diffusely enlarged hypervascular thyroid gland)	

125. A middle aged lady presents with weight gain, constipation, hoarseness of voice and lethargy. On examination, she is found to be having anti TPO antibodies. What is the likely diagnosis in her?


a) Reidel's thyroiditis
b) Graves' disease
c) Hashimoto thyroiditis
d) De quervain thyroiditis


Ans. C

126. A lady has been diagnosed with breast cancer. Her biopsy report revealing the presence of tumor cells with signet ring appearance, mucin deposition and "Indian file pattern" is most likely suggestive of

a) Invasive lobular carcinoma
b) Invasive ductal carcinoma
c) Medullary cancer
d) Mucinous


Ans. A





www.medicalcoupons.in

127. Histological examination of a bone tumor is as follows . What is thelikely diagnosis ?



a) Osteosarcoma
b) Giant cell tumor
c) Ewings sarcoma
d) Chrondrosarcoma

Ans. B



Giant cell tumor

- The typical age group 20-40years.
- Morecommon in females
- Usually occurs at epiphysis
- Most common location is distal femur>proximaltibia
- Usually solitary
- Malignant transformation seen in less than 5% cases.

128. Which of the following anti-diabetic drug does not require dose reduction in renal failure?

a) Linagliptin
b) Sitagliptin
c) Vildagliptin
d) Saxagliptin

Ans. A

- Normally liptins are avoided in renal failure except: linagliptin is the only drug safe in renal failure
- Liptins are dipeptidyl peptidase-4 inhibitor⇨they inhibit breakdown of GLP⇨↑GLP⇨release insulin⇨↓blood sugar

www.medicalcoupons.in

www.medicalcoupons.in

129. Which of the following drugs used in emergency management of asthma, which does not act by causing bronchodilation?

a) Salbutamol
b) Ipratropium
c) Deriphylline
d) Hydrocortisone

Ans. D

- There are 3 major types of bronchodilators:
  - I. Beta2 agonists [eg:salbutamol,salmetrol]
  - II. ACE inhibitors [eg:ipratropium,thiotropium]
  - III. PDE inhibitors [eg:theophylline,deriphylline,aminophylline]

130. Which of the following is the action of dopamine at low dose ?

a) It increases renal blood flow
b) It causes vasoconstriction
c) It has positive chronotropic action
d) It causes hypotension

Ans. A

- Low dose: <1-2µg/kg/min
  - ✓ At low dose dopamine stimulates only D1 receptors
  - ✓ D1 receptors are present on renal blood vessels
  - ✓ So when dopamine stimulate D1 receptors ⇨ renal vasodilation ⇨ increase in renal blood flow
- Intermideatedose : 2-10µg/kg/min
- High dose :>10µg/kg/min

131. A female patient presented with pain and redness in great toe.serum uric acid level is 9.6mg/dl. A part from prescribing analgesics for relieving pain , the physician prescribed a drug which can decrease the formation of uric acid. Which of the following enzyme is likely to be inhibited by this drug ?

a) Xanthine oxidase
b) Thymidylate synthase
c) Phosphpribosyltransferase
d) Dhfr

Ans. A

- Uric acid formed from purines ⇨ which are metabolized to generate xanthine ⇨ and convert xanthine into uricacid by xanthineoxidase

132. Vitamin that increases the absorption of iron from stomach is ?

a) Vitamin A
b) Vitamin B12
c) Vitamin C
d) Vitamin D

Ans. C

www.medicalcoupons.in

- 2 major substances help in increase in absorption of iron :
  - I. Hydrochloric acid
  - II. Vitc/ascorbic acid
- 2 substances which decrease absorption of iron:
  - I. Phytates
  - II. Oxalates
  - III. Tannates

133. longest acting phosphodiesterase inhibitor among the following drugs is ?

a) Sildenafil
b) Vardenafil
c) Tadalafil
d) Phentolamine

Ans. C

PDE inhibitor
USES:

- Erectile dysfunction
- Pulmonary HTN

134. Which of the following statements about these drugs is true?




Fig.2: log dose response curve for potency of drug A & B

a) Drug D is most potent
b) Drug A and B have equal efficacy but B is more potent than drug B
c) Drug C is least potent
d) Drug B has more potency but less efficacy than drug a

Ans. B

- Potency: It defines left and right side of the curve
  - more the drug is on left side more potent the drug
  - more the drug is on right side less the potent the drug
- Efficacy: It defines by height of the curve
  - Higher the curve more efficacy
  - Lower the curve less efficacy

135. Which of the following antimicrobials is contraindicated in a patient with seizure disorder?

a) Ampicillin
b) Ofloxacin
c) Moxifloxacin
d) Cefixime

Ans. B

www.medicalcoupons.in

- 2 major groups of anti-microbials contraindicated in seizure disorders:
  I. Flouroquinolol [floxacindrugs] - ofloxacin > moxifloxacin
  II. Carbapenems [imipenem drugs]

136. Which of the following drug is preffered for closure of ductus arteriosus in a preterm baby with PDA ?

a) Indomethacin
b) Ibuprofen
c) Mefenamic acid
d) Paracetamol

Ans. B

- NSAIDS are used to close ductus arteriosus
- NSAIDS are: ibuprofen[best],indomethacin,aspirin

137. A patient on long term digitalis therapy develops arrythmia. What will be the next step in the management

a) Adenosine
b) Amiodarone
c) Atropine
d) Lignocaine

Ans. D

- Lidocaine: drug of choice for ventricular arrhythmias
- Atropine: drug of choice for atrial arrhythmias
- Amidarone: it causes toxicity
- Digoxin toxicity can emerge during long-term therapy as well as after an overdose. It can occur even when the serum digoxin concentration is within the therapeutic range.Toxicity causes anorexia, nausea, vomiting and neurological symptoms. It can also trigger fatal arrhythmias.
  "Digitalis" arrhythmias were treated with beta-adrenergic blockers: Inderal, Viskene, Eraldin, Trasicor and Aptin. These drugs proved effective in most cases with atrial arrhythmias and in some--with ventricular arrhythmias. Lidocain was more effective in cases of ventricular arrhythmias. Effective drugs of a broad spectrum are also Aimalin, Pulsenorma and Ritmodan.

138. A patient on treatment of myasthenia gravis presents with increasing muscle weakness.which of the following drug is used to differentiate myasthenia gravis from cholinergic crisis ?

a) Edrophonium
b) Pilocarpine
c) Atropine
d) Pyridostigmine

Ans. A

Explanation:

- Edrophonium is ACHesterase inhibitor
- It act only for 10min
- By giving edrophonium if the condition improves then it diagnose as Myasthenia gravis
- By edrophonim if the condition worsens then it diagnose as cholinergic crisis

www.medicalcoupons.in

139. Which of the following drugs can be used for the treatment of undescended testes?
a) Testosterone
b) GnRH
c) Cold Water Compress
d) Anti -MIH Therapy

Ans. B
- Pulsatile GnRH analogues [leuprolide] ⇨↑testosterone ⇨ help in movement of testis to scrotum

140. An elderly male developed the symptoms of dementia and was diagnosed as Alzhiemer's disease. Which neurotransmitter level is reduced in this disease ?
a) Dopamine
b) Acetylcholine
c) Adrenaline
d) Serotonin

Ans. B
- Alzhiemer's dementia occurs in basal nucleus of MEYNERT in brain
- MEYNERT function: is to maintain memory, this area contains plenty of Ach
- So if there is any problem in MEYNERT nucleus then Ach level will decrease

141. Which of the following combination of seizure and its first line drug is not correct ?
a) Generalised seizure – Valproate
b) Myoclonic- Topiramate
c) Focal- Levetiracetam
d) Absence- Ethosuximide

Ans. B
- Myoclonic -valproate

142. Orthotoulidine test is used to measure?
a) Free chlorine & combined chlorine
b) Free chlorine / Residual chlorine
c) Chlorine demand
d) Combined chlorine

Ans. D
Measure:- Total chlorine - free chlorine A/M → combine chlorine
O T test: Measures free chlorine (within 10 seconds) and free + combined chlorine (15-20 mint) in water.

143. Low stationary phase of demographic cycle?
a) Phase 1
b) Phase 2
c) Phase 3
d) Phase 4

Ans. B

Stage 1: (High stationary):- High BR
High DR

Stage 2: (Early expanding): Birth Rate Unchange
Death Rate decrease

Stage 3: (late Expnading): BR: decreases

www.medicalcoupons.in

DR: decreases

Stage 4: (late stationary): BR: low
DR: low

Stage 5: (Declining): BR < DR
Population decreases

Minimum demographic gap: stage 1 & stage 4
Maximum demographic gap: late part of stage 2
Negative D. graph in stage 5
India is in: stage 3 (late expanding stage)

144. A health worker goes house to house for every 15 days taking fever history and preparing malarial slide.
Which form of surveillance is this?
a) Active
b) Passive
c) Sentinel
d) Any of the above

Ans. A
Malarial surveillance was done by MPW (Male)→subcenter level
- Passive: Hospital reporting → Most convenient way of sample
- Sentinel: Done for air & water quality

145. Two vaccines MB and Pentavalent used for immunization round given for the worker which was open. Which of the following is vaccine is discarded?
a) MR discard, USE pentavalent
b) Pentavalent
c) Both MR & Pentavalent
d) MR can be used, Pentavalent must discarded

Ans. A
VACCINES WHICH WILL NOT FOLLOW OPEN VIAL POLICY
- BCG
- MEASLES
- MR
- JE
- YELLOW FEVER

146. Recent transmission of Malaria in community is measured by?
a) Annual Parasite incidence
b) Infant Parasite rate
c) Speed rate
d) Slide positivity rate

Ans. B
API: Most sensitive index for recent transmission of Malaria

Annual parasite index:

$$\frac{\text{comformed cases during 1 year} \times 1000}{\text{Population under surveliance}}$$

Measures:- Burden of Malaria

www.medicalcoupons.in

Slide positivity rate: % of slides found positive for malaria parasite irrespective of the type of speci

147. According to Bio medical wastage compare. Which Bag of Bio medical wastage can be incinerated?

a) Red
b) Blue
c) Yellow
d) White

Ans. C

Yellow: Incinevates
Red
Blue } Disinfected and later discarded
White

148. Given mosquito causes which disease ?



a) JE
b) Dengue
c) Zika virus
d) Malaria

Ans. A

GIVEN IMAGE :.CULEX
Causes:Failarisis
JE

.ANOPHELES
↓



Causes:
Malaria

→.ADES

www.medicalcoupons.in



Causes :
Dengue ,Riftvally fever, Yellow fever, Chicken gunia

149. The average number children that would be born to a woman over her reproductive life span refers to?

a) Total fertility rate
b) Gross reproduction rate
c) Net reproduction rate
d) General fertility rate

Ans. A

TFR:- Give magnitude of complete family size
GRR: Measures the no. of daughters a woman would have in her life time
NRR: No. of daughters a newborn girl will bear during her life time assuming fixed age specific fertility and mortality rate
NRR: 1
To active NRR=1, CPR> 60%
GFR:- No. of CB for 1000 people who are in reproductive age group

150. A pregnant lady was bit by a rabid dog, what should be the next step?

a) Treat the wound locally
b) Give vaccine immediately
c) Local treatment of wound + immunoglobulin + vaccine
d) Local treatment of wound + vaccination

Ans. D

151. Under Ayushman Bharat what is the insurance package provided?

a) Rs. 1 lk/family/year
b) Rs. 5 lk/family/year
c) Rs. 3 lk/family/year
d) Rs. 2lk/family/year

Ans. B

www.medicalcoupons.in

Image:- PM-JAY
Full form :Pradhan mantra janarogyayojana

152. School health is a responsibility by which level of healthin RURAL areas?

a) Sub center
b) PHC
c) CHC
d) Sub district hospital

Ans. B

Population covered by **PHC**
Plain :30000
Hilly: 20000
Population covered by **Sub center**
Plain :5000
Hilly: 3000
Population covered by **CHC**
Plain :120000
Hilly: 80000
CHC is the first referral level

153. All are principals of Primary Health care EXCEPT?

a) Appropriate technologies
b) Community participation
c) Inter sectoral coordination
d) Universal Health coverage

Ans. D

4th principle :Equity of distribution

154. Survey of 4 cities. A, B, C, D reported IFR as 2.1, 2.4, 2.6, 2.8. which city had achieved desire goal of TFR?

a) 2.1 (A)
b) 2.4 (B)
c) 2.6 (C)
d) 2.7 (D)

Ans. A

TOTAL FERTILITY RATE : Single most important factor in population growth

155. Which vaccine follows open vial policy?

a) BCG
b) Pentavalent
c) JE
d) MMR

Ans. B

Open vial vaccines can be used for 4 weeks
Vaccines follows open vial policy are:
PENTAVALENT .OPV
. HEP B . IPV
. DPT . Td

www.medicalcoupons.in

156. Which vaccine has seroconversion time less than incubation period for that disease?

a) Flu
b) Measle
c) Rubella
d) Mumps

Ans. B

- Flu:- I.P:- 7 days after vaccine AB develops > 7 days
- Measles: I.P:-10-14 days:- after exposure If we give vaccines Ab will develops within 7 days
- Rubella:- I.P:- 14-21 days: AB take time: 2-4 weeks

Measles vaccine:

a) Pre exposure:

- Live attenustaved
- Strain: Edmostron ZAEREB
- Powered: We use: distilled water
- 0.5 ml s/c to Rt shoulder
- 2 dose 1) = 9-12 months
  2) = 1 1/2 year (16-24 months)

b) Pos Exposures

1) Measles vaccine
   Within 3 days of exposure
2) Ig (Immuno compromised child)
   0.25 mg/kg
   IM

157. A person reported to ophthalmic OPD with vision of <3/6 in RT eye and finger movement at 1 meter distance in LT eye. His state of vision is?

a) Severe visual impairment
b) Economical Blindness
c) Social Blindness
d) Manifest Blindness

Ans. C

Normal: 6/6 – 6/18

Low vision: < 6/18 – 6/60
Economic Blindness:-< 6/60 – 3/60

Social Blindness: <3/60 – 1/60

158. A man traveled to Assam and after returning back he developed fever, giddiness, and on investigation confirmed to be a case of PI falciparum, Tx of choice?

a) Primaquine
b) Chloroquine
c) A/V artemether
d) Artesunale, Salfadoxine, Pyridethamine

Ans. C

P. vivax:-Chloroquin in all trimesless
No primaquive

P. falciparum: Quinine → 1st trment ACT(artemisinin combination therapy)


www.medicalcoupons.in

NE state: ACT-L
Otur: ACT – SP
No primaquin
} 2nd& 3rdtrimes

159. A worker in the wheat section of a food facility eats peanutsregularly and develops hepatic carcinoma after
many years.Which of the following is implicated in the pathogenesis of his condition?
a) BOAA
b) Aflotoxin
c) Pyrolizidine
d) Sanguanarine

Ans. B

BOAA:-Latherysm
Aflatoxin:- Aspergillus flavours fungus : causes :hepatocellular carcinoma
Pyrocizidine:- Endemic asitis
SANGUANARINE:- Epidemic drop (mustard oil+ argenomone oil)



160. In a hospital the management wants to know that how much time that each doctor is spending with patient
on an average. They want to know increase or decrease in the number of doctors as per requirement, to
Improve the quality of care. Which management technique should be used?
a) FGD
b) Systemic Analysis
c) Network Analysis
d) Work Sampling

Ans. D

161. A case control study was done in a group of people who are ordering food and who are preparing food to see
in which group chances of Obesity is more what sample should the investigator should take?
a) Person with or without ordering food
b) Person with or without exerces
c) Person with and without ordering food
d) Person without ordering food

www.medicalcoupons.in

Ans. C

162. Demographic Bonus is?
a) Population density
b) Reduction in population in last few years due to contraception
c) Fertility rate
d) Literacy

Ans. B

163. Reproductive health promoting agencies in world is?
a) UNDP
b) UNICEF
c) UNFPA
d) UNESCO

Ans. C





UNFPA: United nation fund
Year of formation: 1967
HEADQUATORS: NEW YORK, USA

164. Which agency give support to active sustandable development goals?
a) WHO
b) UN
c) UNESCO
d) UNDP

Ans. D



UNDP: UNITED NATIONS DEVELOPMENT PROGRAM
Year of Formation :1965


www.medicalcoupons.in

Head quators; NEW YORK.USE

165. Which agency is funding for TB program in India?
a) SIDA
b) WHO
c) WORLD BANK
d) GLOBAL FUND

Ans. A

166. Study the graph given below of different occupational group's. Analyse and mark who has the highest chance of developing fever?



a) Industrial Worker
b) White collar Worker
c) Farmer
d) Housewives

Ans.

167. A person with depression prescribed a serotonin and norepinephrine reuptake inhibitor. Identify the drug?
a) Fluoxeitine
b) Venlafaxine
c) Sertraline
d) All of the above

Ans. B
Serotonin nor-epinephrine reuptake inhibitors:
- Venlafaxine
- Desvenlafaxine
- Duloxetine

Selective serotonin reuptake inhibitor:
- Escitalopram
- Paroxetine
- Fluoxetine

168. Earliest symptoms of alcohol withdrawal is:
a) Tremors
b) Hallucinations
c) Seizures
d) Delirium

Ans. A
Alcohol withdrawal
Symptoms
- After 6-8 hours: coarse tremors (m/c), nausea, vomiting, anxiety.

www.medicalcoupons.in

- After 12- 24 hours: alcoholic hallucinations (auditory).
- After 24-48 hours: alcohol withdrawal seizures (GTCS)—comes in clusters.
- After 48-72 hours: delirium tremens—(lillputian hallucination-patient sees everything small)

169. A chronic alcoholic patient develops delirium tremens after stopping alcohol. Patient should be treated with?

a) I/V methycobalamine
b) I/V thiamine
c) I/V thiamine + lorazepam
d) I/V methylcobalamine + lorazepam

Ans. C

170. Not a symptom of panic attack:

a) Suicide attempt
b) Sudden anxiety
c) Episodic anxiety
d) Feeling of impending doom

Ans. A

Panic attack →sudden, discrete period of intense fear or discomfort that are accompanied by significant physical and cognitive symptoms.
C/F: palpitations, chest discomfort, dizziness,discomfort, sweating, trembling, shaking.
Cognitive symptoms : disorganized thinking, irrational fears, depersonalization, decreased ability to communicate.
Fear of death: mimics heart attack
Peak at 10 minutes, lasts upto 30 minutes.

171. A 32 years old women presented with symptoms of mild depression first time. What should be the initial treatment?

a) Cognitive behaviourtherapy(CBT)
b) Antidepressant
c) CBT+ antidepressant
d) Electro convulsive therapy

Ans. A

Treatment of depression
Mild depression
For 1st time:

- CBT- Multiple sessions maybe needed.
- Exercise-early morning
- Active monitoring

If not getting better start anti-depressants
Antidepressents
SSRI'S, SNRI'S, TCA'S, MAOI'S, Atypical antidepressant (choosen on the basis of side effects).


172. A patient suffering from RTA 6 hrs back he was brought to the emergency with altered sensorium. his Glasgow coma score was 12. CT scan of head done which showed the following. he is most likely suffering from –

a) Sub dural hematoma
b) Epidural hematoma
c) Subarachnoid hemorrhage
d) Intracerebral hematoma

www.medicalcoupons.in

Ans. B

Explanation: EPIDURAL HEMATOMA




- ✓ ***Location:*** lies between skull and outer endosteal layer of duramater
- ✓ ***Vessels involves***:middle meningeal artery
  Anterior ethmoidal artery
  Transverse or sigmoid sinus
  Superior sagittal sinus
- ✓ ***Presentation:***
  Symptoms develop soon after injury Lucid interval in 20% of cases
- ✓ ***CT appearance***: Biconvex lens

173. Barium swallow image demonstrates

a) Achalasia cardia
b) Zenker's diverticulum
c) Corkscrew esophagus
d) Cancer esophagus

Ans. A


ACHALASIA CARDIA



- ✓ Rat-tail appearance/bird-beak tapering
- ✓ Dilated sigmoid esophagus

www.medicalcoupons.in

174. What is the probable cause of the condition shown in the given X-ray?




a) Pleural effusion
b) Perforation peritonitis
c) Sigmoid volvulus
d) Intestinal obstruction

Ans. B

PERFORATION PERITONITIS

✓ Air under diaphragm⇨pneumoperitoneum

175. Following image is a suggestive of



a) Congenital diaphragmatic hernia
b) Pneumothorax
c) Cardiac tamponade
d) Tracheo esophageal fistula

Ans. A

Congenital diaphragmatic hernia

✓ Depending on the location and size of the defect retroperitoneal or intra-abdominal organs and tissues can prolapse intothoraciccavity due to negative intra-thoracic pressure

www.medicalcoupons.in

176. Barium enema image demonstrates



a) Meckel's diverticulum
b) Diverticulosis of colon
c) Sigmoidvulvulus
d) Colon cancer

Ans. B

Explanation: DIVERTICULOSIS


- ✓ SAW-TOOTH APPEARANCE
- ✓ ***Description***: serrated/concertina-like appearance
- ✓ ***Mc site***: sigmoid colon
- ✓ ***Seen in***: Diverticulitis

177. A chronic smoker presented to the emergency . His X-ray chest was done which showed the following.
Possible diagnosis in this patient

a) Pneumothorax
b) Tension pneumothorax
c) Hemothorax
d) Hydropneumothorax

Ans. B


TENSION PNEUMOTHORAX



- ✓ Bronchovascular margins are absent
- ✓ Lung collapse towards hilum
- ✓ Heart shifted towards right side

www.medicalcoupons.in

178. Alcoholic male presented with pain in epigastrium radiating to the back.what is the likely diagnosis in the given CT image.




a) Amoebic liver abscess
b) Pancreatic pseudocyst
c) Acute cholecystitis
d) Hydatid cyst

Ans. B

PANCREATIC PSEUDOCYST

- ✓ ***Duration***:>4weeks
- ✓ Fibrous wall is seen
- ✓ Homogenous collection is seen

179. T1 and T2 images in MR



a) Caudate
b) Thalamus
c) Lentiform nucleus
d) Insular cortex

www.medicalcoupons.in

Ans.

- ✓ T2 is the time taken for hydrogen atoms to diphase
- ✓ T1 is the time taken for return to relaxed state (and release energy)

	T1 Image	T2 Image
Time to echo (TE)	40 ms	120 ms
Time to repetition (TR)	400 ms	1000 ms
Useful for	Demonstrating anatomy	Demonstrating pathology
Bright for	Fat, methaemoglobin, contrast, calcium, blood	Fluid (oedema), fat

180. What is most responsive to radiotherapy?

a) Mucosa
b) Bone
c) Nerves
d) Muscle

Ans. A

181. A 65 year old female presented with imbalance while walking. On examination there is loss of vibration and proprioception. What will be next line of treatment

a) Iron
b) Vitamin B12
c) Niacin
d) Folic acid

Ans. B

Vitamin B_{12}deficiency symptoms include:

- strange sensations, numbness, or tingling in the hands, legs, or feet.
- difficulty walking (staggering, balance problems)
- anemia.
- a swollen, inflamed tongue
- difficulty thinking and reasoning (cognitive difficulties), or memory loss.
- weakness.
- fatigue

182. A patient presented with complaints of fatigue, dyspnea on exertion.Lab tests shows macrocytic RBCS. Serum methylcobalamin levels within normal limits.What will be next line of treatment

a) Iron
b) Folic acid
c) Vitamin B12
d) Pyridoxine

Ans. B

The symptoms of anemia that occur due to folate deficiency include:

- Persistent fatigue.
- Weakness.
- Lethargy.
- Pale skin.
- Shortness of breath.
- Irritability.

www.medicalcoupons.in

Laboratory Diagnosis

Test	B12 deficiency	Folate deficiency	Comment
Serum B12	↓	Nl	Less reliable for B12 deficiency
Serum folate	Nl	↓	If > 4 ng/mL, NOT folate deficiency
Methylmalonic Acid	↑	Nl	More reliable for B12 deficiency
Homocysteine	↑	↑	More reliable for B12 deficiency

183. Which of the following presentations has worst prognosis

a) Paralysis
b) Parasthesia
c) Pallor
d) Pulselessness

Ans. D

Pulseless electrical activity (PEA), a cardiac arrest rhythm scenario with an associated poor prognosis, is defined as cardiac electrical activity without a palpable pulse.

184. Central venous pressure is increased in which type of shock

a) Hypovolumic
b) Septic
c) Neurogenic
d) Cardiogenic

Ans. D

Central venous pressure (CVP) has a normal range of 5–7 mm Hg in an adult spontaneously breathing patient while supine. The **CVP** is elevated in obstructive or cardiogenic **shock**, while it is decreased in septic, neurogenicand hypovolemic **shock**.

Type	CO	HR	CVP	PCWP	SVR	02 Sat
Cardiogenic	↓	↑	↔ ↑	↑	↑	↓
Hypovolemi	↓	↑	↔ ↓	↓	↑	↓
nti	↑	↑	↔ ↓	↓	↓	↑
ogenic	↓	↓	↔ ↓	↓	↓	↓
Anaphylactic	↓	↑	↔ ↓	↓	↓	↓

www.medicalcoupons.in

185. A patient diagnosed to have empyema. Which of the following is the best treatment

a) Antibiotics only
b) Thoracotomy + antibotics
c) Pleurodesis + antibiotics
d) Intercostal chest tube drainage + antibiotics


Ans. D

Empyema is a collection of pus in the space between the lung and the inner surface of the chest wall (pleural space).
Treatment is aimed at removing the pus and fluid from the pleura and **treating** the infection. Antibiotics are used to **treat** the underlying infection. The specific type of antibiotic depends on what type of bacteria is causing the infection. The method used to drain the pus depends on the stage of the **empyema**.

186. A female presented with symptoms of lethargy, weight gain, constipation, hair loss, dry skin and intolerance
tocold. What is the most likely diagnosis

a) Hypothyroidisim
b) Hyperthyroidism
c) Hypoparathyroidism
d) Hyperparathyroidism

Ans. A



187. An elderly male presented with dyskinesia and rigidity. There was some pathology in substantia nigra.Which
neurotransmitter is most likely involved

a) Acetylcholine
b) Serotonin
c) Glycine
d) Dopamine

www.medicalcoupons.in

Ans. D

The **substantia nigra** (SN) is an area of deeply pigmented cells in the midbrain that regulates movement and coordination. ... Neurons of the SNc produce Dopamine, which stimulates movement. In contrast, GABAergic neurons of the SNr can stimulate or inhibit movement depending on the input signal.

188. Elderly male with low backache and sclerotic lesions in vertebra is most likely having which condition

a) Ca prostate
b) Multiple myeloma
c) TB of vertebra
d) Waldenstrom macroglobulinemia

Ans. A

Osteoblastic **lesions** are asso- ciated with increased bone formation and appear **sclerotic**. Bone **lesions** resulting from **prostate cancer** are primarily osteoblastic but are also associated with increased bone resorption

189. A patient has serum K of 3.2 meq / L. What will be next line of treatment

a) IV KCl infusion
b) Oral KCI
c) IV Calcium gluconate
d) IV Insulin

Ans. B

Potassium supplements are generally prescribed for **low potassium levels**. If the situation is severe, **potassium** might be given as an intravenous (IV) solution.

Category & Clinical presentation		
State	**Serum K level**	**Clinical presentation**
Mild	3-3.5 mEq/L	Asymptomatic
Moderate	2.5-3 mEq/L	Cramping, weakness, malaise, and myalgias
Severe	< 2.5 mEq/L	ECG changes arrhythmias
ECG charges: ST-segment depression of flattering. T-wave inversion, and U-Wave elevation		

190. A young female with malar rash and fever.On Echocardiography there are vegetations on both sides of mitral valve. What is the likely diagnosis

a) A.RHD
b) B. Infective endocarditis
c) C. Non bacterial thrombotic endocarditis
d) Libman Sacks endocarditis

Ans. C

www.medicalcoupons.in




Figure 12-24 Comparison of the four major forms of vegetative endocarditis. The rheumatic fever phase of rheumatic heart disease (RHD) is marked by small, warty vegetations along the lines of closure of the valve leaflets. Infective endocarditis (IE) is characterized by large, irregular masses on the valve cusps that can extend onto the chordae (see Fig. 12-25A). Nonbacterial thrombotic endocarditis (NBTE) typically exhibits small, bland vegetations, usually attached at the line of closure. One or many may be present (see Figure 12-26). Libman-Sacks endocarditis (LSE) has small or medium-sized vegetations on either or both sides of the valve leaflets.

191. A patient had diarrhea associated with blood in stools. Physician diagnosed shigella as the etiological agent of diarrhea. Which clinical feature supported shigella over cholera

a) Abdominal pain
b) Fever
c) Vomiting
d) Stool frequency

Ans. B

Symptoms	Cholera = acute watery diarrhoea	Shigella = acute bloody diarrhoea
Stool	> 3 loose stools per day. Watery like rice water	> 3 loose stools per day, with blood or pus
Fever	No	Yes
Abdominal cramps	Yes	Yes
Vomiting	Yes a lot	No
Rectal pain	No	Yes

192. A young boy was found to have wing beating tremors.Which is the most likely diagnosis

a) Sydenham Chorea
b) Wilson disease
c) Huntington chorea
d) Neurodegenative disease

Ans. B

Wing-beating tremor is a coarse, irregular **tremor** that is most prominent when the limbs are held outstretched, reminiscent of a bird's **flapping** its **wings**; due to up and down excursion of arm at abducted shoulder. Seen mainly with Wilson disease.


193. What deformity is present as shown in the figure

a) Boutonniere deformity
b) Swan neck deformity
c) Mallet finger
d) Z deformity



www.medicalcoupons.in

Ans. B



194. Which of the following agent doesn't cause bronchodilatation still used in emergency management of bronchial asthma

a) Salbutamol
b) lpratropium
c) MgSO4
d) Hydrocortisone

Ans. D

Corticosteroids reduce the mucus secretion by inhibiting the release of secretagogue from macrophages. Corticosteroids inhibit the late phase reaction by inhibiting the inflammatory response and interfering with chemotaxis. This action may be due to the inhibition of LTB4 release.

195. Corticospinal pathway crosses to opposite side at what level

a) Midbrain
b) Pons
c) Upper medulla
d) Lower medulla

Ans. D

The **corticospinal tract**, along with the corticobulbar **tract**, form two pyramids on either side of the medulla of the brainstem—and give their name as pyramidal **tracts**. ... The lateral **corticospinal tract** neurons **cross** the midline at the level of the lower medulla of medulla oblongata, and controls the limbs and digits.

196. A young patient brought to the ED after road traffic accident. CT head shows evidence of raised intracranial
pressure. What will be the next line of management

a) Mannitol
b) Furosemide
c) Acetazolamide
d) Glycerol

Ans. A

www.medicalcoupons.in

TABLE 301-2 Stepwise Approach to Treatment of Elevated Intracranial Pressure (ICP)[a]

Insert ICP monitor—ventriculostomy versus parenchymal device

General goals: maintain ICP <20 mmHg and CPP ≥60 mmHg. For ICP >20–25 mmHg for >5 min:

1. Elevate head of the bed; midline head position
2. Drain CSF via ventriculostomy (if in place)
3. Osmotherapy—mannitol 25–100 g q4h as needed (maintain serum osmolality <320 mosmol) or hypertonic saline (30 mL, 23.4% NaCl bolus)
4. Glucocorticoids—dexamethasone 4 mg q6h for vasogenic edema from tumor, abscess (avoid glucocorticoids in head trauma, ischemic and hemorrhagic stroke)
5. Sedation (e.g., morphine, propofol, or midazolam); add neuromuscular paralysis if necessary (patient will require endotracheal intubation and mechanical ventilation at this point, if not before)
6. Hyperventilation—to $Paco_2$ 30–35 mmHg (short-term use or skip this step)
7. Pressor therapy—phenylephrine, dopamine, or norepinephrine to maintain adequate MAP to ensure CPP ≥60 mmHg (maintain euvolemia to minimize deleterious systemic effects of pressors). May adjust target CPP in individual patients based on autoregulation status.
8. Consider second-tier therapies for refractory elevated ICP
   a. Decompressive craniectomy
   b. High-dose barbiturate therapy ("pentobarb coma")
   c. Hypothermia to 33°C

197. A 30 year old presented with vomiting and pain epigastrium which is radiating to back.On examination the findings asshown in the figure are noted.What is the diagnosis



a) Acute pancreatitis
b) Acute cholecystitis
c) Acute hepatitis
d) Ectopic ruptured pregnancy

Ans. A

Cullen's sign is **superficial edema and bruising in the subcutaneous fatty tissue around the umbilicus** is named for gynecologist Thomas Stephen Cullen (1869–1953), who first described the sign in ruptured ectopic pregnancy in 1916. This sign takes 24–48 hours to appear and can predict acute pancreatitis, with mortality rising from 8–10% to 40%

Symptoms of acute pancreatitis

- Fever.
  Higher heart rate.
- Nausea and vomiting.
- Swollen and tender belly.

www.medicalcoupons.in

- **Pain** in the upper part of your belly that goes into your back. Eating may make it worse, especially foods high in fat.




198. A patient with deep X descent and rapid Y descent in JVP is most likely having

a) Constrictive pericarditis
b) Tamponade
c) Acute pericarditis
d) D.TR

Ans. A

Constrictive pericarditis

- JVP is elevated
- a wave is usually normal
- v wave is usually equal to a wave
- x descent –prominent
- y descent – rapid descent
- Kussmauls sign is usually positive



199. A middle aged male presented with fatigue and weakness more in evening.His symptoms improved after administeringedrophonium.What is the most likely diagnosis

a) ALS
b) LES
c) Myasthenia Gravis
d) Botulism

Ans. C

The Tensilon **test** uses the drug Tensilon (**edrophonium**) to help you to **diagnose myasthenia gravis**. Tensilon prevents the breakdown of the chemical acetylcholine, a neurotransmitter that nerve cells release to stimulate your muscles.

200. A patient while being operated for a thoracic surgery have a sudden cardiovascular collapse.There are decreasedbreath sounds on right side.What is the likely cau

a) Tension pneumothorax
b) ARDS
c) Cardiac tamponade
d) Atelechesis

Ans. A

Tension pneumothorax is a major life-threatening condition that must be recognized by clinical findings and immediately treated. Air from a leak in the thoracic wall or the lung becomes trapped in the pleural space, decreasing venous return to the heart and **collapsing** the opposite lung.

www.medicalcoupons.in


201. A male patient presented to ED with plasma glucose of 350 mg / dl and positive urine ketones. Serum K 3.8
mmol / L,serum na- 130 mmol / L. Patient is dehydrated. You start normal saline. What will be added in normal
saline

a) Glucose
b) NAHCO3
c) Insulin
d) KCI

Ans. C

Diabetic ketoacidosis is a serious complication of **diabetes** that occurs when your body produces high levels of blood acids called ketones. The condition develops when your body can't produce enough insulin

202. A patient presented with these eye examination findings. He has behavior abnormalities and poor hand
writing. Whichgene mutation is responsible for this presentation



a) ATP 7A gene on chromosome 13
b) ATP 7B gene chromosome 13
c) ATP 7C gene on chromosome 13
d) ATP 7B gene on chromosome 6

Ans. B

Wilson disease Symptoms:

- Fatigue, lack of appetite or abdominal pain.
- A yellowing of the skin and the whites of the eye (jaundice)
- Golden-brown eye discoloration (Kayser-Fleischer rings)
- Fluid build up in the legs or abdomen.
- Problems with speech, swallowing or physical coordination.
- Uncontrolled movements or muscle stiffness.

203. The definition of steroid resistant nephrotic syndrome is absence of remission even after taking steroids for

a) 4 weeks
b) 6 weeks
c) 8 weeks
d) 12 weeks

Ans. A

Steroid-resistant nephrotic syndrome (SRNS) is defined as NS **resistant** to **steroid** therapy, defined by the absence of complete remission after four weeks of daily **prednisone** therapy at a dose of 60 mg/m^2 per day

www.medicalcoupons.in

204. Which of the following is true about severe vitamin D deficiency

a) Low calcium, high PTH
b) Low calcium, Low PTH
c) Normal calcium, High PTH
d) High calcium, high PTH

Ans. A

When **vitamin D level** is **low**, the absorption of **calcium** in the intestines becomes less, which then causes the **level** of **calcium** in the blood to go down. As a consequence the **parathyroid** glands become more active and produce more **PTH** that causes **calcium** to come out of the bones, therefore weakening the bones.

205. A 5 year old boy has hyperflexible fingers and can touch back of hand with wrist extended. What is the most likelydiagnosis

a) Marfans syndrome
b) Ehler Danlos syndrome
c) Alport syndrome
d) Homocystinuria

Ans. B

Ehlers-Danlos syndrome is a group of inherited disorders that affect your connective tissues — primarily your skin, joints and blood vessel walls. Connective tissue is a complex mixture of proteins and other substances that provide strength and elasticity to the underlying structures in your body.

206. A 22 year old male presents with history of recurrent fall and difficulty in ambulation.On examination atrophied legmuscles are observed, Elder brother has same complaints.What is the diagnosis



a) Duchhene muscle dystrophy
b) Becker's muscle dystrophy
c) Charcot Marie Tooth disease
d) Myotonic dystrophy

Ans. C

A group of hereditary disorders that damage the nerves in the arms and legs.
Charcot-Marie-Tooth is a degenerative nerve disease that usually appears in adolescence or early adulthood.
Muscle weakness, decreased muscle size, decreased sensation, hammer toes and high arches are symptoms.
The main treatments are physiotherapy and occupational therapy. Medication may reduce pain.
Muscular: abnormality walking, flaccid muscles, instability, lack of coordination, muscle weakness, rhythmic muscle contractions, or loss of muscle
Foot: hammer toe, high foot arches, constantly walking on tip toe, or difficulty raising the foot
Sensory: pins and needles or reduced sensation of touch
Also common: hand contractures, slow reflexes, or tremor

207. A young female presents in her first trimester of pregnancy. She is taking warfarin for prosthetic heart valve. What should be next appropriate step

www.medicalcoupons.in

a) Continue warfarin
b) Add vitamin K
c) Stop warfarin and switch to heparin
d) MTP

Ans. C

Warfarin therapy should be avoided during **pregnancy**. If **warfarin**therapy is essential, it should be avoided at least during the first trimester (because of teratogenicity) and from about 2 to 4 weeks before delivery to reduce risk of hemorrhagic complications.

Upto 12 weeks - LMWH
12- 36 weeks - Warfarin
➢ 36 weeks - LMWH

208. What is the arthropathy in diabetes with given x ray of ankle known as



a) Osteomyelitis
b) Charcoat arthropathy
c) Gout
d) Osteoarthritis

Ans. B

Charcot foot, also called **Charcot arthropathy**, is a disease that attacks the bones, joints, and soft tissue in your feet. When it starts, you may not realize something's wrong. But eventually, it can cause painful sores or change the shape of your foot.

209. What is the acid base abnormality in patient presented with multiple episodes of vomiting
a) Metabolic acidosis
b) Metabolic alkalosis
c) Respiratory alkalosis
d) Respiratory acidosis


Ans. B

Vomiting or nasogastric (NG) suction generates **metabolic** alkalosis by the loss of gastric secretions, which are rich in hydrochloric acid (HCl). Whenever a hydrogen ion is excreted, a bicarbonate ion is gained in the extracellular space.

210. An arterial blood gas is performed and reveals-pH-7.2, paCO2-68 mmHg, and serum HCO3- 26 mmol / L,
Diagnosis?
a) Respiratory acidosis partially compensated
b) Respiratory acidosis fully compensated
c) Metabolic acidosis with respiratory acidosis
d) Respiratory acidosis uncompensated

Ans. D

www.medicalcoupons.in



211. What is the GCS in a patient with eye openi to p in appropriate words and limb withdrawl is

a) E2, M4,V3
b) E2, M3.V4
c) E2, V2,M4
d) E1, V3, M3

Ans. A

Glasgow Coma Scale

BEHAVIOR	RESPONSE	SCORE
Eye opening response	Spontaneously	4
	To speech	3
	To pain	2
	No response	1
Best verbal response	Oriented to tim pl e, a person	5
	Confused	4
	Inappropr e wo	3
	Incomp ehe ble sounds	2
	No res onse	1
Best motor response	Ob eys co mands	6
	ves to lo ized pain	5
	Fle n withdrawal from pain	4
	Abno l flexion (decorticate)	3
	Abnormal extension (decerebrate)	2
	No sponse	1
Tot e:	*Best response*	15
	Comatose client	8 or less
	Totally unresponsive	3

212. A female presents with chronic diarrhea, fatigue and dyspnea on exertion. On examination the findings are shown inthe figure. Lab test shows low Hb. What will be the appropriate treatment

www.medicalcoupons.in




a) Doxycycline
b) Folic acid
c) High protein diet
d) Iron

Ans. D

Koilonychia affects the shape of nails. The nails start to curve like a spoon. Possible causes include nutritional deficiencies and autoimmune conditions.

Iron deficiency is the most frequent cause of koilonychia. Iron deficiency anemia is the world's most common nutritional deficiency disease. It most commonly affects children and women of childbearing age. Koilonycia is a nail finding seen in iron deficiency anemia, syphilis, ischemic heart disease.



213. All the following conditions are associated with the sign shown in figure except



a) Pregnancy
b) Hypoestrogen state
c) Rheumatoid arthritis
d) Cirrhosis

Ans. B

Palmar erythema is a skin condition that makes the palms of your hands turn red, palms have turned red because of dilated capillaries, which are the smallest blood vessels in the body.Mostly associated with **Pregnancy,Liver disease,Autoimmune diseases,Thyroid disease and it is a hyper estrogenic state.**

www.medicalcoupons.in

214. Where is this pacemaker lead located ?




a) LA
b) RA
c) LV
d) RV

Ans. D

There are three basic kinds of pacemakers:
1-Single chamber: One lead attaches to the upper or lower heart chamber.
2-Dual-chamber: Uses two leads, one for the upper and one for the lower chamber.
3-Biventricular pacemakers(used in cardiac resynchronization therapy).

215. A patient presented with recurrent syncope episodes. ECG strip shows the following findings. What is the most appropriate treatment option for him



a) Pacemaker
b) ICD
c) Anticoagulant
d) Antiarrhythmic drugs

Ans. A

Sinus bradycardia is a type of slow heartbeat. A special group of cells begin the signal to start your heartbeat. These cells are in the sinoatrial (SA) node. Normally, the SA node fires the signal at about 60 to 100 times per minute at rest. In **sinus bradycardia**, the node fires less than 60 times per minute.

Treatment of post infectious **bradycardia** usually requires permanent pacing. In patients with hypothermia who have confirmed **sinus bradycardia** with a pulse, atropine and pacing are usually not recommended because of myocardial irritability.

www.medicalcoupons.in

216. A 30-year-old patient was having breathingdifficulty on doing day to day activity. His physiciannoticed
dullness onpercussion in right infra-axillaryarea with reduced air entry. CXR showed pleuraleffusion and
pleural tapping was done. What findings will suggest an exudative pleural effusion?

a) Pleural fluid protein = 3.5 gm% and LDH= 100 IU
b) Pleural fluid protein=4.5 gm% and glucose= 30mg%
c) Pleural fluid LDH =90 IU and glucose=60mg %
d) Pleural fluid protein=3.5gm% and glucose=90 mg%


Ans. A

Light's Criteria		
	Transudate	Exudate
Pleural:Serum Protein	< 0.5	≥ 0.5
Pleural:Serum LDH	< 0.6	≥ 0.6
Pleural fluid LDH	< 2/3 upper limit of normal	> 2/3 upper limit of norma
Main Causes	• CHF • Cirrhosis • Nephrotic syndrome • Pulmonary embolism	• Malignancy • Bacterial/Viral pneumoni • Tuberculosis • Pulmonary embo sm • Pancreatitis • Esopha eal rup re • Collag rascu disease • Chylotho Hemo

217. In ED a patient becomes unresponsive and ECG showed arrhythmia. DC shock is immediately given. But he is
still unresponsive and arrhythmia is persisting. What is the next step

a) Repeat DC shock AV
b) CPR
c) Start bag and mask ventilation
d) Check carotid pulse

Ans. B



www.medicalcoupons.in

218. In ICU a patient is on mechanical ventilator. His ABG shows features of hypoventilation leading to respiratory acidosis. What will you do next to improve the ventilation of the patient

a) Increase FiO2
b) Increase respiratory rate
c) Increase PEEP
d) Increase expiratory time

Ans. B

Respiratory failure is divided into **type I** and **type II**. **Type I respiratory failure** involves low oxygen, and normal or low carbon dioxide levels. **Type II respiratory failure** involves low oxygen, with high carbon dioxide.

Type 1 respiratory failure may require only supplementary oxygen, but type 2 failure may require additional support such as continuous positive airway pressure (CPAP) or biphasic positive airway pressure (BiPAP) to increase exchange of both gases and, where possible, reverse any causes for low tidal volumes or low respiratory rates.


219. Which blood vessel is involved in Type II Takayasu arteritis

a) Arch of aorta
b) Arch of aorta and ascending aorta
c) Arch of aorta and descending aorta
d) Abdomen aorta

Ans. B

Takayasu arteritis can be divided into the following six types based on angiographic involvement:

- Type I - Branches of the aortic arch
- Type IIa - Ascending aorta, aortic arch, and its branches
- Type IIb - Type IIa region plus thoracic descending aorta
- Type III - Thoracic descending aorta, abdominal aorta, renal arteries, or a combination
- Type IV - Abdominal aorta, renal arteries, or both
- Type V - Entire aorta and its branches



Type	Vessel Involvement
Type I	Branches from the aortic arch
Type IIa	Ascending aorta, Aortic arch and its branches
Type IIb	Ascending aorta, aortic arch and its branches, thoracic descending aorta
Type III	Thoracic descending aorta, abdominal aorta, and / or renal arteries
Type IV	Abdominal aorta and / or renal arteries
Type V	Combined features of types IIb and IV

www.medicalcoupons.in

220. Which of the following will show pulsus paradoxus

a) Obstructive sleep apnea
b) Pleural effusion
c) Pulmonary hypertension
d) Acute exacerbation of asthma

Ans. D

Pulsus paradoxus (a decrease in the systolic blood pressure during inspiration) results from a decrease in cardiac stroke volume with inspiration due to greatly increased left-ventricular afterload. **Pulsus paradoxus** can be observed in cardiac tamponade and in conditions where intrathoracic pressure swings are exaggerated or the right ventricle is distended, such as severe acute **asthma** or exacerbations of chronic obstructive pulmonary disease.

221. 20-year-old Boy presented with pain in scrotum with red discoloration of scrotal skin. (On lifting the testis pain was decreased) Diagnosis?

a) Torsion testis
b) Inguinal hernia
c) Hydrocele
d) Epididmyo-orchitis

Ans. D

Prehn's sign is an evaluation used to determine the cause of testicular pain. It is performed by lifting the scrotum and assessing the consequent changes in pain. A positive **Prehn's sign** indicates relief of pain upon elevation of the scrotum and is associated with epididymitis but exacerbates the pain of **torsion.**

222. A Patient has been admitted to the ICU since he is having fever for past (4 days with new onset of breathlessness which was unresponsive to incremental flow rate of oxygen and saturation is progressively decreasing. What can be the diagnosis?

a) Atelectasis
b) Idiopathic Pulmonary fibrosis
c) ARDS
d) Mediastinitis chest pain

Ans. C

Refractory hypoxemia, and this term usually considered when there is inadequate arterial oxygenation despite optimal levels of inspired oxygen.
A hallmark of COVID-19 is the rapid development of refractory hypoxemia with a poor response to oxygen supplementation, suggesting intrapulmonary shunting. **Refractory hypoxemia** in **Covid** 19 patients occurs due to moderate to severe ARDS.

223. A 5-year old boy starts choking at dinner table and is not able to cough or speak. What is the immediate management to be done in this child?

a) Abdominal Thrust
b) Drink water / Carbonated drinks fast
c) Give back blows
d) Identify the food consumed

www.medicalcoupons.in

Ans. C



224. 46 xx, absent uterus, vagina, normal secondary sexual characteristics. Ovary present diagnosis, fsh and Lh
within normal limits?

a) MRKH
b) AIS
c) TURNER syndrome
d) Kliniefilter Syndrome

Ans. A

Bilateral mullerian duct agenesis

- ✓ Mullerian duct is absent ⇨ so fallopian tube (distal FT is present) , uterus, cervix, vagina 2/3
- ✓ Gonad ⇨ ovary
- ✓ Lower 1/3 rd of vagina is present (bcz it is from urogenital sinus)
- ✓ ***Phenotype*** : female
- ✓ ***Karyotype*** : 46 xx
- ✓ ***Clinical features*** :⇨primary amennorhea
  - ⇨no menses(no uterus)
  - ⇨no cyclic abdominal pain
  - ⇨normal secondary sexual characters (breast development)
- ✓ ***Examination*** :⇨per vaginal : blind pouch vagina
  - USG : gonads present , no uterus
- ✓ ***Gold standard investigation*** :laproscopy
- ✓ ***Treatment*** : vaginoplasty (done just before or after marriage) ⇨Williams , Mc indoe's

225. A 25 year old female attends gynae OPD with c/o secondary amenorrhea. She has history of previous d and c
and her FSH levels are 6 IU/L . The probable cause of amenorrhea is

www.medicalcoupons.in

a) Sheehan syndrome
b) Ashermann syndrome
c) Premature ovarian failure
d) Pregnancy


Ans. B

Ashermansyndrome :

- ✓ It is a rare acquired condition of the uterus
- ✓ Scar tissue or adhesions form due to some form of trauma
- ✓ ***Causes*** :➪c- section
  ➪fibroids/ polyps

- ✓ ***Symptoms*** :➪no periods , pain but no bleeding bcz the blood is unable to leave the uterus bcz the exit is blocked by scar tissue
- ✓ Asherman syndrome increases risk during pregnancy :➪placenta previa
  ➪placenta increta
  ➪excesssive bleeding
- ✓ ***Diagnosis*** : Hysteroscopy
- ✓ ***Treatment*** : operative hysteroscopy : small instruments are attached to the end of the hysteroscope and used to remove the adhesions

226. 16 yr old girl with primary amenorrhea. Ssc well developed. On examination there is bluish bulging
hymen. What is the best management ?



a) Cruciate incision and drainge of blood
b) long term treatment with estrogen
c) B / l gonadectomy
d) OCP

Ans. A

Imperforate hymen :

- ✓ Means when the hymen covers whole opening of vagina
  Symptoms :➪cryptomenorrhea
  ➪lack of menstrual cycle
  ➪secondary sexual characters are present
  ➪cyclic abdominal pain

www.medicalcoupons.in

- ✓ ***Diagnosis*** : on P/V : bluish hymen is seen
- ✓ ***Treatment*** : cruciate incision : x-shaped cut made in the hymen

227. A 55 year old women was found to have a carcinoma cervix figo stage 3b. What is the management?

a) Concurrent chemoradiotherapy
b) Radiotherapy + hpv vaccine
c) Schauta procedure
d) Intracavitary brachytherapy followed by ebrt

Ans. A

Stage 0	Carcinoma in situ, CIN
Stage I	Invasive carcinoma confined to the cervix
Stage I	Diagnosed only by microscopy
IA1	Micro-invasive carcinoma with stromal invasion <3 mm depth, <7 mm width
IA2	Micro-invasive carcinoma <5 mm depth, <7 mm width
Stage IB	Clinically visible or microscopic lesion >IA2
IB1	Clinical lesion <4 cm
IB2	Clinical lesion >4 cm
Stage II	Extension beyond cervix but not to sidewall
IIA	Involvement of upper two-thirds of vagina
IIB	Parametrial involvement
Stage III	Extension to pelvic wall and/or lower third of vagina; hydronephrosis
IIIA	Involvement of lower third of vagina
IIIB	Pelvic sidewall involvement; hydronephrosis
Stage IV	Extension beyond true pelvis or involving bladder or rectum
IVA	Involvement of bladder or rectal mucosa
IVB	Spread outside true pelvis or metastasis to distant organs

Management of cervical cancer :

- ✓ Surgery : is done for stages up till 2a2 except if size of tumour is >4cm (eg : 1b3,2a2)
- ✓ Radiotherapy: done in stage 2 onwards + stages 1b3 , 2a2.

Cisplatin(radiosensitizer): is given before radiotherapy →chemoradiation

228. A couple presented with infertility semen analysis is normal, female has menorrhagia. On hsg there is distal dilatation with fimbria What is the likely cause?

a) Tubal polyp
b) Tubal endometriosis
c) Tubal spasm
d) Tubal salpingitis

Ans. D

Gonorrhea : distal dilatation of fimbria is seen

- ✓ ***Etiology*** :neisseriagonorrheae (gram –ve diplococcic)
- ✓ ***Site*** : endo cervix >bartholinglan> urethra
- ✓ ***clinical features*** :➪ asymptomatic
  - ➪mucopurulent endocervical discharge
  - ➪urethral syndrome →dysuria
  - ➪pelvic pain
  - ➪post coital bleeding
- ✓ ***Investigation***: ➪culture : MC COY / HELA CELLS
  - ➪NAAT (nucleic acid amplification test) - best test
  - ➪urine test (1st in the mrng)

www.medicalcoupons.in

- ✓ ***complications*** : 1.FITZ-HUGH CURTIS SYNDROME
  2.REITER'S SYNDROME
  3.ACUTE SALPINGITIS
  4.ECTOPIC PREGNANCY
  5. INFERTILITY
- ✓ ***Treatment*** : doxycycline or azithromycin
- ✓ In pregnancy cyclines are contraindicated so RX of choice in pregnancy zithromycin

229. Women with history of previous cs presented with prolonged labor. Patient is hypotensive. Fhr is not audible, fetal parts are easily palpable on pa examination, what is the diagnosis?

a) Abruption
b) Uterine rupture
c) Fetal intrauterine death
d) Polyhydramnios

Ans. B

- ✓ Most common cause is previous ***caesarean section***
- ✓ Mcc innon scarred uterus is ***obstructed labor***
- ✓ Sign of impending rupture is ***fetal tachycardia***
- ✓ Sigh of rupture is ***fetal bradycardia***

Management:

- ✓ Uterine rupture : immediate laparotomy ⇨ if not possible do hysterectomy
- ✓ Impending rupture : immediate caesarean section

230. P212 postpartum presents with a few weeks of spotting and history of bleeding *per vaginum* since 3 months now comes with heavy bleeding. She gives h / o d and c in some other hospital but still continues to bleed. What could be the reason?

a) Normal Lochia
b) DIC
c) Retained bits of placenta
d) Gestational trophoblastic neoplasia

Ans. D

Normal lochia- is present for only first 14 days (no sudden bleeding will be seen in this case)
Retained bits of placenta is removed by **Dilation and curettage** (**D&C**)
Continuous bleeding ***per vaginum*** beyond 12weeks or beyond
Puerperium suspects Choriocarcinoma or Gestational Trophoblastic Neoplasia.

231. Complication associated with iv bolus oxytocin?


a) Hypotension
b) Hypoglycemia
c) Hyperglycemia
d) Hypertension

www.medicalcoupons.in

Ans. A

Oxytocin is given to women during Caesarean section to decrease blood loss. When given as a rapid i.v. bolus, it causes hypotension and tachycardia.

232. Which of the following peaks is responsible for ovulation



a) A
b) B
c) C
d) D

Ans. B

Usually around the midpoint of your cycle, LH secretion surges to really high levels(LH peak). This hormone **surge** is what triggers ovulation about 24 to 36 hours later.

233. A 23 year old primigravida presents at 19 weeks of pregnancy gestation for the first time. She ahs no
significant medical history. Her family history is significant for diabetes running in the family. Her ecg is
normal. Urinalysis is normal
Her FBS- 126 mg/dl
Hba1c- 6.5
Likely diagnosis is

a) Pregestational diabetes
b) Gestational diabetes
c) Normal pregnancy
d) Glucose intolerance

Ans. A

Women with **pregestational diabetes** mellitus defined by fasting ≥7 mmol/L (126 mg/dL) or 2 h ≥11.1 mmol/L (200 mg/dL),hba1c >6.5

234. False statement regarding following image?


www.medicalcoupons.in



a) Do not use electricity
b) Creates a negative pressure of 660 mm hg
c) Capacity is 30cc
d) Can be used up to 12 weeks

Ans. C

Manual vacuum aspiration (MVA) is a way of surgically treating miscarriage when there is pregnancy tissue remaining within the womb. MVA uses a narrow tube to enter and empty the womb using **aspiration** (gentle **suction**).There are two methods of **vacuum aspiration** (also called **suction aspiration**): **Manual vacuum**. This **procedure** can be used around 5 to 12 weeks after the last menstrual period (early first trimester)



235. A 24 year old primigravida attends anc clinic with h / o bp 160/120 mmhg. Her lab findings shows increased liver enzymes and increased LDH. Low platelet count. Diagnosis?

a) AFLP
b) HELLP
c) Hepatitis B
d) Obstetric cholestasis

Ans. B

HELLP syndrome is a potentially life-threatening disorder that's usually associated with preeclampsia

- **H**emolysis
- **EL**: elevated liver enzymes

www.medicalcoupons.in

- **LP**: low platelet count

236. Which type of pelvis is associated with persistent op position?

a) Android pelvis
b) Anthropoid pelvis
c) Gynecoid pelvis
d) Platypelloid pelvis

Ans. B

Android pelvis- MC OP position and associated with deep transverse arrest it's a triangular or a male type pelvis.
Anthropoid pelvis- its anterio posteriorly oval which is associated with persistent op position and face to pubis delivery.
Gynecoid pelvis- MC and the best pelvis.
Platypelloid pelvis- Rarest pelvis, it is Flat and transversely oval, it associated with face presentation and Asynclitism

237. An intern conducting delivery observes mother developing breathlessness, hypotension, tachycardia and collapsing. Shock is out of proportion to bleeding what is the cause?

a) PPH
b) Uterine inversion
c) DIC
d) Amniotic fluid embolism

Ans. B

Uterine inversion means the placenta fails to detach from the **uterine** wall, and pulls the **uterus** inside-out as it exits.Uterine inversion is an unusual and potentially life-threatening event occurring in the third stage of labour. It is associated with significant blood loss, and shock, which may be out of proportion to the haemorrhage, When managed promptly and aggressively, uterine inversion can result in minimal maternal morbidity and mortality.

238. A 25 year old primigravida female is on lithium for treatment of her psychiatric illness for past 2 years . The m/c anomaly seen in fetus whose mother is on lithium during pregnancy is

a) VSD
b) PDA
c) Ebstein anomaly
d) NTD

Ans.

- Lithium causes Ebstein's anomaly if used during pregnancy
- Ebstein's anomaly is a congenital cardiac malformation

www.medicalcoupons.in

Table I : Drugs long recognized to have teratogenic effects in humans at therapeutic doses (continued of facing page)

Drug	Maternal Condition	Most susceptible period post conception	Nature of adverse effect
Ace inhibitors : benazepril, captoprill, cilazpril, enalapril, enaprilat, fasinopril, lisinopril, moexipril, peridopril, quinapril, ramipril,trandolapril	Hypertension	Second or third trimester (13th wk-term)	Oligohydraminios, intrauterine growth retardation, neonatal renal failure, hypotnension, pulmonary hypoplasia, hypoclavaria, joint contracture, death
Amiodarone	Thyroid disorder	10 wk-term	Neonatal thyroid dysfunction or goitre
Aminopterin (≥ 1 -3 mg/d)	Cancer	Organigenesis (18-60 d)	CNS limb and skeletal defects
Antiepileptic drugs : carbamazepine, ethosuximide, phenobarbital, phenetoin, primidone, trimethadione, valporic acid	Epilepsy	Organigenesis (18-60 d)	CNS, cardiac, gastrointestinal and genitourinary defects, facial dysmorphism and digital hypoplasia, growth retardation
Coumarin derivatives: dicumarol, warfarin	Thromboembolic disorders	For CNS defects: unknown: for other defects: second part of first trimester (6-9wk)	Nasal hypoplasia, stipped epiphyses, vertebral abnormalities, CNS and ocular defects, cutaneous hematomas, intracranial hemorrhage, growth retardation, stillbirth
Cyclophosphamide	Cancer, transplant rejection	Organigenesis (18-60 d)	Skeletal and ocular defects, cleft palate
Danazol (≥ 200 mg/d)	Endometriosis, fibrocystic breast disease, hereditary angiodema	Unknown	Virilization of external genetalia in female fetuses
Diethylstilbestrol (1.5-150 mg/d)	Ovarian insufficiency, postcoital contraception	First and second trimester (6-14wk)	Vaginal/cervical carcinoma in females and genetal tract abnormalities in males and females
Indomethascin	Fever, inflammation, premature labour, hydrammios	Second or third trimster (18 wk-term)	Oligohydramnios, anuria, premature closure of the ductus arteriosus, necrotizing entrocolitis
Lithium	Mental illness	Organogenesis (18-60 d)	Cardiac defects, partucularly Ebstein's anomaly of the tricuspid valve

239. A 25 year old women visits labour room at 37 weeks. Pt has mild labor pain for 10 hrs and cervix is
persistently 1 cm dilated and unefaced. Management?

a) LSCS
b) Arm
c) Sedation and wait
d) Augmentation with oxytocin

Ans. C

After sedation false labor pain subsides
Braxton Hicks contractions are the "**false**" **labor pains** that a **pregnant** woman might have before "true" **labor**. They're your body's way of getting ready for the real thing. But they don't mean **labor** has started or is about to begin.Before "true" **labor** begins, you may have "**false**" **labor pains**.

240. A pregnant women with prosthetic valve on warfarin tests positive for pregnancy. What is the advice to be
given?

a) Discontinue warfarin
b) To replace warfarin with heparin
c) Add vit k with warfarin
d) To continue warfarin throughout pregnancy

www.medicalcoupons.in

Ans. B

Warfarin therapy should be avoided during **pregnancy**. If **warfarin**therapy is essential, it should be avoided at least during the first trimester (because of teratogenicity) and from about 2 to 4 weeks before delivery to reduce risk of hemorrhagic complications.

Upto 12 weeks	- LMWH
12-36 weeks	- Warfarin
> 36 weeks	- LMWH

241. A G4 P0+3 , 24 year old female presents at 22 weeks with previous history of three midtrimester abortions,
On usg her cervical length is 20 mm What could be the most probable cause of her recurrent pregnancy loss

a) Cervical Incompetence
b) Uterine Anomaly
c) Hormonal Disturbance
d) Chromosomal Anomalies


Ans. A

Common causes of **miscarriage in** the **second** trimester include cervical insufficiency (the premature dilation of the cervix) or preterm labor (also known as premature birth). With cervical insufficiency (also known as an incompetent cervix), the baby may be born too early to survive.

242. A pregnant women comes for checkup at 18 weeks. On examination it was found that the uterine size
corresponds to only 16 weeks. On usg oligohydrominos was found. Which of the following is likely cause?

a) Fetal Anemia
b) Anencephaly
c) Renal agenesis
d) DM

Ans. C



www.medicalcoupons.in


243. Which is most cost effective cervical cancer screening test?


a) VIA
b) HPV DNATesting
c) PAP smear
d) Colposcopy

Ans. A

Cost effective

1. Lugol iodine
2. Visual inspection of Acetic acid
   BEST :Pap smear





244. A 55 year old post menopausal female attends obgy clinic with history of chronic pelvic pain. An Xray pelvis showed the following image The most likely diagnosis is



a) Ovarian mass
b) PID
c) Calcified fibroid
d) Bladder stone

Ans. C

Leiomyoma is a benign lesion in **postmenopausal** age group. It causes diagnostic confusion with solid **calcified** adnexal mass and large bladder calculi at the pelvic region. A **calcified fibroid** is when a **fibroid** has reached the final stage of degeneration, or cell death and calcium deposits develop on the remaining **fibroid** tissue Clinical and radiological diagnoses were confirmed by histopathology of the hysterectomy specimen.

www.medicalcoupons.in

245. Which of the following have a false positive Barr body

a) Turners syndrome
b) Trisomy 21
c) Klinefelter syndrome
d) Androgen insensitivity syndrome

Ans. C

A **Barr body** (named after discoverer Murray **Barr**) is an inactive X chromosome in a cell with more than one X chromosome, rendered inactive in a process called lyonization, in species with XY sex-determination.
Turners syndrome-45XO
Klinefelter syndrome-47XXY
Androgen insensitivity syndrome-46XY

246. What organs are removed in hysterectomy ?

a) Ovary
b) Fallopian tube
c) Cervix
d) Vagina

Ans. C

A total hysterectomy is the removal of the uterus and cervix. A total hysterectomy with bilateral salpingo-oophorectomy is the removalof the uterus, cervix, fallopian tubes (salpingo) and ovaries (oophor)

247. A female after delivery complains of cramps during breast feeding Which of the following hormone is responsible for it

a) Prolactin
b) Oxytocin
c) Estrogen
d) Progesterone

Ans. B

Nipple stimulation during breastfeeding causes a hormone known as oxytocin to be released into your bloodstream. This hormone causes the contraction of all smooth muscles and helps your uterus contract back into its pre-pregnancy shape and size. These contractions also help reduce postpartum blood loss, so although you may be uncomfortable, this cramping is helping your body heal.

248. After binge drinking of alcohol a young male presented with vomiting and upper Glbleed.Most likely cause is-


a) Boerrhave syndrome
b) Mallory Weiss tear
c) Duodenal ulcer perforation
d) Tension pneumothorax

Ans. B

In Mallory Weiss syndrome, vigorous vomiting produces a vertical split in the gastric mucosa, immediately below the squamocolumnar junction at the cardia in 90% of the cases. In only 10% is the tear in the oesophagus.

www.medicalcoupons.in

Mallory Weiss tear occurs just below GE junction, partial tear involving mucosa and submucosa.
Patient present with painless hematemesis.




249. Following stage is done for which of the following malignancies?



a) Oral cancer - AJCC
b) Bladder cancer- WHO
c) Gastric cancer - Borrmann
d) Color cancer -Duke's


Ans. C



Type 4 (diffuse infiltrate so known as Linnitisplastica aka leather bottle stomach)

www.medicalcoupons.in

250. A 50year old woman presents to the emergency room for nausea, vomiting, loss of appetite and abdominal pain. She has a history of crohn's disease, on infliximab, but she reports that this did not feel like a flare of her disease. On physical examination, there is tenderness to palpation of her abdomen without any peritoneal signs. An abdominal radiograph shows valves of kerckring. She is started in intravenous fluids, given anti-emetics, and put on bowel rest. What is the diagnosis?



a) Distal obstruction in the small intestine
b) Proximal obstruction in the large intestine
c) Pseudo obstruction
d) Distal colonic obstruction

Ans. A

In image we can see dilated bowel loops with valvulae conniventes. Valvulae conniventes is seen in jejunum in means obstruction is in ileum. Bowel loops are placed centrally (in large bowel obstruction bowel loops are placed peripherally and we can see haustration)

Small bowel obstruction	Large bowel obstruction
• Colicky pain • Pain in peri-umbilical region • Multiple episodes of bilious vomiting • Obstipation present	• Colicky pain • In hypogastrium • Distention is seen • Obstipation present

251. A 35 year old patient presented with a one year history of low grade fever, anorexia and weight loss. During investigation with a contrast evaluation of the bowel, following images were seen. Possible diagnosis?

a) Ileocaecal tuberculosis
b) Volvulus
c) Cancer colon
d) Diverticulosis of colon

Ans. A

In image the position of caecum is subhepatic (almost located in the right hypochondrium). Generally the ileo-caecal angle is 90^0. In tuberculosis we see pulled up caecum i.e., obtuse ileo- caecal angle which is known as GOOSE NECK DEFORMITY.

252. A female patient presented to the hospital with complaint of vomiting, diarrhea, wheezing, flushing and palpitations. On laboratory evaluation, high serotonin was found. What is the diagnosis?

a) VIPoma
b) Somatostatinoma
c) Pheochromocytoma
d) Carcinoid syndrome

www.medicalcoupons.in



Ans. D

carcinoid syndrome can produce number of vasoactive peptides, most commonly 5- hydroxytryptamine (serotonin), but also histamine, prostaglandins and kallikrein. When they metastasize to the liver, the 'carcinoid syndrome' can become evident, because the vasoactive substances escape the filtering actions of the liver. The clinical syndrome itself consists of reddish- blue cyanosis, flushing attacks, diarrhea, borborygmi, asthmatic attacks and, eventually, pulmonary and tricuspid stenosis. Classically, the flushing attacks are induced by alcohol.

other options-

VIPoma →is aka Vernal Morrisons syndrome/ WDHA syndrome.

Somatostatinoma→ these patients have DM, steatorrhea and gall stones.

Pheochromocytoma→ headache + diaphoresis + palpitations.

253. In an asymptomatic patient, on routine screening a small mass was detected in the rectum. Carcinoma rectum was suspected. What is the next step in the management?

a) Colonoscopy with lesion biopsy
b) Sigmoidoscopy with lesion biopsy
c) CT pelvis
d) Barium enema

Ans. B


Rigid sigmoidoscopy can be performed in the outpatient clinic and is useful to identify the neoplasm and possibly obtain biopsies. However, it requires the rectum to be empty of faces and may require a prior rectal enema, which may not be practical in the outpatient setting. As colonoscopy is almost always required to visualize the whole colorectum, it is often easier and safer to obtain biopsies at this time.

Proctoscope	10-12 cm
Rigid sigmoidoscope	25 cm
Flexible sigmoidoscope	60 cm
Colonoscope	160

254. What is the site of maximum pain in acute appendicitis

a) McBurney's point
b) Flank pain
c) C)Right upper abdomen
d) D)Umbilicus

Ans. A



www.medicalcoupons.in

255. Patient presented with history of fever and pain in right lower abdomen for last 5 days. On palpitation, painful lump was palpable in right iliac fossa. On laboratory investigations, leukocytosis was found. What is the most probable diagnosis?

a) Hydronephrosis
b) Appendicular lump
c) C.Tuberculosis
d) D.Carcinoma colon


Ans. B

256. A 34 year old female presented with a mass in the inguinal region which produced gurgling sound on reduction. The mass reduced completely on pressing it through the deep inguinal ring. Possible diagnosis in this lady?

a) Indirect inguinal hernia
b) Direct inguinal hernia
c) Femoral hernia
d) Pantaloon hernia

Ans. A

Indirect inguinal hernia
Other option—
Direct inguinal hernia (not seen in females)
Femoral hernia (hernia occurs in femoral ring)
Pantaloon hernia (have dual sac)



257. Very old male presented with a ballotable mass in the left flank region. On examination painless hematuria,
RBC's were found in urine, while there were no WBC's in urine. Most probable diagnosis in this patient?

a) Hydronephrosis
b) Pyelonephritis
c) Renal cell carcinoma
d) D.Polycystic kidney disease

Ans. C

mass+ pain(in lumbar/loin region) + hematuria

258. An old man presented to hospital with low back pain and sclerotic lesion in vertebrae and pelvic bone. What
is the most likely diagnosis in this patient?

a) Testicular tumor
b) Lung cancer
c) Colon cancer
d) Prostate cancer

Ans. D

Prostate cancer metastasis to lumbar vertebra.(pelvic bones are also involved)
RCC, colon cancer—lungs

www.medicalcoupons.in

259. Type of bladder cancer seen in the patient with history of working with dye- related industry?

a) Adenocarcinoma
b) Squamous cell carcinoma
c) Transitional cell carcinoma
d) Papillary carcinoma

Ans. C


260. A patient with history of trauma has ruptured bulbar urethra. What is the location for accumulation of urine?

a) Superficial pouch
b) Deep pouch
c) Retrouterine pouch
d) Uterovesical pouch

Ans. A

In bulbar urethral injury there is superficial extravasation.
Urethral injury is of 2 types
Anterior and posterior (C/F- urinary retention, blood in meatus and hematoma)
In anterior urethral injury – perineal hematoma
In posterior urethral injury – pelvic hematoma

261. 37 years old women came to the hospital with the folloing presentation (image) what is the t-staging ofof the cancer?



a) T4a
b) T4b
c) T4c
d) T4d

Ans. B

pT	
Tis	DCIS LCIS Paget nipple
T1 mic	≤ 0,1 cm
T1a	≤ 0,5 cm
T1b	> 0,5 - 1 cm
T1c	> 1 cm - 2 cm
T2	> 2 cm - 5 cm
T3	> 5 cm
T4a	Extension to chest wall, not including pectoralis muscle invasion
T4b	Ulceration, ipsilateral satellite skin nodules, or skin oedema - including peau d'orange.
T4c	a+b
T4d	Inflammatory ca

pN	
pN1mi	Micrometastasis > 0,2 mm to 2 mm
pN1a	1-3 axillary nodes
pN1b	Internal mammary nodes with microscopic/macroscopic metastasis by sentinel node biopsy but not clinically detected
pN1c	1-3 axillary nodes and internal mammary nodes and internal mammary nodes with microscopic/ macroscopic metastasis by sentinel node biopsy but not clinically detected
pN2a	4-9 axillary nodes
pN2b	Internal mammary nodes, clinically detected, without axillary nodes
pN3a	≥ 10 axillary nodes or infraclavicular
pN3b	Internal mammary nodes, clinically detected, with axillary node(s) or > 3 axillary nodes and internal axillary mammary nodes with microscopic metastasis by sentinal node biopsy but not clinically detected
pN3c	Supra-clavicular

pTNM-Stage	
0	DCIS
IA	T1No
IB	To-1N1mi
IIA	To-1N1 T2No
IIB	T2N1 T3No
IIIA	To-2N2 T3N1-2
IIIA	T4No-2 T3N1-2
IIIC	anyT N3
IV	systemic

www.medicalcoupons.in

262. 60 years old lady presented with a hard breast mass of 5x6cm in size, single lymph node mass present. There was no evidence of metastasis? What should be the ideal treatment?

a) Radical mastectomy with chemotherapy
b) Modified radical mastectomy followed by adjuvant chemotherapy
c) Neoadjuvant chemotherapy with Modified radical mastectomy with radio ther...
d) Palliative mastectomy.

Ans. A

cTNM	Neo-adjuv	Sentinel Node	BCT	Mastectomy
DCIS	**No**	When possible invasive component > 4cm < 55 years Solid component on mammogram DCIS grade II and III	Excision possib with good cosmetic result Unifocal lesio Multifoc ithi 5 cm DCIS < cm	Go cosmetic re lt not possible Multicentric DCIS > 4 cm
Localized Cancer Tumor < 5 cm ***and*** < 4 axill nodes	**Sometimes** to make a BCT possible in large tumors	**Always** unless US-guide d fine needle aspiration is positive	E cision possible with g od sm etic result Unifocal lesion M ltifocal within 5 cm Invasive Ca < 5 cm	Good cosmetic result not possible with BCT Multicentric Invasive Ca > 5 cm
Loco-regionally Advanced Tumor > 5 cm ***or*** > 4 axill nodes	**Yes** to make surgery possible	**No** reliable	Usually mastectomy is performed. Sometimes BCT is possible.	

263. A women with breast malignancy as shown in image,it is not associated with lymph node metastasis, not movable and attached to the rib ,what is the diagnosis?



a) Phyllodes tumour
b) Brest abscess
c) Inflammatory breast cancer
d) Leiomyosarcoma

Ans. A

www.medicalcoupons.in

264. A lactating female with history of breast feeding for 15 days presented to emergency with fever. On examination, breast was red as shown in the image .what is the probable diagnosis?



a) Breast abscess
b) Milk fever
c) Tuberculosis
d) None of the above

Ans. A

265. Which of the following radio-isotop is used for carcinoma thyroid?
a) I-123
b) I-125
c) I-127
d) I-131

Ans. D
I-123 =13 Hrs used for diagnosis RAI(radio active iodine up take scan)
I-131 =8 days used for diagnostic and therapeutic
Also used for Radio active ablation

266. A patient came with the pain in right hypochondrium .bloodinvestigantion shows AST-22 IU/l and ALT-27 IU/ L,serum bilirubine levels -6mg% and ALP-200 mg % what should be the investingation in thus patient
a) USG abdomen
b) CECT abdomen
c) Hepatitis A and Hepatitis B markers
d) HIDA scan

Ans. A

267. A female patient presented to the hospital with history of recurrent pain in the right hypochondrium for 1 year. what is the diagnosis?
a) Hydatid cyst
b) Acute pancreatitis
c) Acute cholecystitis
d) Amobic liver abscess

Ans. C

www.medicalcoupons.in

268. A 60 years old male who is heavy alcoholic present with 12 hours history of sharp central abdominal pain and vomiting, preferred to nsit, rather lying down as pain increases when lying down. pulse rate 132/min, bp 118/82 mmhg, temperature is 38.3 c. on examibnation he has mild jaundice. There is periumbilical and flank discoloration as shown in the image and minimum tenderness in epigastrium. What is the diagnosis?

a) acute pancreatitis
b) pyelonephritis
c) acute viral hepatitis
d) perforated duodenal ulcer

Ans. A

key points

BLUISH DISCOLERATION OF THE FLANKS =GREY TURNER SIGN
DIS COLERATION AROUND UMBILICUS =CULLEN SIGN


269. Alcoholic male presented with pain in epigastrium radiating to the back. What is the likely diagnosis in the given CT image ?



a) Amoebic liver abscess
b) Pancreatic pseudocyst
c) Acute cholecystitis
d) Hydatid cyst

Ans. B

270. What is the name of surgery, whose image is given below?




www.medicalcoupons.in

a) Pancreatiojejunostomy
b) Whipples surgery
c) Pancreaticoduodenostomy
d) Duodenojejunostomy

Ans. A
also called : PUESTOW'S PROCEDURE LATERAL PANCREATIOJEJUNOSTOMY
Involves a retrocolic side-to-side roux-en-ypancreaticojejunostomy

271. What is the name of procedure, whose image is given below?



a) Pancreaticojejunostomy
b) Whipples procedure
c) Radical pancreatectomy
d) Duodenojejunostomy

Ans. B

272. In a new born male baby mother found empty scrotum.on examination, left testis is in inguinal region and
right testis is in perineum. Which of the following is correct?
a) Right testis is undescended and left testis is ectopic
b) Left testis is undescended and right testis is ectopic
c) Both are undescended
d) Both are ectopic

Ans. B
left – is at inguinal region –undescended
right is at perineuinal- ectopic

273. In a new born bady with empty scrotum, testis was was found to be laying close to the peritoneum.which of
the following helps the testis to reach the scrotum?
a) GnRH analogues
b) Testosteron
c) Cord compression
d) Manually bring testis down to the bottom of scrotum

Ans. A
Maternal chorionic gonadotrophin growth of the testis and may stimulate its migration so we give baby GnRH analog that helps the testis to reach the scrotum


www.medicalcoupons.in

274. 18 years old boy presented in agony with sudden onset of severe pain in the groin region associated with redness and swelling of the scrotum. On examination the right testis as gone up. Possible diagnosis in this boy is?


a) Testicular torsion
b) Hydrocele
c) Strangulated inguinal hernia
d) Acute epididymo-orchitis

Ans. A


Testicular torsion:



Hydrocele:




Strangulated inguinal hernia:



www.medicalcoupons.in

epididymo-orchitis:



275. A 35 years old male presented with scrotal enlargement on investigation, LDH was significantly high but the HCG and alpha fetoprotin were within normal limits .histopathology image is shown below what is the most likely diagnosis?

a) Teratoma of testis
b) Seminoma of testis
c) Choriocarcinoma
d) Yolk sac tumor

Ans. B

276. A young male underwent lumbar sympathectomy and bilateral L1 ganglions were removed. What is the most commion side effect?


a) Impotence
b) Retrograde ejaculation
c) Bladder dysfunction
d) Incontinence

Ans. B

patients with **burgers disease** who have rest pain we perform lumbar sympathectomy,
in this we remove L1 and L2 ganglion on one side
if we remove both sides the patient will have **retrograde ejaculation**
it can be prevented by sparing of L1 at one side

277. A patient underwent surgery, from which she was immobilized.patient started complaining of leg pain on the 5th post operative day, on examination Homans sign was found to be positive. What is the most probable diagnosis?

a) Deep venous thrombosis
b) Urinary sepsis
c) Peripheral vasculsar disease
d) Lymphangitis



Ans. A

www.medicalcoupons.in

278. A patient with old burn scar develop symptoms over burnt area . it was indicative of malignancy.what is the type of malignancy?



a) Basal cell carcinoma
b) Squamous cell carcinoma
c) Keratocanthoma
d) Sebaceous horn

Ans. B
given condition-MARJOLINS ULCER

279. A patient with blunt trauma was brought to the emergency with the pulse rate 120/min bp80/50. What is the next best step?
a) FAST
b) DPL
c) Exploratory laprotomy
d) CECT

Ans. A
Patient whether stable or unstable first FAST should be done
(FAST-focused assessment with sonography for trauma)

280. A patient of blunt abdominal trauma was brought in the emergency with cold clammy skin and increased puls rate. What is the next step of management?
a) CECT
b) Exploratory laprotomy
c) Resuscitation with blood products
d) Resuscitation with intravenous crystalloids


Ans. D
given only one liter of warm isotonic crystalloids ***for adults***
child <40kg given 20ml/kg of warm isotonic crystalloids

281. A patient underwent a surgery for ingunial hernia 5 days before. Now comes to the hospital with complains of fever. What can be the cause?
a) Atelectasis of lung
b) Surgical site infection
c) Thromboembolism
d) Chest infection

Ans. B
cause of post-op fever
POD 1-2: Atelectasis
POD 2-3: UTI
POD 3-7: Wound infection (ssi)
POD 5-7: DVT/Thrombophlebitis
POD >7: drug fever

www.medicalcoupons.in

282. Identify the movement taking place in the intestine?



a) Segmented movement
b) Peristaltic movement
c) Passive diffusion
d) Intersegmental contraction

Ans. B

GI – Motility

↓ Types

2- types

Food

Major inititoing response for persislassis is – local stretching of gest

Yes — Fed state pattern food

No — Fasting state pattern

Para & of each

→ (Food) (F)

Contraction felaxation (2)

1 Peristalasis 'food'

Oral ⟶ Anal

food

Contraction

1 Arch – I

2 Substance – P

↑↑↑↑ Ach ⟷ Balance

↳ LES → Absence of

No

Relaxation

Nitric Oxide (No)


Rx :

Inhibits ⇒ ← Botulinum toxin

Achalasa Cardia


www.medicalcoupons.in

283. Pressure volume graph given and asked to find out B accordingly?




a) Opening of aortic valve
b) Opening of mitral valve
c) Closing of aortic valve
d) Closing of mitral valve

Ans. A



284. Which part of nephron does water reabsorption takes place more?



a) A
b) B
c) C
d) D

www.medicalcoupons.in

Ans. A

About 2/3rd of the filtered water is reabsorped in proximal convoluted tubule along with salt absorption.other molecules absorped here are bicarbonates, glucose, amino acids and phosphates.

285. Inhibition of Na+ - K+ ion channel results in which of the following?

a) First depolarise and then hyperpolarise
b) Hyperpolarisation
c) Depolarisation
d) Repolarisation

Ans. C

286. Hormone that regulate serum calcium levels

a) Calcitonin
b) Vitamin D
c) Parathyroid hormone
d) All

Ans. D

Calcium Regulation :

3 HORMONES
1. Parathormone
2. Vit D
3. Calcitonin →

↑Ca^{2+}
↓Ca^{2+}

3 organs modal
↓
1. Bone (Bar... for Ca^{2+})
2. GI...
3. K...dneys

Parathormone (Action)

1. Bone → 2 cells PTH
Osteo blast
PTH receptor ↑
↓
RANK ligand
Rx : Osteoporosis
↓
Rank Ligand Bl...
DENOSUMAB

PTH
Oste..Clast → Bone resorption
+
Ca^{2+} - PO_4^{2-}
Crystal

↑Ca^{2+} ↑PO_4^{2-}
Bone (PTH)

2) Kidneys G→ PTH
↓ Excreation o... Ca^{2+}
↑ Excreati... n of P... $^{2-}$
PTM
↓ Phosphaturic action
Over all ...tion
PTH → PO42- Terminating Hormone

3) GIT → ... Vit D → ↑ Ca2+ absortion
&
PTH + Indiract action

2... 7. Why can't hemoglobin pass through glomerulus?

... Large molecule bounded with protein
b) Cone of protein decrease glomerular capillaries
... Negative charge of bowmans capsule
d) None of the above

Ans. A

www.medicalcoupons.in

Protiens → not filted by GC Glossullary capillary
3 reason

1. Charge — Repals -Ve — Protein Anions

Gc
?
Heparan, sulfate
Heparan SO_4^{2-}
(Proteoglycans)

2. Size
GC – Small gap
Fenestrations → Fenestrated Endothelium
10 – 90 nm

3. Slit diaphram :
Structure
Proteins
i. Neprin → NPHS 1 geno
mutation
Congenital. N syndrome = (protein Excretion)
↓
Finish type

ii. Podocin → NPHS2
↓ gen
Mutation
AR. Steroid resistant Nephrotic syndrome ARSRNS

iii. Alpha actin
→ Mutation
FSGS. Focal Segmeatal glomullory sclerosis

288. In gestational life surfactants are produced by which cells?
a) Type I pneumocyte
b) Type II pneumocytes
c) Basal cells

Ans. B

Type I pneumocytes	Type II pneumocytes
Large **Less in number** **More area**	**Small** **More in number** **SURFACTANT** **Stem cells**

Surfactant :
Lipids (major—lecithin), (minor –spingomyelin)
Protein :A,B,C,D
In amniotic fluid—bubble test, shake test, clement test.
Function of surfactant:

www.medicalcoupons.in

- ↓ in surface tension
- Prevent collapse of alveoli
- Stability of alveoli
- SP : A and D → lung maturity

Surfactant stored in Lamellar body.

Regulation of surfactant :

↑surfactant : steroids, glucocorticoids, cortisol, T3 and T4.

289. In child which appears first?

a) Palmar grasp
b) transfering objects
c) uni dexterous approach
d) bi dexterous approach

Ans. A

Palmar grasp (immature) should logically also appear first follo d by nsferring objects
Bi dexterous appear by 4th month
Unidextrous appear by 6th month

290. A 4 years old child,presented with thisty,drink eager'y,irritable with dehydration, what the treatment ?

a) ORS
b) Ringer lactate
c) Plain normal saline
d) Refer to district hospital

Ans. A

DEHYDRATION:

Skin pinch test

NO dehydration	No sign / symptoms	< 1 sec
Some dehydration	Thirst ++,Irritabi	slow <2 sec
Severe dehydration	leth rgi ina tiv	very slow >2 sec

Calculation of fluid by H LIDAY SEGAR REGMIN
First 0-10 kg = 100 /kg
Next 10-20 kg +50 m /kg
>20 kg = +20 m /kg

TREATM NT PL AN FOR DEHYDRATION:

N dehydration	Some Dehydration	Severe dehydration (lethargy)
PLAN A	PLAN B	Plan C

www.medicalcoupons.in

• ORS 100-200ml/loose stool (ongoing losses 10-20ml/kg)	• Rehydration (75ml/kg in 1st 4 hrs) + maintenance (replacing on going loss-10-20ml/kg)	• Start IV fluids • Ringer lactate >>Normal saline Give 100 ml/kg as : Age | 30ml/kg | 70ml/kg <1yr | 1hr | 5 hr >1yr | 30 min | 2 ½ hr

Patient comes with an enlarged leg with sulphur crystals coming out (madura foot as shown in the image). Organisum responsible is:



Explanation :
Actinomycotic mycetoma: Sinus ,swelling,granules(discharging)

Actinomycetoma	**Eumycetoma**
➢ Numerous sinus and swelling ➢ Colour of granules→ yellow/white except in Actinomadural pellitri when they have red granules ➢ Respond to antibiotics	➢ Single sinus and swelling ➢ Black →madurella mycetomatis ➢ White ➢ Acremonium ➢ Pseudallescheria boydii ➢ Amputation

CTEV






✓ Denis brown splint ⇨used to correct foot
✓ **Use** C A V E⇨ C: cavus
A: adduction
V: varus correction
E: equines (tibia and thalus involved)
Keep the foot in over corrected position


www.medicalcoupons.in

Right Midble Lobe Consolidation




- ✓ Image 1: Area of consolidation middle lobe which silhouette right heart border
- ✓ Image 2: cardiac shadow is seen on the middle lateral lobe

Diffuse Esophageal Spasm



- ✓ Cork-screw/rosary bead esophagus on barium studies

Q. Implantable Cardioverter-Defibrillator [Icd]



Plain radiograph

- ✓ For implantable cardio-defibrillator (ICD) , a chest x-ray should be obtained to confirm lead position and examine lead integrity
- ✓ Most pulse generators, both pacemakers and ICD's, can be identified from their appearance under radiography: most helpful method of identifying unknown devices

www.medicalcoupons.in

Q. What is the radioisotope used in PET scan ?

Ans: 18FDG

ANNEXURE 5. HALF – LIVES OF RADIOACTIVE ISOTOPES	
Substance	**Half – life**
Iodine – 132	2.3 hours
Technetium – 99	6 hours
Iodine – 123	13 hours
Gold – 198	2.7 days
Gollium -67	3.3 days
Iodine – 131	8 days
Phosphorus -32	14.3 days
Iodine -125	59 days
Iridium – 192	74 days
Cobalt -60	5.3 days
Strontium – 90	29.1 years
Caesium -137	30 years
Radium – 226	1600 years
Uranium – 238	4.5 billion years


✓ Half- life: 110min

Q. Oneliner: To know correct position of IUCD ⇨ USG is best

Q. Identify the cysyt?



Q. Identify the fibroid .




www.medicalcoupons.in

FMGE DEC 2021 RECALLS

1. A patient having the history of leprosy has presented with the deformity as shown in the image. Which nerve is involved in this case?

A. Axillary nerve
B. Radial nerve
C. Median nerve
D. Ulnar nerve



2. Identify the bone marked in the given x-ray



A. Talus
B. Navicular
C. Cuboid
D. Cuneiform


www.medicalcoupons.in

3. Which type of joint is present at the pointed area in the given X-Ray



A. Primary cartilaginous
B. Secondary cartilaginous
C. Fibrous
D. Synovial

4. Identify the marked structure?



A. Vocal fold
B. Epiglottis
C. Esophagus
D. Tongue

www.medicalcoupons.in

5. **A Patient with Supraspinatous Tendinitis, what degree of movement would be painful?**
   A. 15 degree
   B. 90 degree
   C. 60 degree
   D. 30 degree

6. **What is the dermatome for the pointed area by the arrow?**



A. L2
B. L3
C. L4
D. L5

www.medicalcoupons.in

7. A newborn is having a congenital defect shown in the picture. Find out the embryological developmental defect out of the following options?



A. Failure in closure of anterior neuropore of neural tube
B. Failure in closure of posterior neuropore of neural tube
C. Failure in closure of complete neural tube
D. Failure in closure of caudal neuropore of neural tube


8. Which structure is palpated in the given image?



A. Anterior tibial artery
B. Posterior tibial artery
C. Plantar artery
D. Dorsalis pedis artery


www.medicalcoupons.in

9. A patient present with the symptoms (below the level of lesion) as shown in image. What is the most probable reason for this?



A. Brown Sequard syndrome
B. Wallenberg syndrome
C. Millard Gubler syndrome
D. Weber's syndrome


10.Identify the marked structure?



A. Right atrium
B. Right ventricle
C. Left atrium
D. Left ventricle


www.medicalcoupons.in

11.Identify the arrow marked structure the image below?



A. Primordial follicle
B. Primary follicle
C. Preantral follicle
D. Antral follicle


12.Identify the correct location of auscultation areas for heart?



A. a-aortic area, b-pulmonic area, c- tricuspid area, d- mitral area
B. a-pulmonic area, b-aortic area, c- tricuspid area, d- mitral area
C. a-aortic area, b-pulmonic area, c- mitral area, d- tricuspid area
D. a-aortic area, b- tricuspid area, c- pulmonic area, d- mitral area

www.medicalcoupons.in

13.Identify the arrow marked structure in the transverse section shown below?



A. Trachealis
B. Posterior inter-arytenoid
C. Circular muscle layer of esophagus
D. Longitudinal muscle layer of esophagus

14.The section along the frontal plane is termed as?

A. Sagittal section
B. Coronal section
C. Transverse section
D. Oblique section

15.Which epithelium is present in fallopian tube?

A. **Ciliated columnar epithelium**
B. Simple cuboidal epithelium
C. Simple squamous epithelium
D. Pseudostratified epithelium

16.What is the location of neurovascular bundle in anterior abdominal wall?

A. Between external oblique & internal oblique
B. Between internal oblique & transversus abdominis
C. Between internal oblique & fascia transversalis
D. Between transversus abdominis & transversus abdominis

www.medicalcoupons.in

17.A patient was referred by dr. to a radiologist for CT scan. Radiologist later visited the physician with a thank you letter & money envelop. Which is true regarding above situation.

A. Drs. should be punished under law
B. Medical dichotomy
C. Association of Doctor
D. Covering

18.What type of toxin is released by the snake shown below?



A. Hematotoxin
B. Myotoxic
C. Neurotoxin
D. Vasculotoxic

19.Which of the following is the true statement from the image given below:



www.medicalcoupons.in

A. A case of homicide
B. A case of suicide
C. Rigor mortis
D. Cadaveric spasm

20. What type of odour is produced by the decomposed body shown in the image?



A. Ammoniacal
B. Odorless
C. Offensive
D. Rancid

21.Identify the test shown in the image below:



www.medicalcoupons.in

A. Barberio's test
B. Florence test
C. Takayama's test
D. Teichmann's test

22. A couple did not want a girl child and aborted by using an abnormal method What could be the method used for criminal abortion?

A. Vacuum aspiration
B. Prostaglandins
C. Dilatation & curettage
D. Abortion stick

23. Finding in the image shown is suggestive of



A. Scar
B. Tattoo
C. Mole
D. Ligature mark

24. A male patient presented hospital with tachycardia, hypertension, palpitations and pupil dilatation. He is found to be a substance abuser. Which of the following drug withdrawal can cause the following symptoms?

A. Alcohol
B. cannabis
C. Cocaine
D. Opioid

www.medicalcoupons.in

25.Number of maternal deaths per 1,00,000 live birth is

A. Maternal Mortality Rate
B. Maternal Mortality Ratio
C. IMR
D. Under-5 Mortality Rate

26.In a population of 10,000; What does it mean that the sex ratio is more than 1000?

A. Male population less than 500
B. Male population less than 5000
C. Female population less than 500
D. Female population less than 5000

27. Couple has given birth to their first child, the family is entering in which phase


A. Formation
B. Expansion / Extension
C. Contraction
D. Dissolution

28.In 14th century there was a disease called plague, quarantine of ship sailor was done for how many days?

A. 7 days
B. 14 days
C. 40 days
D. 60 days

www.medicalcoupons.in

29. What is the percentage of 1 SD in the given graph?



A. 34%
B. 68.3%
C. 4.6%
D. 95.4%

30. After dressing of a patient of RTA Patient, cotton swab will be disposed in which colour bag

A. Yellow
B. Red
C. White
D. Black

31. A worker posted in the same village of age 25-45 years, 8th std pass, widow, with good communication skill. Identify the worker

A. MPW
B. Accredited social health activist
C. Anganwadi worker
D. Avachi /TBA

32. Pregnancy kit for follow up at home

A. NIKSHAY
B. NISHCHAY
C. Free Days
D. Today

www.medicalcoupons.in

33.Burden of disease is

A. DALY
B. SULLIVAN INDEX
C. HALE
D. QUALY

34.For teaching of how to prepare ORS in urban slum area which appropriate method will be used

A. Role play
B. Demonstration
C. Group discussion
D. Flash card

35.If you want to explain nutritive value of various food items to a patient with help of chart, the following chart is?

A. Histogram
B. Bar chart
C. Pie chart
D. Pictogram

36.A 40-year patient from Rajasthan working in stone breaking industry presenting with dry cough, weight loss, with possibility of TB, which pneumoconiosis is associated with his condition

A. Silicosis
B. Asbestosis
C. Byssinosis
D. Bagassosis

37.If you want to check for incidence of deafness caused by noise effect in office workers and factory workers after 1 year, study conducted will be

A. Case control
B. Cohort
C. Ecological
D. Demographic

www.medicalcoupons.in

38.Land area required to fill trench of 2-meter depth with compacted refuse for population of 10,000 is

A. 1 acre
B. 2 acre
C. 4 acre
D. 3 acre

39.Maximum number of candidates in group discussion to avoid overcrowding is

A. 3-4
B. 4-8
C. 6-12
D.12-15

40.Reinfection in presence of antibody in blood will flare up symptoms in

A. Measle
B. Rubella
C. Dengue
D. Malaria

41.A child presenting with fever, swelling in neck and testis, 2 more children in village were having same symptoms, diagnosis is

A. Measle
B. Rubella
C. Mumps
D. Pertussis

42.If you want to reduce selection error in a clinical trial conducted among two groups, best method will be

A. Matching
B. Blinding
C. Randomization
D. Stratification

www.medicalcoupons.in

43.Evaluation of birth and death rate in National Level by

A. Civil registration
B. SRS
C. NFHS
D. Census

44.A Farmer comes to emergency with complaints of his wife consuming Bajra seeds which were abnormal and discoloured and she developed allergy, it was found to be adulterated with black grains. how can we separate the adulterant from the food?

A. Distilled H2O
B.10% sodium
C. 20% sodium
D. Normal Saline

45.In rural areas, patient don't have insurance. But under ESI. Registered medical practitioner can give which of the following service to beneficiary

A. Direct medical benefit
B. Indirect medical benefit
C. Sickness benefit
D. Extended sickness benefit

46.Lung malignancy is associated with

A. Silicosis
B. Asbestosis
C. Bagassosis
D. Byssinosis

47.A 2-year-old child reported to PHC with complains of fever and breathlessness. On examination respiratory rate is 36 /min and presence of chest indrawing. management protocol is

A. home management
B. Antibiotic & Follow up
C. Immediate referral
D. Antibiotic and immediate referral

www.medicalcoupons.in

48.All are true for vasectomy except

A. Immediate sterilization
B. Permanent method
C. Contraceptive result after 3 months

49.Information specially of a biased or misleading nature used to promote a political cause or point of view is

A. Propaganda
B. Health Promotion
C. Advocacy
D. Health Education

50.Lowest Glycaemic index is of

A. Watermelon
B. Papaya
C. Sweet corn
D. Potato

51.More sodium content in

A. Soup
B. Green vegetable
C. Bread
D. Canned food

52.A 67 years old hypertensive man presents with acute painless loss of vision. examination reveals v/a hand movements & afferent pupillary defect. On examination Fundus was pale in colour and Macula was dark. Probable diagnosis is: -

A. CRAO
B. CRVO
C. Cystic edema
D. Macular edema

www.medicalcoupons.in

53.A female feeling weak having dysphagia, retrosternal pain, regurgitation, Cough and burning sensation. What is the most likely diagnosis?

A. Paterson Kelly disorder
B. Thyroid Disorder
C. Reflux Esophagitis
D. Barrette Esophagus

54.Chronic smoker for 20 years came to OPD with the complaints of following Image. What is the diagnosis?



A. Buerger's disease
B. Fournier's gangrene
C. Reynaud's phenomenon
D. Wet Gangrene

55.The following instrument is used in?



A. Dacryocystorhinostomy
B. Ectropion
C. Entropion
D. Chalazion

www.medicalcoupons.in

56.A person enters in a theatre and there is dim light. What are the pupillary changes seen in him?

A. Pupil dilatation with 11 cis retinol connecting scotopsin
B. Pupil dilation with 11 cis retinoid connecting scotopsin
C. Pupil dilation with all cis retinol connecting scotopsin
D. Pupil dilatation with all cis retinoid connecting scotopsin

57.V/A improves on looking through pinhole. What disorder does this indicate for the patient to have?

A. Refractive errors
B. Retinal diseases
C. Macular diseases
D. Astigmatism

58.A farmer presented with pain, photophobia & watering from eye since last 36 hours. On examination 1x1 cm ulcer is seen as shown in image. Identify the lesion marked



A. Satellite lesions
B. Hypopyon
C. Dendrite ulcer
D. Ring infiltrate

www.medicalcoupons.in

59.A child comes with defect in vision but fundus is normal. Diagnosis is. -

A. Astigmatism
B. Amblyopia
C. Myopia
D. Hypermetropia


60. A child presents to an ophthalmologist with complaint of hazy vision while trying to watch the blackboard in the class room. The doctor prescribes him concave lens. What could be the diagnosis?

A. Myopia
B. Hypermetropia
C. Astigmatism
D. Night blindness

61.M/C cause of blindness in children is

A. Cataract
B. Glaucoma
C. Vitamin A deficiency
D. Refractive Error

62.Identify the condition: -



A. Entropion
B. Ectropion
C. Distichiasis
D. Trichiasis

www.medicalcoupons.in

63.A child 1.5 years comes with leukocoria and Flexner winter Steiner rosettes. Most probable diagnosis is:



A. Retinoblastoma
B. Rhabdomyosarcoma
C. Congenital cataract
D. Congenital glaucoma

64.After a catastrophe from metal works, a mechanic got injured from metal foreign body. I.O.C. is

A. MRI
B. CT
C. B Scan
D. Nuclear radiology

65.All Extra Ocular Muscles are paralyzed. Most probable site of lesion is

A. Upper medulla
B. Pons
C. Lower medulla
D. Midbrain

www.medicalcoupons.in

66.Difference in Visual Acuity between two eyes. Most probable cause is:

A. Refractive errors
B. Anisometropia
C. Amblyopia
D. Anisometropic

67.A patient comes with History of contact lens use complains of foreign body sensation. Most probable diagnosis?

A. Corneal abrasion
B. VKC
C. Giant Papillary Conjunctivitis
D. Keratitis

68.Patient had a fight & he suffered a blow to the eye and now presents with non-reactive pupil with painful ocular movements. Most probable diagnosis-

A. Cataract
B. Optic neuritis
C. Glaucoma
D. Papilledema

69. Trotters' triad includes all except?
A. Conductive hearing loss
B. Retroorbital pain
C. Palatal palsy
D. Pain in the area of distribution of 5th nerve.

70. A Patient presented with whitish membrane that bleeds on scraping What is the diagnosis:
A. Diphtheria
B. Ludwig Angina
C. Mumps
D. Rubella

www.medicalcoupons.in

71. While examining a patient's External ear syncopal attacks occurred. Which nerve is responsible for this?

A. Jacobson's nerve
B. Glossopharyngeal nerve
C. **Arnold's nerve**
D. Cauda Tympani

72. A 65-year-old Female patient presented with a red mass behind tympanic membrane with pulsatile tinnitus the diagnosis could be

A. Cholesteatoma
B. **Glomus tumor**
C. Otosclerosis
D. Meniere's disease

73. A 15 years old male presented with bleeding from nose and on examination there is a mass, probable diagnosis is

A. Rhinosporidiosis
B. **Juvenile Nasopharyngeal angiofibroma**
C. Nasopharyngeal ca
D. Rhinoscleroma

74. A young boy presented with mouth breathing, failure to thrive high arched palate, and hearing loss. Management is

A. Adenoidectomy with grommet
B. Myringotomy
C. Myringotomy with grommet
D. Adenoidectomy

75. A 50-year-old patient of type 2 diabetes mellitus was controlled on oral hypoglycemic drugs. This patient presented to hospital with hypoglycemia. Which among the following drugs can result in hypoglycemia in this patient?

www.medicalcoupons.in

A. Metformin
B. Vaglibose
C. Vildagliptin
D. Glipizide

76. A patient is taking platinum based anti-cancer therapy for treatment of ovarian cancer. Which of the following drug should be used to prevent the nausea and vomiting in this patient?

A. Domperidone
B. Promethazine
C. Palonosetron
D. Metoclopramide

77. In a patient with pregnancy induced hypertension, Drug used for treatment is

A. Hydralazine
B. Alpha methyl dopa
C. Lisinopril
D. Telmisartan

78. A female patient was treated with clindamycin and developed diarrhea most probably pseudomembranous colitis. Which of the following is the likely organism responsible for this?

A. Clostridium perfringens
B. Clostridium difficile
C. Clostridium tetani
D. Clostridium septicum

79. A 56-year-old patient who is a known case of hypertension was being treated with antihypertensive drugs. He is having chronic cough since 2 weeks. He was started on antimicrobials but cough did not resolve. Which of the following antihypertensive drug may induce chronic cough in this patient?

A. Lisinopril
B. Minoxidil
C. Propranolol
D. Amlodipine

www.medicalcoupons.in

80. A 30-year-old man travelled to Assam. After coming back to his hometown Delhi, he developed fever, chills and rigors and altered sensorium. The peripheral smear demonstrated the presence of P. falciparum. What is the treatment of choice for this patient?

A. Chloroquine, Sulfadoxine pyrimethamine
B. Intravenous Artesunate
C. Quinine
D. Primaquine

81. A patient with chronic obstructive pulmonary disease is on salmeterol therapy but there is no improvement. Which of the following drug should be added to the regimen?

A. Inhaled Corticosteroids
B. Tiotropium
C. Oxygen therapy
D. Promethazine

82. A person presented to emergency in comatose state after consuming some unknown poison. Excessive sweating is apparent and on examination there is bradycardia and BP is 80/60 mmHg. Which of the following is antidote for treatment of this patient?

A. Neostigmine
B. Atropine
C. Physostigmine
D. N-acetylcysteine

83. A patient has developed tuberculosis in which mycobacterium is resistant to rifampicin, isoniazid, kanamycin and levofloxacin. This tuberculosis is referred to as

A. MDR TB
B. Rifampicin resistant TB
C. Extensively drug resistant TB
D. Extremely drug resistant TB

www.medicalcoupons.in

84.A patient presented with unilateral hemiparesis and presented to hospital after 8 hours. On examination and investigations, the blood pressure is 220/110 mmHg and HbA1c is 8%. CT scan of brain shows no hemorrhage. Which is the appropriate management of the patient?

A. Conservative management
B. tPA
C. IV labetalol
D. Sublingual nifedipine

85.A 25-year-old male presented to emergency with head trauma due to road traffic accident. In hospital, the patient developed seizures. Emergency CT scan revealed widespread cerebral edema. Which of the following is diuretic of choice for cerebral edema in this patient?

A. Mannitol
B. Spironolactone
C. Furosemide
D. Hydrochlorothiazide

86.A 28-year-old male went to Ladakh. He developed shortness of breath. Which of the following drug will you use for the treatment of mountain sickness at high altitude?

A. Acetazolamide
B. Hydrochlorthiazide
C. Eplerenone
D. Salbutamol

87.A young boy was brought to hospital with complaints of few episodes of fast blinking and staring at something for around 20 seconds. ECG shows spike and wave pattern. Which of the following drug should be used for treatment?

A. Phenytoin
B. Sodium valproate
C. Diazepam
D. Carbamazepine

www.medicalcoupons.in

88. A professor in medical college is teaching students about the adverse effects of a drug. Which of the following drug can cause gynecomastia in men and menstrual irregularities in women?

A. Spironolactone
B. Propranolol
C. Misoprostol
D. Oxytocin

89. What is the mechanism of action of tPA?

A. Promote Fibrinolysis
B. Inhibition of factor V and VII
C. Inhibition of factor III and V
D. Activation of anti-thrombin III

90. Which of the following drug will you omit from the prescription of a patient with XDR tuberculosis in pregnancy?

A. Kanamycin
B. Cyclosporine
C. Moxifloxacin
D. Linezolid

91. A 30-year-old female presented with weight gain and cold intolerance. On investigation, Value of T3 is 70 ng/dl (reference value 100-200 ng/dl), T4 is 3 mcg/dl (reference value 5-12 mcg/dl) and TSH is 20mIU/L (reference value 0.5-5 mIU/L). Which of the following statement is correct regarding the management of this patient?

A. Measure T3, T4 and TSH after one month
B. Start L-thyroxine (T4) treatment 100 mcg once daily
C. Start daily L-thyroxine (T4) 100 mcg with Liothyronine (T3) 5
D. Start L-thyroxine (T4) treatment 25 mcg daily and gradually

92. A 72-year-old patient with mitral valve disease was started on long term warfarin therapy. What should be the target INR value in this patient?

www.medicalcoupons.in

A. Less than 1
B. 1-2
C. 2-3
D. 3-4

93. A 50-year-old female presented to oncologist with breast lump. On examination and investigations, breast cancer was diagnosed. Echocardiography of the patient shows ejection fraction of 25 %. Which of the following drug will you avoid in this patient?

A. Epirubicin
B. Irinotecan
C. Cyclophosphamide
D. Methotrexate

94. Which of the following is an adverse effect of orlistat?

A. Suicidal tendency
B. Flatulence
C. Weight gain
D. Hyperglycemia

95. Which of the following statements about the anti-psychotic drug, Aripiprazole is most appropriate?

A. It is a D2 and 5HT 1A partial agonist with less extrapyramidal symptoms and less weight gain as compared to other anti-psychotic drugs
B. It is a D2 and 5HT 1A partial agonist with more extrapyramidal symptoms and more weight gain as compared to other anti-psychotic drugs
C. It is a D2 antagonist and 5HT2 agonist with less extrapyramidal symptoms and less weight gain as compared to other anti-psychotic drugs
D. It is a D2 antagonist and 5HT2 agonist with less extrapyramidal symptoms and less weight gain as compared to other anti-psychotic drugs

96. A 5-year-old child presented with complaints of hematuria with history of sore throat 2 weeks. On assessment, no fever and cola colored urine was noted. What is the diagnosis?

www.medicalcoupons.in

A. PSGN
B. Membranous nephropathy
C. IgA Nephropathy
D. Minimal change disease

97. A 10-month-old is undergoing treatment for diarrhea with hypokalemia. However, in spite of potassium correction, the child does not improve and floppiness is noted. Which of the following condition should be considered?

A. Albumin deficiency
B. Phosphate deficiency
C. Calcium deficiency
D. Magnesium deficiency

98. A 3-year-old child presented with fever and cough and inability to feed for 2 days. On examination, chest in drawing is noted with respiratory rate of 42/minute. What is the severity grading of respiratory infection in this child?

A. No pneumonia
B. Pneumonia
C. Severe pneumonia
D. Very severe pneumonia

99. A 5-month child presented with difficulty in breathing, chest in drawing and wheeze of 3 day duration. Previous episodes of wheezing/ MDI usage are present. What is the next step in management of this condition?

A. Oxygen inhalation + Salbutamol nebulization
B. Oxygen inhalation + Inj. hydrocortisone
C. Oxygen inhalation only
D. Adrenaline nebulization

www.medicalcoupons.in

100. A 11-year-old child is found to be anemic with previous history of multiple blood transfusions. Lab values reveal:

- **Hb - 8 g/dL**
- **MCV - 70 fL**
- **MCH, MCHC - normal**
- **Blood smear - microcytic, hypochromic RBCs**

Which of the following investigation should be done next?

A. Serum iron
B. Hb electrophoresis
C. Direct Comb's test
D. Osmotic fragility

101. In a neonate who presented with incessant cry and irritability, cyanosis was noted which subsides when the baby starts crying. What should be done next?

A. Endotracheal intubation
B. Oropharyngeal airway insertion
C. Oxygen inhalation
D. Nasal CPAP

102. In an adolescent patient presenting with short duration illness and starvation, lab evaluation reveals unconjugated hyperbilirubinemia with normal liver enzymes. Which condition should be suspected?

A. Chronic pancreatitis
B. Acute viral hepatitis
C. Gilbert syndrome
D. Rotor syndrome

103. In a child presenting with short stature, GH levels were found to be high but IGF-1 levels are low. Likely abnormality?

a) GHRH deficiency
b) GHRH receptor defect
c) GH receptor defect
d) Panhypopituitarism

www.medicalcoupons.in

104. A 10-year-old child presents with rash in lower limbs with complaints of abdominal and joint pain. Lab findings reveal hematuria with normal platelet count. Likely diagnosis?

A. HSP
B. IgA nephropathy
C. ITP
D. PSGN

105. In a child presenting with febrile seizures, first line drug to control seizures?

A. Midazolam
B. Phenytoin
C. Valproate
D. Phenobarbitone

106. Which of the following is not true about colostrum?

A. Contains IgA
B. Lemon yellow color
C. Rich in minerals and amino acid
D. High amounts of sugar and fat

107. Dystrophin protein is absent in?

A. Becker muscular dystrophy
B. Duchenne muscular dystrophy
C. Emery Dreifuss muscle dystrophy
D. Limb girdle muscle dystrophy

108. A Baby presented with fever for 2days and has slapped cheek appearance, what is the diagnosis?



www.medicalcoupons.in

A. Erythema Infectiosum
B. Erythema Subitum
C. Erythema Marginosum
D. Dermatitis


109. A 25-year-old female who is a STD patient came with chief complaint of vaginal discharge with lower abdominal pain and tenderness, which colour kit will you prescribe to her

A. Grey kit (kit 1)
B. Red kit (kit 5)
C. Yellow kit (kit 6)
D. Black kit (kit 7)

110. A patient came to OPD with complaints of diarrhea, dermatitis and mental retardation. It was also found that he was a maize eater. Which of the following could be most probably diagnosis?

A. Beri Beri
B. Night blindness
C. Pellagra
D. Kwashiorkor

111. All are seen in this condition except?




A. Munro micro abscess/ Kogoj
B. Wick hams striae
C. Saw tooth acanthosis
E. Civatte bodies /shiny plaques

www.medicalcoupons.in

112. Patient on treatment with 13-CIS RETINOIC ACID. Which of the following is one of side effects of the drug?

A. Teratogenic
B. Neural tube defect
C. Follicular hyper keratosis
D. Photo onycholysis

113. A patient presented with hypo pigmented macula over back and upper trunk, on KOH mount spaghetti and meat ball appearance was seen. What can be the causative agent?



A. Malassezia fur fur
B. Tinea nigra
C. Trichosporon
D. Candida

114. Patient presented with lesions as shown in the image, later patient also developed arthritis. Treatment?



www.medicalcoupons.in

A. Methotrexate
B. Cyclophosphamide
C. Mycophenolate mofetil
D. Rituximab

115. Identify the plant shown in the image



A. Parthenium
B. Ginkgo biloba
C. Poison ivy
D. Vita trifolia

116. On which princip[illegible] th[illegible] following device functions?



www.medicalcoupons.in

A. Bohr effect
B. Bernoulli effect
C. Venturi effect
D. Boyles principle

117. Which of the following drug combinations contains all IV hypnotic agents?

A. Propofol, Thiopentone sodium, Etomidate
B. Halothane, sevoflurane, desflurane
C. Sugammadex, Sulfentanil, tramadol
D. Vecuronium, Rocuronium, Atracurium

118. Intravenous anesthetic agent of choice for cardiac patient

A. Propofol
B. Ketamine
C. Etomidate
D. Sevoflurane

119. What is the following Image referring to?



A. Mallampatti Score
B. Apgar Score
C. Duke's Criteria
D. Jone's Criteria

www.medicalcoupons.in

120. Level of ending of spinal cord in infants?

A. L1
B. L3
C. S1
D. S3

121. A patient suffered from Blunt trauma to the abdomen and was brought to the Emergency room. He is unstable having Blood pressure of 90/60 mm of Hg. You are the doctor, basic resuscitation is done with intravenous fluids, FAST is positive. He is not improving despite Intravenous fluids. Next step in the management is?

A. CECT
B. Exploratory Laparotomy
C. Laparoscopy
D. MRI

122. A female patient after resection of lipoma on the neck, develop elevated scar with itching and burning sensation. And scar was contained within the wound margin. What is the diagnosis?

A. Keloid
B. Hypertrophic scar
C. Suture granuloma
D. Infection

123. A pregnant lady with 7th week of POG with right lower abdominal quadrant pain. She got diagnosed with acute appendicitis. What is the best management in her case?


A. Immediate Laparoscopic appendectomy
B. Wait till 2nd trimester
C. Ochsner sherren regimen
D. Abortion

124. An alcoholic patient who is diagnosed with chronic hepatitis has been admitted to emergency room for vomiting of blood since last 2 days due to variceal bleeding. Patient is in shock. You are the doctor and you have started the first line treatment, but the patient died. What is the underlying cause of death in this patient?

www.medicalcoupons.in

A. Hepatic failure
B. Variceal bleeding
C. Chronic Hepatitis
D. Hypovolemic Shock

125. A patient was brought to the emergency with hypotension, raised JVP, heart sound was not properly audible, neck veins were engorged. He died in the hospital and autopsy was done which showed the following image. Most likely cause of death in this patient would be?



A. Fat necrosis
B. Cardiac tamponade
C. Perforation of intestine
D. Pneumonia

126. Lucid Interval is seen in?

A. Extradural hematoma
B. Subdural hematoma
C. Subarachnoid hemorrhage
D. Scalp hematoma

www.medicalcoupons.in

127. Battle sign is seen in:



A. Bruising on mastoids
B. CSF rhinorrhea
C. Periorbital staining
D. Frontal Bone abscess

128. A patient was brought to the emergency following a car accident. His Blood Pressure is 90/60 mm of Hg. His Glasgow coma scale is 9. He is having visceral injuries. Least likely cause of hypovolemia in this patient is?

A. Splenic injury
B. Sub dural hematoma
C. Retroperitoneal hematoma
D. Hemothorax

129. A patient had a central line insertion in the Subclavian vein and the patient collapsed during insertion. On examination, breath sounds suddenly were found to be absent on the side of insertion of central line, On Percussion hyper resonance sounds were heard on the same side of insertion of central line. Most likely cause in this patient?

A. Pneumothorax
B. Hemothorax
C. Tension pneumothorax
D. Neurogenic Shock

www.medicalcoupons.in

130. A Soldier was participating in a marathon in hot temperatures of 40 degrees Celsius, suddenly collapsed. What first aid should have been provided to him by the local people?

A. Immediately ice-cold water should be put on this soldier
B. Put foot of the soldier in ice immediately
C. Injection of Benzodiazepine
D. Wrap a cool water-soaked blanket over the person at around 25-degree Celsius temperature and switch on the fan.

131. A patient suffered from Femur neck fracture for which he was immobilized for a period of 3 months. Suddenly he developed breathlessness, cyanosis and was brought to the emergency. What can be the likely cause of this condition?

A. Fat Embolism
B. Venous thromboembolism
C. Systemic thromboembolism
D. Air Embolism

132. A lactating woman comes in the hospital with 3 days' history of painful, red, fluctuant, 2cm mass in the breast. USG shows fluid collection. What will you do in the management of this patient?

A. Antibiotics and Follow-up after 3 days
B. Incision and Drainage
C. Hot fomentation
D. Empty the breast every 3 hours

133. 32-year-old lady, 2cm single, mobile, asymptomatic breast lump with well-defined margin, noticed by the patient herself on self-breast examination, presented to the OPD. She was diagnosed as having a benign lesion. You are the doctor and what would you recommend to this patient

A. Reassure, continue to follow up the patient
B. Surgical Excision
C. Biopsy
D. Give Danazol

www.medicalcoupons.in

134. In a Type 3 Branchial fistula, internal opening is located in which part of pharynx?


A. Tonsil
B. Pyriform Sinus
C. Cricopharyngeus
D. External auditory canal

135. What should be the treatment for the pathology as shown in the given image:



A. Incision and drainage
B. Sclerotherapy
C. Radiotherapy
D. Excision

136. Following is _____ image of Polycystic Kidney disease:



www.medicalcoupons.in

A. PET-Scan
B. CT
C. MRI
D. USG

137. A 3-day old baby, is passing greenish yellow faecal discharge from the umbilicus. What shall be the Embryological cause of this condition?

A. Patent Allantois
B. Patent Vitello intestinal duct
C. Meckel's diverticulum
D. Patient umbilical end of omphalomesenteric duct

138. 5-year-old baby having constipation. We are giving stool softeners also but is still able to pass only one stool in one week. There is a history of passage of meconium after 48 hrs of birth, what investigation would you do initially in this baby?

A. Ask mother to give high fiber diet
B. Plain X-ray of abdomen
C. Barium enema and Anorectal manometry
D. Surgical Treatment

139. A patient had a laparoscopic cholecystectomy and Gall Bladder was removed and was discharged on the same day, he later came with severe abdominal pain and tenderness, LFT was normal. What is the diagnosis?

A. Bile Leak
B. Perforated peritonitis
C. Infection of the surgical site
D. Abscess

140. 5-year-old boy is having continuous dribbling of urine and also normal micturition Most likely cause?


A. Posterior urethral valves
B. Ureteric opening distal to external sphincter
C. Urinary tract infection
D. Vesicoureteric reflux

www.medicalcoupons.in

141. Mother while giving bath to her 7-day old baby noticed that the scrotum of the baby is empty, you are the doctor, what treatment you will offer in this baby?

A. Wait and watch for spontaneous resolution
B. Give GnRH for 1 month, if testis does not descend down on its own, then do surgery before 2 years
C. Give GnRH for - month, if testis does not descend down on its own, then do surgery before 5 years)
D. Do immediate Surgery.

142. Identify the site of foreign body as shown in the image?



A. Trachea
B. Esophagus
C. Mediastinum
D. Pleural cavity

143. A Patient had ileal resection. Which of the following deficiencies would be seen in this patient?

A. Vitamin B12
B. Vitamin A
C. Vitamin B1
D. Vitamin C

144. A known case of Rheumatic arthritis was taking NSAIDs for long period of time. He presented to the emergency with severe pain. His X-Ray was done which showed the following. Most likely cause of his presentation would be?

www.medicalcoupons.in



A. Pericardial effusion
B. Pneumothorax
C. Pleural effusion
D. Perforation of Peptic ulcer

145. A lady was incidentally diagnosed with a 0.5cm solid mass in her Gall bladder on USG which was non mobile, diagnosed as GB polyp. She is asymptomatic. Next step in the management?

A. Reassure the patient and follow up
B. Laparoscopic Cholecystectomy
C. Open Cholecystectomy
D. CECT abdomen

146. What do you expect in a lady after GB removal most commonly?

A. Weight loss
B. Steatorrhoea
C. High cholesterol
D. Hypoglycemia

147. Miss X, 35yrs old presented to the emergency with upper abdominal pain radiating to the back, On X-Ray Colon cut off sign is seen. Likely diagnosis?

A. Acute Cholecystitis
B. Acute appendicitis
C. Acute Pancreatitis
D. Chronic Pancreatitis

www.medicalcoupons.in

148. Whitaker test is performed for:

A. Posterior urethral valves
B. Hydronephrosis to compare abdominal and urinary bladder pressures
C. Ureterocele
D. Benign prostatic hyperplasia

149. A patient suffering from Inflammatory bowel disease presented with Peritonitis following ileal perforation. Exploration done. You as a doctor would prefer to do which of the following:

A. Ileostomy and definite surgery later
B. Meal pouch anal anastomosis
C. Resection of the diseased segment and anastomosis of the normal segment
D. Right hemicolectomy and ileo transverse anastomosis

150. A patient has presented with herniation of intestinal loop which passes along the spermatic cord & reaches the scrotum/cremaster muscle?

A. Femoral hernia
B. Direct inguinal hernia
C. Indirect inguinal hernia
D. Paraumbilical hernia

151. 3 Testicle appearance is seen in?

A. Vaginal hydrocele
B. Spermatocele
C. Epididymo orchitis
D. Torsion Testis

152. An old age patient felt heaviness in the left scrotum, he was suspecting cancer, came to ER. There was no pain or tenderness and transillumination was negative, most likely diagnosis in this patient would be:

A. Varicocele
B. Hydrocele
C. Testicular Torsion
D. Epididymo orchitis

www.medicalcoupons.in

153. The internal opening in Anal fistula usually begins at?

A. Crypt at dentate line
B. Perineal body
C. Anal verge
D. Anorectal junction

154. Pressure to cause the pressure sore is:

A. 10mm of Hg
B. 20mm of Hg
C. 25mm of Hg
D. 30mm of Hg

155. 65-year male presented with a mass in the left lumbar region which was ballotable. He was having abdominal discomfort and on investigations he was found to have a 7 X 5 X 3cm which was confined within the kidney and diagnosed as Renal cell carcinoma. What would be the treatment in this patient?


A. Partial nephrectomy
B. Radical nephrectomy
C. Radiotherapy
D. Chemotherapy

156. A patient was operated for Benign prostatic hyperplasia. On evaluation, incidentally cancer of prostate was detected and on further work up was found to have metastasis. Which of the following drug is used?

A. Goserelin
B. Adriamycin
C. Cisplatin
D. BCG

157. Young male presented with scrotal enlargement. On investigations, LDH was significantly high but the HCG and Alpha fetoprotein were with in normal limits. Histopathology as shown. Most likely diagnosis in this patient.

www.medicalcoupons.in



A. Teratoma of testis
B. Seminoma of testis
C. Choriocarcinoma
D. Yolk sac tumor

158. A young male presented with enlargement of the scrotum along with lump in the paraumbilical region and swelling in the supraclavicular fossa. Most likely diagnosis?

A. Testicular tumour
B. Hydrocele
C. Testicular Torsion
D. Epididymo orchitis

159. Best prognosis is of which of the following Breast cancer:

A. Luminal A
B. Luminal B
C. Triple Negative Breast cancer
D. Her 2 neu enriched

160. A 7-year-old child with recurrent H/O jaundice comes to OPD. Lab tests detected that UDG is decreased. Diagnosis could be:


A. Gilbert Syndrome
B. Neonatal jaundice
C. Crigler-Najjar syndrome
D. Down syndrome

www.medicalcoupons.in

161. Role of Sunlight in vitamin D3 synthesis?

A. 7-Dehydrocholesterol to Cholecalciferol
B. Cholecalciferol to 25-Hydroxycholecalciferol
C. 25-Hydroxycholecalciferol to 1,25-Dihydroxycholecalciferol
D. Cholecalciferol to 1,25-Dihydroxycholecalciferol

162. Patient had gastrectomy 10 months ago has anaemia. While evaluating for cataract, she presents with the image shown below. Cause is due to deficiency of;



A. Vit B 12
C. Niacin
D. Biotin
D. Vit. C

163. A patient presented with delayed wound healing, bleeding gums and blood pooling around the eyes

A. Vitamin C deficiency
B. Vitamin B3
C. Vitamin D
D. Vitamin B6

www.medicalcoupons.in

164. Fat sugar baby appearance seen in

A. Kwashiorkar
B. Scurvy
C. Marasmas
D. Osteopetrosis

165. Deficiency of enzyme glucose-6-phosphatase

A. Glucose accumulation in liver
B. glucose accumulation in skeletal muscle
C. Pyruvate accumulation in liver
D. Gycogen accumulation in liver

166. Menke's disease is due to disorder of metabolism in

A. Copper
B. Zinc
C. Selenium
D. Iron

167. According to MTP act 2021 guidelines, MTP can be done up to _____ weeks?
A. 20
B. 26
C. 24
D. 28

168. A patient presents with 45 days of amenorrhea. She wants abortion, if you are the treating doctor, what would you do?

A. Mifepristone 200mg and methotrexate after 48 hours
B. Misoprostol 200mg and methotrexate after 48 hours
C. Mifepristone 200mg and misoprostol 400microgram after 48 hours
D. Methotrexate and misoprostol after 48 hours

www.medicalcoupons.in

169. Blood testis barrier is made up of:

A. Sertoli cells
B. Leydig cell
C. Interstitium
D. None of the above

170. 31 years old g3p2 woman at 34 weeks gestation presents with nausea &"not feeling well". Bp is 110/82. Fetal HR pattern is category 1. Hb level is 9g/dl, leukocyte count is 8000 /cc ml & platelet count is 84000 /cu mm, alt of 500 IU/l, AST 550 IU/l & bilirubin of 2.5mg/dl. Which of the following is the most likely diagnosis?

A. Acute leukemia
B. Hellp syndrome
C. Hepatitis infection
D. Pre-eclampsia with severe features

171. Breast feeding contraindicatio ns'

A. CMV
B. Herpes
C. HIV
D. Galactosemia

172. Lady comes for termination of pregnancy at 8 weeks. What measurement of Karman cannula is used?



A. 3 mm
B. 7 mm
C. 4 mm
D. 6 mm

www.medicalcoupons.in

173. Patient was advised to take pregnancy test regularly for:

A. ABO incompatibility
B. Fetal viability
C. Molar pregnancy
D. Sex determination of fetus

174. Chromo perturbation is done by:

A. Saline
B. Methylene blue dye
C. CO2
D. Phenolphthalein


175. A Pregnant woman complaints of urinary leakage on coughing and Laughing. Which of the following can cause this?

A. Stress Incontinence
B. Urge Incontinence
C. Overflow incontinence
D. Neurogenic Incontinence

176. Hormone responsible for pain during breast feeding is

A. Estrogen
B. Prolactin
C. Progesterone
D. Oxytocin

177. 15-year old girl comes with primary amenorrhea with well developed breast and pubic hair. Local examination of the external genitalia shows the following finding. Most probable diagnosis is



www.medicalcoupons.in

A. MRKH Syndrome
B. Testicular Feminization Syndrome
C. Imperforate Hymen
D. Aschermann's Syndrome

178. Amniotic fluid level is highest during

A. 28 weeks
B. 32 weeks
C. 36 weeks
D. 38 weeks

179. A 26-year-old primigravida with 1[illegible] weeks POG attends ANC OPD with history of lower quadrant p[illegible]in [illegible]nd fever. USG reveals10mm thickness of appendix and WBC =1[illegible]000 cells/cumm. Which of the following is done?

A. CECT as a next measure
B. Antibiotics and appendectom[illegible]y after 20 weeks
C. Antibiotics and appendectomy during delivery
D. Antibiotics and immediate surgery

180. A lady treated with O1 drugs for infertility comes with abdominal pain, loss of appetite & difficulty in breathing clinical examination reveals tense ascites. Investigations reveals large hydrothorax with haematocrit of 60%. USG shows the following finding. Diagnosis is



www.medicalcoupons.in

A. Ectopic pregnancy
B. Molar pregnancy
C. **Ovarian Hyper Stimulation Syndrome**
D. Endometriosis

181. 36-year-old comes with 10 weeks amenorrhea with severe vomiting and persistent spotting P/V. O/E She is pale and uterus is enlarged to 16 weeks size. Ultrasound examination shows the below picture. What is the most probable diagnosis?



A. Hydatidiform mole
B. Adenomyosis
C. Leiomyoma
D. Endometriosis

182. Ideal dose of folic acid for women with previous H/O NTD is
A. 1mg
B. 0.4 mg
C. 4 mg
D. 0.5 mg

183. After delivery cardiac output returns to normal after how many weeks

A. 7 days
B. 14 days
C. 45 days
D. 60 days

www.medicalcoupons.in

184. 25-year-old sexually active woman has come with complaints of profuse dirty white discharge per vaginum. However, there is no significant associated itching. Microscopic examination revealed the presence of clue cells. Ideal treatment for the above condition is

A. Metronidazole
B. Fluconazole
C. Azithromycin
D. Doxycycline

185. Filter used in colposcopy is

A. Red filter
B. Blue filter
C. Green filter
D. Yellow filter

186. Safest drug for hypertension in pregnancy is

A. Methyl dopa
B. Labetalol
C. Nifedipine
D. Enalapril

187. Which of the following is an indication for surgical management in ectopic pregnancy?

A. Sac size 2.5cm
B. beta HCG 30001U/L
C. Tubal ectopic
D. Presence of Fetal cardiac activity


www.medicalcoupons.in

188. Diagnose the condition based on the given x-ray:



A. Anterior dislocation of hip
B. Posterior dislocation of hip
C. Fracture neck of femur
D. **Perthes disease**

189. Diagnose the pathology in the given on the barium image ?



A. Achalasia cardia
B. Diffuse Esophageal spasm
C. Esophageal Polyps
D. Esophageal carcinoma

190. An RTA patient suffered Blunt trauma to the abdomen and was brought to the Emergency room, where you are the doctor He is unstable having Blood pressure of 90/60 mm of Hg. Basic resuscitation is done with intravenous fluids and FAST Performed is positive.

www.medicalcoupons.in

He remains unstable despite intravenous fluids. What would be the next step in the management of this patient?

A. CECT
B. Exploratory Laparotomy
C. Laparoscopy
D. Ultrasound

191. A 20-year old male presented with pain and swelling around wrist joint. There is no history of injury. X-ray taken from the patient is shown below, what is the likely diagnosis




A. Osteosarcoma
B. Rickets
C. Colle's Fracture
D. Giant cell tumour

192. Popcorn calcification in Chest X-ray seen in:
A. Pulmonary Hamartoma
B. Fibroadenoma
C. Bronchitis
D. TB

www.medicalcoupons.in

193. Identify the investigation given below



A. Barium meal follow through
B. Enterography
C. **CT Enteroclysis**
D. Mr Enteroclysis

194. A chest x-ray from a patient of covid-19 would show

A. Ground Glass opacification
B. Halo around nodules
C. Apical cavitation
D. Calcification

195. Cannon-Ball appearance is seen in



www.medicalcoupons.in

A. Sarcoidosis
B. Metastasis
C. TB
D. Pulmonary fibrosis

196. A 60-year-old man presents with bony pain. His calcium level is 14 mg/Dl. As per given image, what is the diagnosis?



A. Histiocytosis
B. Paget's disease
C. Hperparathyrodism
D. Multiple myeloma

197. Tall T wave, wide QRS interval, increased PR interval. Diagnose the case as per ECG findings:

A. Hyperkalemia
B. Hypokalemia
C. Hyponatremia
D. Hypernatremia

198. A patient with temporal lobe seizure, 90% lymphocytes, 10% polymorphic cells, 90 mg% protein in CSF examination. Temporal lobe EEG spike shown

A. Start IV Acyclovir
B. Start ATT
C. Start Ceftriaxone
D. Start ceftriaxone + Vancomycin

www.medicalcoupons.in


199. Cause of Mucormycosis is most commonly associated with;

A. Diabetic ketoacidosis
B. Steroids
C. Bacterial Infection
D. Diabetic Insipidus

200. Patient complaints of weakness, proptosis and his condition worsen by the end of the day.


A. Myasthenia Gravis
B. Lambert Eaton Syndrome
C. Alzheimer's
D. Reactive Arthritis

201. Treatment of the Diagnosis of following ECG:



A. Urgent defibrillator
B. Iv Adrenaline
C. CPR
D. Amiodarone

202. Diagnosis of the following ECG



www.medicalcoupons.in

A. Sinus arrythmia
B. WPW
C. Atrial flutter
D. Atrial fibrillation

203. A Patient came to the hospital with a H/O of persistent or worsening cough, hemoptysis, chest pain that gets worse when you take a deep breath, laugh, or cough, shortness of breath, frequent or persistent attacks of bronchitis or pneumonia and weight loss, what will the diagnosis?

A. TB
B. Bronchogenic carcinoma
C. Miliary TB
D. Pulmonary TB

204. Patient has renal tubular acidosis. Treatment:

A. Infusion of sodium bicarbonate
B. Infusion of NS
C. Infusion of Spironolactone
D. Blood Transfusion

205. Fixed wide split S2 heart sounds seen in:

A. TOF
B. Bicuspid aortic valve
C. ASD
D. TAPVC

206. Wheat eater patient came to the Hospital with the complaints of diarrhoea, Abdominal pain, heart burn, Indigestion, Nausea, Vomiting, Flatulence. Lab Diagnosis showed Anti TTG Antibodies: What could be the diagnosis:

A. Celiac Sprue
B. Pseudomembranous Colitis
C. Whipple's Disease
D. Crohn's Disease

www.medicalcoupons.in

207. A Patient 26 years old, came to the hospital with the complaints of fever malaise and head ache. Doctor observed the Prescence of Malar rash, and lab diagnostics showed Carditis, and Immunological tests revealed the presence of Anti Sm- Antibodies:

A. SLE
B. Rheumatoid Arthritis
C. Reactive Arthritis
D. AIDS

208. A 23-year-old male with high BP, profuse sweating and headache. What is the initial investigation?

A. 24 Hr urinary metanephrines
B. VMA
C. MIBG scan
D. CT scan

209. MC type of renal stone:

A. Calcium oxalate
B. Urate
C. Triple phosphate
D. Cysteine

210. Identify the following condition shown in the image?



A. Syndactyly toes
B. Polydactyly
C. Brachydactyly
D. Ectrodactyly

www.medicalcoupons.in


211. Which of the following is not an obstructive lung disease?

A. COPD
B. Asthma
C. Interstitial Lung Fibrosis
D. Bronchitis

212. Which of the following is not a component of Meigh s Syndrome?

A. Pleural effusion
B. Benign ovarian tumor
C. Ascites
D. Testicular tumor


213. A patient presented to ER is comp!aining that he is facing the worst headache in his life. What can b< the orol able cause?



A. Tens on headache
B. S ba achnoid hemorrhage
C. Ex adural hemorrhage
D. Intracerebral hemorrhage

www.medicalcoupons.in

214. What is the diagnosis as per given ECG?



A. Inferior wall MI
B. Posterior wall MI
C. Anterior wall MI
D. none of the above

215. Lymphogranuloma venerum causative agent:

A. Klebsiella
B. H. Ducrei
C. Chlamydia Trachomatis
D. T. Pallidum



Note: If granuloma inguinale is mentioned the answer will be Klebsiella

216. Tumbling motility is seen in-

A. Vibrio
B. Clostridium
C. Listeria
D. Salmonella

www.medicalcoupons.in

217. Banana shape gametocyte and ring form in Rbc's:





A. Ovale
B. Vivax
C. Falciparum
D. Malaria

218. A child present with multiple episode of bloody diarrhea. He was suspected to have E. coli O157 H7. Which of the following culture media can be used to identify?

A. TCBS
B. Sorbitol McConkey
C. Blood Agar
D. Chocolate Agar

219. Which hormone is produced by these cells?



A. Parathormone
B. Calcitonin
C. T3/T4
D. TSH

www.medicalcoupons.in

220. Which of the following helps in adhesion of leucocyte to the endothelium?

A. Selectin
B. F 13
C. F 9
D. C5a

221. A 7-year-old child has recurrent infection. He has past history of recurrent visits for infection. Thymus and parathyroid glands atrophy, facial defects are present, microdeletion in chromosome 22q11.2. What is the diagnosis?

A. DiGeorge syndrome
B. Down's syndrome
C. Klinefelter's syndrome
D. Turner's syndrome

222. Thermogenic Uncoupler is found in:
A. Mitochondria
B. Lysosome
C. Golgi bodies
D. Endoplasmic reticulum

223. 65-year-old patient with thrombocytopenia, neutropenia, eryhtrocytopenia. Splenomegaly seen. On bone marrow aspiration, there was increased level of blast cells. What is the chromosomal translocation?

A. t (8;14)
B. t (14;18)
C. t (11;17)
D. t (9;22)

224. SCID is due to deficiency of which enzyme?

A. Adenosine Deaminase
B. Collagen synthase
C. Glucocerebrosidase
D. homogentisate 1,2-dioxygenase

www.medicalcoupons.in

225. A man presented with red rashes after bee sting. Reaction is due to;

A. IgA
B. IgE
C. IgM
D. IgG

226. A case of Rh Incompatibility. Which type of hypersensitivity reaction is this?

A. Antigen-antibodies.
B. Cytotoxic reaction
C. Immune complexes.
D. Cellular response.

227. Normal level of ALP is seen in:

A. Osteoporosis
B. Paget's disease
C. Multiple myeloma
D. Vit K deficiency

228. A 55-year-old male presented to OPD with fever, weight loss, fatigue and pallor.
CBC shows:


- **Hb - 9gm/dl**
- **TLC - 1,00,000/cmm**
- **Basophil - 4%**
- **Platelet count 1.60,000/cmm**
- **Myeloid blasts <11%**

Diagnosis?

A. HCL
B. CLL
C. CML
D. NHL

www.medicalcoupons.in

229. What changes occur during left shift of the curve?



A. Increase Temperature, Decrease 2,3 DPG, Alkalosis
B. Decrease Temperature, Increase 2,3 DPG, Acidosis
C. Decrease Temperature, Decrease 2,3 DPG, Alkalosis
D. Increase Temperature, Increase 2,3 DPG, Acidosis

230. Pancytopenia with hepatosplenomegaly, blast cells in the bone marrow in a 23-year-old female.

A. Aplastic anemia
B. AML
C. CML
D. ALL

231. A post-menopausal patient has cervical cancer. She underwent radiotherapy. Which of the following is most radiosensitive organ?


A. Ovary
B. Vagina
C. Urethra
D. Bladder

232. A patient is diagnosed with tumor in abdomen and got operated. On histopathological examination, tumor was found to be originated from all three germ layers. What is the diagnosis?

www.medicalcoupons.in

A. Teratoma
B. Adenoma
C. Mixed
D. Choriostoma

233. A female is on a Vegan diet, her peripheral blood smear is given below. What is the diagnosis



A. Vitamin B12 deficiency
B. Folic acid deficiency
C. Iron deficiency anemia
D. Sickle cell anemia

234. What findings would be seen on Chest Xray of a patient of silicosis who is having respiratory difficulty?

A. Honey comb appearance
B. Starry sky appearance
C. Tram track appearance
D. Snow storm appearance

235. Which gene mutation is seen in hereditary hemochromatosis disease?

A. Ceruloplasmin
B. Transferrin receptor
C. ATP7B
D. HFE

www.medicalcoupons.in

236. A person 47 years old is on medication and that he is recovering from the symptoms but he is complaining of having hyper sexual desire. Which condition he is suffering from?

A. Nymphomania
B. Satyriasis
C. Fetishism
D. Frotteurism

237. A patient presented to the psychiatric OPD with complains of recurrent episodes of palpitations, sweating, Dyspnea and insomnia. He was very scared of death. He has a history of being on a ventilator when he was COVID 19 positive one year back. What could be the diagnosis?

A. PTSD
B. GAD
C. Panic attack
D. Dementia

238. A chronic alcoholic man, takes alcohol every day, presented with altered sensorium, dementia and hallucinations. Last intake history was 2 days ago. What is the diagnosis?

A. Korsakoff syndrome
B. Wernickes encephalopathy
C. Delerium tremens
D. Alcoholic intoxication

239. Identify the splint:



www.medicalcoupons.in

A. Dennis Brown Splint
B. Cock-Up Splint
C. Sugar tong splint.
D. Knuckle bender splint


240. Vocal cord palsy treatment??

⇒ **Treatment: wait and watch**

241. Condyloma accuminata caused by

⇒ **Human papilloma virus**

242. Which inflammatory cells is seen in this image?



⇒ **MACROPHAGES**

243. Which TB drug is contraindicated in pregnancy?

⇒ **Rifampicin**

244. Genetic syndrome associated with Tetralogy of Fallot?

⇒ **DOWN SYNDROME**

245. PSM program NIKSHAY for

⇒ **TB surveys**

246. Marasmus is due to less intake of

www.medicalcoupons.in

⇒ **PROTEIN**

247. Bartholin cyst located at?

⇒ **Vaginal gland**

248. Cholesterol is synthesis from oxaloacetate by

⇒ **ACETYL-COA**

249. Gram positive catalase negative, optochin resistant organisms endocarditis is

A. Streptococcus agalactiae
B. Streptococcus pneumoniae
C. Streptococcus sanguis
D. Streptococcus epidermidis

250. In the setting of poor resource and village setting, which of the following is a cost-effective method to diagnose syphilis?

A. Culture
B. Dark field microscopy
C. Light microscopy
D. Immunofluorescence

251. A 7-year-old child while playing fell down. He has developed nasal blockage on both sides and there is swelling around the nose. Further Rx?

A. Aspiration of septal swelling
B. Anterior nasal packing
C. Start intravenous steroids
D. Start topical corticosteroid nasal spray

252. A 65-year-old patient is complaining of long standing mucopurulent discharge from left eye and a swelling near medial canthus. Best Mx for this patient?

www.medicalcoupons.in

A. Dacryocystectomy
B. Syringing and probing
C. Dacryocystorhinostomy
D. Lacrimal massage

253. A patient with CSOM with positive fistula sign reported few days later with negative fistula sign. What is the expected tuning fork test finding in this patient?

A. False negative rinne test
B. False positive rinne test
C. True positive rinne test
D. True negative rinne test


254. True about little's area

A. Keisselbach's plexus is present in little's area
B. It is present on lateral wall of nose
C. It is present in the posterior part of nasal cavity
D. It is rare site of epistaxis

255. A couple brought their son to doctor. They were worried because their son likes to eat foods from multinational companies. What we will call this shift from consumption of traditional natural foods to western fast foods?

A. Food transition
B. Acculturation
C. Social pathology
D. Demographic change

256. Which is the following is an absolute contraindication to the below device?




www.medicalcoupons.in

A. Past H/O PID
B. Past H/O ectopic
C. Subserious fibroid
D. Unexplained vaginal bleeding

257. A baby born to hyperglycemic mother will have?

A. Hypercalcemia
B. Hypokalemia
C. Hypoglycemia
D. Macrosomia

258. MBBS student was asked to diagrammatically make axis deviation of a patient. The following schemata shown represents?



A. Right ventricular hypertrophy
B. Left ventricular hypertrophy
C. Biventricular hypertrophy
D. Normal axis

259. A known case of sarcoidosis presents with nausea and vomiting. Serum calcium level - 12.5 mg%. Which of the following is least useful for this patient?

A. Mineralocorticoids
B. Bisphosphonates
C. Glucocorticoids
D. Normal saline

www.medicalcoupons.in

260. CLD patient presented to OPD with fever and gross abdominal distention. On abdominal examination tenderness is elicited. Ascitic fluid reveals presence of 500 cells/ cu.mm. Which is the next best step?

A. Large volume paracentesis
B. IV albumin plus midodrine
C. IV Octreotide
D. IV ceftriaxone

261. A patient has presented with features of cardiac tamponade. Which is a true finding?

A. Pulsus paradoxsus is always present
B. Kussmaul sign is always present
C. Tall A wave is present
D. Jugular veins column is visibly distended without pulsations

262. A patient is having sudden exacerbation of dyspnea and can't talk. On chest auscultation prolonged expiration is noted. ABG report shows ph= 7.3 PC02= 68 mm Hg. HCO3= 3 meq/L. Which of the following intervention should be done in this child?

A. Intravenous soda bicarbonate
B. Ammonium chloride
C. NI-PAP
D. Fluid resuscitation

263. A 45-year-old man with k/c/o emphysema and 20 years' history of working in factory presented with daily symptoms of chest tightedness. which factory is he likely to have been working in?

A. Textile
B. Shipyard
C. Glass
D. Cardboard

www.medicalcoupons.in

264. Investigations for Pulmonary embolism?

A. CTPA (CT- Pulmonary angiography)
B. CECT chest
C. Doppler
D. USG

265. Patient came in emergency room with breathlessness. On lab examination, hypokalemia causes T waves on the ECG. Potassium levels are critically low (< 1.7). This condition can lead to?

A. Torsades de pointes
B. SVT
C. Atrial fibrillation
D. Multifocal atrial tachycardia

266. A patient presented with polyuria and with no presence of sugar in urine, what could be the diagnosis?

A. SIADH
B. Diabetes insipidus
C. Type 2 DM
D. Type 1 DM

267. Mechanism of action of doxycycline is


A. DNA gyrase inhibitor
B. Cell wall synthesis inhibitor
C. Protein synthesis inhibitor
D. DNA dependent RNA polymerase inhibitor

268. Since last 3 months a non-alcoholic patient started believing that his daughter is trying to kill him. what is the diagnosis?

A. Schizophrenia
B. Bipolar disorder
C. Delusional disorder
D. Xenophobia

www.medicalcoupons.in

269. Earliest inflammatory cell out of the given cells in the image is



A. Lymphocyte
B. Neutrophil
C. Fibroblast
D. Macrophage

270. Which of the following activities are low in riboflavin deficiency?

A. Transketolase
B. Glycogen phosphorylase
C. Propionyl CoA carboxylase
D. Glutathione reductase

271. Enzyme deficit in galactosemia?

A. Galactose – 1 – phosphate uridyltransferase (GALT)
B. ADP-ribosyl transferase
C. Glycosyl-transferase
D. TDP daunosamine

272. What happens after a meal?

A. Increase in insulin and decrease in glucagon
B. Increase in glucagon and increase in insulin
C. No change in both
D. Decrease in insulin and glucagon

273. A Child Presented with Low BP and Had Clitoromegaly. Which Enzyme Deficiency Is Suspected Here?

www.medicalcoupons.in

A. 21-β Hydroxylase deficiency
B. 11- β Hydroxylase deficiency
C. 3- β Hydroxylase dehydrogenase deficiency
D. 17-α Hydroxylase deficiency

274. A Vitamin A dose > 12-month old baby weight 8kg?

A. 50000 IU
B. 100000 IU
C. 2 lakh IU
D. 150000 IU

Kindly email us on arisemedicalacademy@gmail.com to help improve the recall and to submit the questions missed out, your feedback, suggestions and support is highly appreciable

www.medicalcoupons.in